# CHARLES HERBERT BEST

CBE, M.A., MD, DSC. (LOND) FRS, FRCP (CANADA)
Professor and Head of Department of Physiology, Director of the BantingBest Department of Medical Research University of Toronto

### NORMAN BURKE TAYLOR

V D., M D., T R S. (CANADA), F R C S. (SENY) F R C P. (CANADA), M R C S. (SEW) L R C P. (LOND)
Formerly Professor of Physiology, University of Toronto, and Professor of the
History of Medicine and Medical Literature, University of Western Ontario,
London Canada

# The Liv<u>i</u>ng Body

# A TEXT IN HUMAN PHYSIOLOGY

Fourth Edition Reprinted



ASIA PUBLISHING HOUSE

BOMBAY CALCUTTA NEW DELHI . MADRAS

## PREFACE TO THE THIRD EDITION

THE DYNAMIC nature of physiological study, though burdensome to those who must revise its textbooks, is one of its chief fascinations The subject is never static but forever changing, always flowing and ricestless with fresh discoveries and new conceptions which sweep away without regret many old ideas and theories. The rapid advances made by physiology and the contributory sciences since the last revision of this book have made necessary many changes and additions. Two new Chapters (1 and 5) have been written. Chapter 1 has been devoted to "Elementary Physical and Chemical Principles," and should be a help to those readers who have had no instruction in the basic sciences; the to those readers who have not all of instruction in the oaste setences, the use of radioactive isotopes in physiological research, and in the treatment of disease has been lightly touched on. In Chapter 5, entitled "The Defensive Mechanisms of the Body against Disease and Injury," the processes of immunity and related questions, the use of the "sulfa" drugs and antibiotics, as well as the problems of shock, anaphylaxis, drugs and antibiotics, as well as the problems of snock, anaphyaxis, and allergy, have been outlined. Chapter 14, dealing with "The Physiology of Reproduction," has been enlarged in scope by the addition of several new sections and the expansion of some of the old ones. All this new material together with that required to bring the text up to date has led inevitably to an increase in the size of the book. But the increased size is due also, and in large measure, to new illustrations, the total number of figures having been raised from 283 in the last edition to 400 in the present one.

In writing upon a subject such as that of mammalian physiology, which is so largely a study of interactions and integrations of processes

and systems, it is not always possible to avoid some anticipation of the reader's knowledge. An endeavor has been made to offset this tendency by providing many cross-references in the text. But this provision may sometimes be found inadequate, and readers are advised to make free use of the index which is rather comprehensive.

Much of the rewriting in this edition has been undertaken to meet the suggestions and kindly criticisms of many colleagues, to whom sincere thanks are extended More than a few of the changes which are believed to have improved the book are due to suggestions made by Professor J B Stier of the University of Indiana. His interest and care in reading the second edition from cover to cover, and his comments from the view of an American teacher are gratefully acknowledged.

Recognition of the painstaking care with which Mrs. Dorothy Spicer has given her secretarial assistance in preparing the manuscript for the press is also warmly expressed.

N. B. T.

21 Ardwold Gate Toronto March, 1952

### 1. Elementary Physical and Chemical Principles

### 2. Protoplasm. The Cell. The Tissues

The Physiological Properties of Protoplasm The Cell The Tie sues

# The Blood, Lymph, and Tissue Fluids; the Spleen and the Reticulo-Endothelial System

Composition of the Blood The Red Blood Cells or Erythrocyte. The Volume of the Blood Hemoglobin The Formation and History of the Red Cells Anemia Hemorrhage Transfusion History of the Laking of Blood The White Cells or Leucocyte. The Blood Platelets or Thrombocytes The Retuculo-Endothelis System The Spleen The Lymph and the Lymphatic System The Coagulation of Blood and Lymph Hemorrhagic States Due t Failure of the Clotting Mechanism. The Regulation of the Read tion of the Blood and Other Body Fluids

### 4. The Circulation of the Blood

General Description of the Circulatory System Hemodynam Principles The Velocity of the Blood The Arterial Blood Pre sure Capillary and Venous Pressures The Effect of Gravi upon the Circulation The Effects of Acceleration upon the Ci culation The Arterial Pulse The Structure and Physiologic

### CONTENTS Properties of Cardiac Muscle. Perfusion of the Excised Heart,

The Origin and Conduction of the Heartbeat. The Action of the Heart, the Cardiac Cycle. The Opening and Closing of the Heart Valves. The Regulation of the Heart's Action. Nerve Control of the Blood Vessels Special Features of the Circulation in Certain Parts The Defensive Mechanisms of the Body Against Disease and In-

210

225

268

288

350

403

jery; Shock, Anaphylaxis, Allergy Shock ~ 6. Respiration An-Outline of the Structure of the Respiratory Organs. The Me-

schanics of Respiration. The Subdivisions of the Lung Air The Chemistry of Respiration The Control of Respiration Physiological Modifications of Respiration Abnormal Types of Respiration Oxygen Want-Anoxia or Hypoxia The Voice 7. The Physiology of the Kidney. Micturition

Micturition or the Voiding of Urine 2. Digestion Enzymes or Ferments and Their Action Digestion in the Mouth.

Digestion in the Stomach. Digestion in the Intestine The Production of Bile by the Liver: The Composition of Bile, Its Storage in, and Expulsion from, the Gall Bladder Absorption from the In-

testinal Tract

9. Metabolism and Nutrition

General Metabolism. The Metabolism of Protein The History of Protein in the Body. Purine Metabolism. Carbohydrate Metabo-

lism Fat Metabolism. The Regulation of Body Temperature. The

Vitamins, The Principles of Dietetics

10. The Endocrine Glands

The Thyroid Gland, The Parathyroid Glands The Adrenal Glands (or Suprarenal Capsules). The Pituitary Body or Gland

(Hypophysis Cerebri). The Sex Glands or Gonads. The Thymus. The Pineal Body

CONTENTS	xi
11. The Physiology of Nerve and Muscle	459
The Nature of the Nerve Impulse. The Physiology of Skeletal (Voluntary) Muscle The Physiology of Muscular Exercise	
12. The Central Nervous System	506
Reflex Action. The Spinal Cord. The Brain or Encephalon. The Membranes (Meninges) of the Brain and Spinal Cord The Autonomic Nervous System. Conditioned Reflexes. The Physiology of Sleep	
13. The Special Senses	574
The Physiology of Vision. The Ear. Taste Smell Cutaneous Sensations	
14. The Physiology of Reproduction L	648
The Female Reproductive Organs The Male Reproductive Organs The Processes of Reproduction	
Glossary	681

707 723 <sub>.</sub>

Appendix Index

# Elementary Physical and Chemical Principles'

The elements Substances such as pure iron, copper, gold, silver, so dium, potassium, chlorine iodine, oxygen hydrogen, and a great number of others which are composed entirely of but one type of particle or atom are called elements. A list of the more common elements is given in Table 1 1 (p 2) Those composing the animal body are shown in percentages in Figure 1 1 (p 3)

The conception of an indivisible particle of matter—the atom (lit erally that which cannot be cut or divided)—goes back to ancient Grecian times (Democritus, 600 B c), but its modern history starts with John Dalton (1766–1844), an English chemist

Nevertheless, the atom is not, as was believed for so long, the ultimate particle of matter, it is composed of still more minute particles arranged within it according to a definite pattern. The simplest and lightest—the hydrogen atom—resembles a miniature solar system. It consists of a core or nucleus with a much smaller and lighter particle revolving around it (Fig. 1.2). All atoms of whatever kind, whether of coaygen. Chlorine, won. gold. silver, or any other dement. The constructed upon an essentially similar pattern, but the particles in the central core as well as the lighter revolving particles vary in number.

<sup>1</sup> This chapter has been written for those readers who have had no instruction in chemistry or physics. It may be omitted by others. It is offered merely as an intro duction to and an aid in the understanding of subsequent chapters. Several chemical and physical principles such as hydrogen ion concentration (p. 108) osmosis (p. 64) kinetic theory and diffusion (pp. 244 and 245) the composition of the principal foodstuffs (Chaps. 8 and 9) and the actions of enzymes (p. 285) are described more appropriately in chapters dealing with subjects to which such principles are most directly applicable.

Greek or

Latın Name

Aluminium

Arsenicum

Phosphoricum

Platinum

Kalum

Radium

Argentum

Strontium

Natrium

Sulfur

Stannum

Hranum

Zincum

Silicon

Subium

Barrum

D----

2

Common Name

of Element

Aluminium

Antimony

Phosphorus

Platinum

Radium

Silicon

Silver

Sodium

Sulfur

Tin

Zinc.

Strontium

Uranium

Potassium

Arsenic

Barium

D----

Table	1-1	

Symbol

Αl

Sh

As

Ba

n

Atomic

Weight

26 98

121 76

74 91

137 36

30 97

195 23

39 10

226.05

28 09

107 88

22 99

87 63

32 06

118 70

238 07

65 38

10.02

Atomic

Number

13

51

56

15

78

19

88

14

47

11

38

16

50

92

30

Boron	Boton	В	10 82	5
Bromine	Bromum	Br	79 92	35
Cadmium	Cadmium	Cd	112 41	48
Calcium	Calcium	Ca	40 08	20
Carbon	Carbon	C	12 01	6
Chlorine	Chlorum	Cì	35 46	17
Cobalt	Cobaltum	Co	58 94	27
Copper	Сиргит	Cu	63 54	29
Fluorine	Fluorum	F	19 00	9
Gold	Aurum	Αu	197 20	79
Helium	Helium	He	4 00	2
Hydrogen	Hydrogenium	H	1 00	1
Iodine	Iodum	I	126 91	53
Iron	Ferrum	Fe	55 85	26
Lead	Plumbum	Pb	207 21	82
Lithium	Lithium	Lı	6 94	3
Magnesium	Magnesium	Mg	24 32	12
Manganese	Manganesium	Mn	54 93	25
Mercury	Hydrargyrum	Hg	10 002	80
Neon	Neon	Ne	20 18	10
Nickel	Niccolum	Nı	58 69	28
Nitrogen	Nitrogenum	N	14 00	7
Oxygen	Oxygenium	0	16 00	8

P

Pr

ĸ

Ra

Sı

Ag

Νa

Sr

S

Sn

U

Zα

with the different kinds of atom. The central core or nucleus of the atom contains one or more particles called *protons*. Each proton holds a positive charge of electricity, and, except in the case of the hydrogen atom, is paired with an electrically neutral particle called a *neutron*. The hydrogen atom is a single proton, unpaired with a neutron. The particle or particles revolving in one or more orbits around the nucleus of the atom are electrically negative and are called *electrons*. The chemical properties of the atom are dependent upon the




Fig 1.1 Diagram to show the percentages of the elements composing the body.

electrons in the outer orbit. The number of protons, which as just mentioned varies with the kind of atom, is balanced by an equal number of electrons. Positive and negative charges being equal, the atom as a whole, so long as it loses none of its charges, is electrically neutral. The number of protons (or electrons) possessed by an atom is called the atomic number. The atomic numbers of the naturally occurring elements are from 1 (for hydrogen) to 92 for uranium. The atomic numbers for the other elements run in unbroken sequence between those two extremes.

Atomic weights. The atomic weight of an element is the weight of one of its atoms in relation to the weight of the hydrogen atom taken as 1.008. It is the weight of the smallest quantity of an element present in the molecule of a pure substance. It will be seen from Table 1-1 that the weights of the different kinds of naturally occurring atoms vary widely, radium (226 05) and uranium (238.07) being the heaviest in Table 1-1, but still heavier ones, such as plutonium, ameri-

### FLEMENTARY PHYSICAL & CHEMICAL PRINCIPLES

cium curium and mendelevium, etc., have been produced in the laboratory

Isotopes. Until recent years it was universally believed that elements were atomically homogeneous—that its atoms were all pre




Fig 12 Showing the structure of the atom A hydrogen atom B helium atom C carbon atom.

cisely of the same weight. Since it has become possible to determine the weights of individual atoms, it has been found that most of the elements are composed of atoms of slightly different weights. For example, chlorine, which when its atoms are taken in the aggregate has an atomic weight of 35.46, is constituted of atoms with weights of




Fig 1.3 The isotopes of chlorine. The figures refer to the numbers of protons and neutrons in the nucleus and to the electrons in the three orbits.

35 and 37, there being none having a weight of exactly 35 46. Atoms of the same element but of different atomic weights are called isotopes

The disparity of the atomic weights of isotopes is due to their having different numbers of neutrons. Since neutrons do not earry an electric charge, the nuclear charges (protons) of isotopes are the

Radioactive isotopes. Many isotopes are radioactive like radium and uranium. That is to say, their nuclei are unstable and particles are shot out at very high velocities (20,000 to 100,000 miles per second).

Fig 1.4 An alpha particle it will be noted that it is of the same constitution as the helium nucleus



The particles (see Fig 14), usually referred to as "rays," consist of a group of two protons and two neutrons held together as a unit (alpha particle or alpha "ray") and electrons (beta 'rays') A third type of ray with a high power of penetration is also emitted, rays of this type are known as gamma rays and are identical with or closely similar to hard x-rays

Isotopes of various elements are now produced artificially by bombarding atoms with protons or neutrons produced by an apparatus called a cyclotron, developed by Dr. Lawrence of California The high speed projectiles penetrate the nuclei of some of the bombarded atoms, a nuclear reaction follows, and an isotope of the bombarded element, or of a newly created element, is produced. The production in this way of radioactive isotopes has been a great boon to those carrying out researches into biochemical and physiological probleris. The radioactive isotope can be incorporated into a compound whose history in the body it is desired to follow. When such a compound is fed to an animal, it behaves chemically in the body as would the naturally occurring nonradioactive substance, but on account of its radioactivity it is "labeled" or "tagged" as it were, and can be detected in the body tissues or excreta by means of a Geiger counter, an instrument sensitive to the rays which the incorporated isotope emits

#### ELEMENTARY PHYSICAL & CHEMICAL PRINCIPLES

Radium has long been used in the treatment of cancer and other malignant growths, the emitted rays having a destructive action upon the cancer cells, but they also destroy normal issue and for this reason their action must be strictly localized to the diseased structure Artificially produced radioactive isotopes are now employed for their therapeutic effects, and more especially those which have a selective action upon certain tissues, since they can be injected into the general circulation. The thyroid gland, for example, has a great affinity for ordine, and a radioactive isotope (I<sup>128</sup>, I<sup>130</sup>, or I<sup>131</sup>) of this element concentrates its action upon thyroid tissue, it is, therefore, used in the treatment of certain types of goiter (Chap 10). Radioactive phosphorus is attracted to and exerts its effect upon osseous tissue and is of value in the treatment of some types of bone disease.

Chemical symbols. The chemist employs a kind of shorthand language for the expression of the elemental composition of chemical substances and to indicate the character of chemical reactions. In this chemical language, the elements are designated by the first letter of their Latin names, or, when the initial letter of the names of more than one element is the same, a second letter of the name is added Thus, F is the symbol for fluorine, but since there are a number of elements commencing with the letter C, the latter is used alone as the symbol for carbon, and copper (L cuprum), calcium chlorine, and cobalt are designated Cu, Ca, Cl, and Co, respectively Similarly, while H is the symbol for hydrogen, mercury (Hydrargyrum) is represented by Hg and helium by He (see Table 1-1)

A chemical formula indicates the atomic constitution of the molecules of a substance, the symbols of the elements being written in a certain conventional order (empirical or molecular formula) or according to a specific pritern which shows the spatial relationships of the atoms to one another in the molecule. The latter is called a graphic or structural formula.

An empirical formula indicates merely the number of the different kinds of atoms in the molecule. When there are more than one of the same kind of atom, a small figure is written below the symbol. Thus, sodium chloride is written NaCl, but water, which is formed by the union of two atoms of hydrogen and one of oxygen, is written H<sub>2</sub>O. The two types of formula for the amino acid glycine are shown below

C<sub>2</sub>H<sub>3</sub>NO<sub>2</sub> empirical formula CH<sub>2</sub>—NH<sub>2</sub>
COOH
graphic formula

If we compare the symbols of the elements to the alphabet of ordinary language then chemical formulas may be likened to words Formulas written with appropriate signs  $(+, -, -, \rightarrow)$  are known as equations, by this means, information is given with respect to the na ture of the reactions taking place between the molecules (see below) of chemical substances when brought together Thus, the formation of carbon dioxide by the action of hydrochloric acid on limestone (calcium carbonate) is expressed in the following equation

A molecule of calcium carbonate reacts with 2 molecules of hydro chloric acid to produce a molecule each of carbon dioxide, calcium chloride and water. It will be noted that the two sides of the equation balance—that is the number of calcium carbon, hydrogen, chlorine, and oxygen atoms on the two sides are equal.

Chemical reactions-molecules, Certain atoms have an attraction or affinity for other kinds of atom the two kinds uniting to form sub stances called compounds which have properties quite different from those of the originally separate atoms. Common table salt, for example, which is known chemically as sodium chloride and has the formula NaCl, is formed by the union of one atom of sodium (Na) with one atom of chlorine (Cl) Sodium is a caustic solid element, highly destructive to animal and vegetable tissues chlorine is a green ish intensely irritating, poisonous gas, yet the compound, sodium chloride, which results from their union, is a mild harmless substance and, indeed, is indispensable for the well being of the animal body It is present in a concentration of about 0 6 percent in the body fluids The particles of a compound such as sodium chloride, carbon dioxide. water, etc., are called molecules Any pair or group of atoms which being united form a unit particle of a compound is so termed The molecules of the compounds just mentioned are among the smallest and simplest known, but some molecules such as those composing proteins, are relatively enormous, being constituted of hundreds of atoms of several different kinds The molecular weight (which is arrived at by adding together the weights of the constituent atoms) of these large particles is thousands of times greater than the weight of a water molecule The molecular weight of water (HO), for ex ample, is only 18 (hydrogen 2 + oxygen 16), whereas that of certain proteins is in the neighborhood of 400,000 or more. The hydrogen

molecule  $(H_2)$ , for it is not only atoms of different kinds which form molecules, is the smallest and lightest of all. The atoms of nitrogen or of oxygen also combine molecules to form  $(N_1, O_2, \text{ and } O_3)$ 

If one wished to show graphically the great differences in size be tween the smallest and the largest molecules, one might represent a water molecule by a dot, then one of the largest protein molecules would require for its representation a space larger than this page Molecules also vary in shape, some protein molecules, for example,




Fig. 1.5 Diagram showing the union of a sodium atom and a chlorine atom to form a molecule of sodium chloride. The figures indicate the number of protons and neutrons in the nuclei and of the electrons in the concentric orbits. The hatched circle represents the electron common to both types of atom in a molecule of sodium chloride.

such as serum albumin, are globular in form, while in others, such as muscle protein, the atoms are extended in chains to form long elon gated fiberlike molecules (see p 362)

As implied at the beginning of this section, it is only some atoms which are capable of uniting with one another to form compounds. The attraction or affinity of one atom for another depends upon the instability of the outer orbit of electrons. For example, the number of electrons in the outer orbit of the sodium atom and of the chlorine atom render their outer orbits unstable. The former has one electron too many the latter one too few, for stabilization. But there is always a tendency toward the stable state. This is achieved, when possible by the sodium atom losing an electron and the chlorine atom gaining one. So then, when a sodium atom and a chlorine atom are brought together an electron is transferred from the one. (Na.) to the other (CI). When sodium chloride undergoes ionization (see p. 12) the sodium atom has one less electron than it had originally, and the

chlorine atom one more. In the unionized state, the two atoms are held together, as it were, by the electron in question (Fig. 1.5).

Equivalent or combining weights. When the elements combine to form compounds, they do so in certain definite and unvarying proportions. The smallest amount in grams in which an element combines with 1 gram of hydrogen is called its equivalent or combining weight. An element also replaces 1 gram of hydrogen in a compound or combines with another element to form a compound by this same amount —its equivalent weight—or some simple multiple of it. The equiva-

### Table 1-2

	Equivalent or combining weights
	(number of grams which combine
Element	with or replace 1 gram of hydrogen)

Calcium	20
Carbon	3
Chlorine	35 46
Magnesium	12
Nitrogen	4.67
Oxygen	
Potassium	39 1

lent weights of a few of the commoner elements are given in Table 1-2. It will be seen by comparing Tables 1-1 and 1-2 that the atomic weight of an element is either equal to the combining weight or is a multiple of it. The equivalent weight of oxygen is 8, its atomic weight is  $(8 \times 2 -)$  16; that of carbon is 3, its atomic weight is  $(3 \times 4 -)$  12; the equivalent weight of nitrogen is 4.67, its atomic weight is  $(4.67 \times 3 -)$  14, but the equivalent weight of chlorine and potassium are the same as their atomic weights—namely, 35.46 and 39.1 respectively.

Oxidation and reduction. When oxygen unites with another element, the compound so formed is called an oxide, and the process is referred to as oxidation. If the reverse action takes place—namely, a compound loses oxygen—the substance is said to be reduced and the process is known as reduction. But the loss of hydrogen from a compound is also called oxidation, and the addution of hydrogen to a compound is called reduction.

A compound of an element with hydrogen and oxygen is called a hydroxide. Iron rusts when exposed to moisture and air, the rust being a hydroxide of iron (ferric hydroxide). Many metals and other sub-

stances form oxides or hydroxides when exposed, the oxygen being derived from the atmosphere. The carbon in substances such as wool, coal, etc., undergoes oxidation during combustion, carbon dioxide (CO) being produced and given up to the atmosphere. The carbon of our food also undergoes oxidation in the body, the carbon dioxide so formed being exhaled in the breath. But many substances in the tissues undergo oxidation, not by the addition of oxygen, but by the loss of hydrogen brought about through the action of specific dehydrogenating enzymes (dehydrogenases).

Besides the combination of atoms with the formation of compounds, which so far has been chiefly dealt with, two other types of chemical reaction remain to be described namely, the breakdown of molecules into smaller groups of atoms or into individual atoms—decomposition—and the disruption of the molecules of two or more compounds, followed by the union of the liberated atoms or groups of atoms in new combinations. This reaction is known as double decomposition.

The chemical reaction known as double decomposition can be illustrated by changes occurring when HCI is mixed with a solution of sodium bicarbonate. Upon the instant of mixing the two liquids, vio lent effervescence occurs, for the bicarbonate has been decomposed and carbon dioxide (a gas) is given up to the atmosphere. The hydrochloric acid also splits into its constituent atoms. Atoms of sodium unite with the chlorine atoms to form sodium chloride, carbon atoms with a molecule of oxygen to form carbon dioxide, and hydrogen molecules with oxygen atoms to form water, thus

The indestructible nature of the atom in ordinary reactions. Molecules disintegrate and fresh compounds are formed, but the atoms in the ordinary process of Nature are indestructible. Decay and disintegration of substances are occurring ceaselessly. Organic matter, animal and vegetable, is continually undergoing decomposition, its elements being returned to the earth and there reconstituted into new compounds. Even rocks become eroded and metals wear away, but the immortal atoms reappear in new forms. Though materials composing the world around us disappear and are lost to our senses, the sum total of matter undergoes no reduction in weight. When we watch a blazing log being consumed until nothing but the ash remains, we may doubt the general statement that matter is indestructible. But

were it possible to collect the smoke and escaped gases and add them to the ash and weigh the whole, the total weight (less the oxygen used in the combustion) would be found to be equal to that of the original unburned piece of wood. Compounds composing the wood have been broken up; carbon has been combined with oxygen to form carbon monoxide (CO) and carbon dioxide (CO<sub>2</sub>); hydrogen has




Fig. 1.6 Santorio Santorio carrying out an experiment upon himself seited in a large balance (steel arm).

been oxidized to water (H<sub>2</sub>O); many other compounds have been decomposed and new compounds formed by the recombination of atoms momentarily freed. Did one know all the chemical reactions which had taken place, a balanced equation could be written (p. 7).

Nor is matter created in the sense that new atoms are formed. The growth of plants or of animal tissues is merely the accumulation of pre-existing atoms. A plant, for instance, acquires water, minerals, and nitrogen from the soil, and carbon from the carbon dioxide of the atmosphere; from these it builds its structure and gains in weight. The animal body takes in carbon, hydrogen, nitrogen, oxygen, and

other elements in the food as well as oxygen from the atmosphere Just as in the flaming wood, the rarbon is oxidized to carbon dioxide and the hydrogen to water, part of the introgen of the food in combination with other elements (see amino acids p 360) is built into body tissue, and the remainder excreted. The carbon dioxide and part of the water is exhaled in the breath. If the adult body's intake of all the elemental materials were collected, their total weight would be found to balance the weight of the materials excreted in urine, feces,




Fig 1.7 Illustrating electrolysis When an electric current is passed through a solution of an electrolyte (e.g. sodium chloride) the positively charged ions (cations) move to the cathode and the negatively charged ions (amions) to the anode

skin (perspiration) and breath. The body's weight would be found to have remained unchanged.

An Italian physician (Santorio Santorio) living in the sixteenth and seventeenth centuries (1561–1636) was the first to demonstrate these facts. He spent most of his time in the latter part of his life in a large balance (Fig. 1.6) weighed all that he ate and drank and tried to balance the weight of these with the weight of this exercta (urine and dreces) but found a discrepancy. The weight of that which he ate and drank was more than that of his excreta. He concluded that the difference was caused by the loss of water through the skin (1e. evaporation) and in the expired air. This loss which was not revealed by his weighings he called unsensible perspiration (see pp. 377–8).

Ionization or dissociation. Certain compounds, such as hydrochloric acid (HCl), nutric acid (HNO<sub>2</sub>), phosphoric acid (H<sub>2</sub>PO<sub>2</sub>), sulfure acid (H<sub>2</sub>SO<sub>4</sub>), carbonic acid (H<sub>2</sub>CO<sub>3</sub>) sodium potassium, calcium, and ammonium hydroxides (NaOH KOH Ca(OH)<sub>2</sub>, and NH<sub>2</sub>OH, respectively), sodium and potassium chlorides (NaCl and KCl), cop-

per sulfate (CuSO<sub>4</sub>), and a great number of others when in aqueous solution, separate into their constituent atoms or groups of atoms. The separated atoms are electrically charged and are called ions (Gk. tion, going, traveling), because when an electric current is passed through an aqueous solution of one of these compounds they move at (Fig. 1.7). This property of molecules to break up into electrically charged atoms is called *ionization* or dissociation.

Some ions, such as those of calcium, hydrogen, sodium, and potassium, are positively charged and move toward the negative pole or

Table 1-3	
Name of Electrolyte	Percentage Ionizea
Acetic acid	13
Carbonic acid	0.17
Hydrochloric acid	92.00
Hydrocyanic acid	
Nitric acid	92 00
Sulfuric acid	61 00
Phosphoric acid	8 50
Potassium hydroxide	91 00
Sodium hydroxide	91.00
Sodium chloride	85.00

cathode; they are therefore called cations. Such atoms are positively charged because they have lost the electron that joined them to the other atom of the compound (see p. 8). Others, such as chlorine (chloride ion), having gained the electron lost by the other atom, are negatively charged; they move toward the positive pole or anode and therefore are known as anions. The anions are frequently groups of atoms and are given specific names, such as nitrate (NO<sub>2</sub>), phosphate. (HPO,), sulfate (SO,), bicarbonate (HCO,), or hydroxyl (OH) ions. The nature of the charge carried by the atoms is indicated by a plus or a minus sign to the right and above the last symbol of the formula. Thus, the hydrogen ion is written H+, and the chloride and nitrate ions Cl- and HNO<sub>3</sub>-, respectively. If an atom carries two or more positive or negative charges, the respective sign is repeated; the calcium ion is written Ca++, and the phosphate ion HPO.

The compounds mentioned above ionize to the greatest extent when dissolved in water, and to a much less degree in solvents such as alco-

#### FLEMENTARY PHYSICAL & CHEMICAL PRINCIPLES

hol or acetone, and do not dissociate at all in the great majority of solvents. Water itself is very slightly dissociated into H and OH ions

14

Ionizable substances, since their solutions are capable of conducting an electric current, are called electrolytes. The movement of the two kinds of ion to the respective poles of an electric source whose terminals are immersed in a solution of an electrolyte is called electrolysis, Many compounds, such as cane sugar, do not undergo ionization, and are therefore called nonelectrolytes. A list of electrolytes and the percentages to which they undergo ionization in aqueous solution is given in Table 1-3 It will be observed that the percentages vary widely.

A few of the commoner electrolytes and the ions into which they

separate are shown in the following equations:

CaSO, Ca++ 50.-calcium sulfate calcium ion sulfate ion HCI H+ CIhydrochloric hydrogen chloride ion acid юп HNO. H+ NO,nitric acid nitrate lon hydrogen ion H2SO4 H+ + H+ SO,--suffuric acid sulfate 10n hydrogen ions H,CO, H+ HCO.carbonic scid hydrogen son bicarbonate ion NaOH Na+ OHhydroxyl ion sodium sodium ion hydroxide NaCI Na+ CIsodium chloride sodinii iou chloride ion H.PO. 3H+ PO.phosphoric acid hydrogen ions phosphate ion

Acids, bases, and salts. Anyone even without a knowledge of chemistry knows, in a general way, the nature of an acid. A typical acid is sour or sharp to the taste, and unless very weak gives a burning sensation to the skin; if strong, it is highly destructive to animal and vegetable tissues; it also turns blue litmus paper pink or red. These properties are due to the hydrogen ion (H+) (see p. 108). Every acid contains in its molecule at least one atom of hydrogen.

Lye, known chemically as sodium hydroxide (NaOH), and slaked lime (Ca(OH), are bases. Typical bases in solution have a soapy feel, and most have a bitter taste, they turn red litmus paper blue In strong solution, they also have a destructive action on animal and vegetable tissues. These properties are due to the hydroxyl ion (OH<sup>-</sup>) Bases react with acids to form compounds called salts, a metal or other element of the base replacing one or more hydrogen atom of the acid. Two such reactions are shown below.

	Ca(OH) <sub>2</sub> calcium hydroxide	+ 2HCl = hydrochloric acid	CaCl <sub>2</sub> calcium chloride	+	2H₂O water
or	NaOH sodium	+ HCl =	NaCl	+	H <sub>2</sub> O water

Strong bases, such as sodium, potassium, or calcium hydroxides, are also known as *alkalls* (Arabian, al, the + qaliy, [kali], ashes, which are rich in sodium). They interact with fats or oils to form soaps. The adjectives corresponding to the nouns, acid, base, and alkali, are acidic, basic, and alkaline.

Solutions. When a substance such as sugar or salt is stirred in water, it 'melts' away until the water is again quite clear. We say the sugar or salt has been dissolved, and the water and the dissolved substance are called a solution of sugar, in one instance, or of salt, in the other. The substance which goes into solution is called the solute, the liquid, whether water, alcohol, ether, oil etc., in which the substance is dissolved, is called the solvent. When a crystalloid (see below) is dissolved and the solution well stirred, the solute becomes distributed uniformly throughout the solvent, the molecules being evenly intermingled. Once a substance composed of molecules having a size of the order comprising sugar or salt has been dissolved, it is not possible to separate the solute again from the solvent by mechanical means, as by filtering. The molecules of the solute pass as readily through the pores of the filter as do those of the solvent.

A saturated solution is one which, at a given temperature, holds the maximum quantity of the solute which it is capable of dissolving, it is said to be saturated at that particular temperature. If the solution, at a given temperature, contains less than the maximum amount of solute which it can dissolve at that temperature, it is said to be un saturated

Saturated solutions of certain substances when slowly and carefully cooled remain clear. The solution is then supersaturated at the lower temperature. But solid particles of the solute separate out if one of its crystals is dropped into the solution, or often if the solution is even

slightly agitated. The substance continues to come out of solution until saturation at the lower temperature is reached.

Most substances increase in solubility with heating, but some are rendered more soluble by lowering the temperature. When the former class of substance go into solution, heat is absorbed—that is the tem perature falls. In the latter group, heat is given out, the temperature of the solution being higher than that of either the original solute or solvent.

A normal (N) solution is one which contains an equivalent (p 9) in grams of solute per liter. Thus, a normal calcium chloride (CaCl.) solution contains 55 5 grams of the salt (equivalent of calcium 20, and chlorine 35 46). A solution containing a fraction  $(\frac{1}{2}, \frac{1}{10}, \frac{1}{100})$  of the amount of the solute that is contained in a normal solution or one in which twice the quantity is dissolved, is referred to, respectively, as half normal one tenth normal (or decinormal), one hundredth normal twice normal, and so on, and is abbreviated N/2, N/10, N/100, or 2N

A solution containing per liter an amount of a substance in grams equal to its molecular weight is called a molar or a molecular solution Thus, a molar solution of calcium chloride (CaCl.) contains (40 + 71 --) 111 grams per liter Solutions containing fractions or multiples of this quantity may be made up giving tenth molar, twice molar etc, solutions

A physiological saline solution contains about 0.9 percent of sodium chloride or about the concentration of the mineral salts of human blood plasma

Suspensions and emulsions, Solid substances composed of relatively large particles of a diameter of 0.1 micron or more will remain suspended in water for a short time and then settle to the bottom of the container. But in suspensions of which the liquid medium is heavier and of a more viscous nature, such as mucilage, the particles settle very slowly, remaining suspended for long periods Blood is a suspension of cells in a fluid called plasma (p. 57) which is some five times more viscous than water. The cells are about 7 microns in their largest diameters and remain suspended for a relatively short time after the blood has been drawn from a vessel and placed in a test tube. The size of the particles of any substance as well as the consistency of the liquid medium is a factor in the rate of settling the larger particles settling more rapidly than the smaller ones. Particles of a size less than 0.1 micron remain suspended almost indefinitely, even in water (see suspensions, p. 17).

Liquid particles or globules of about the same size as the solid particles in a suspension, when dispersed in another liquid, constitute an emulsion. The two liquids do not lose their chemical individuality. As with suspensions, the smaller the particles the longer they will remain dispersed in the medium. Fat, oil, resins, gums, and many other substances of both animal and vegetable origin form emulsions with variable size of particles. Milk is a familiar example of an emul sion of fat globules in an aqueous medium. Emulsification of the fatty substances of the food occurs with the aid of the bile in the small intestine (p. 334).


Crystalloids and colloids Substances with molecules of relatively minute size—less than 1  $0\mu\mu^2$  in diameter—such as sodium chloride, sugar, copper sulfate, and the majority of chemical compounds, form true solutions, as described on page 15 They are called crystalloids Materials composed of larger molecules—from 0.1 u to 1.0 uu—or of smaller molecules clumped together to form particles of such a size, are called colloids (Gk kollae, glue) Glue, gelatin, egg white and other proteins, starch, rubber, various enzymes (p 288), many dyes, fine carbon particles as in smoke, or India ink, and a host of other sub stances, are colloidal in nature The animal body-blood plasma and the protoplasm comprising the tissue cells-is composed largely of solutions of colloids Solutions of colloids, also called emulsoids or suspensoids are frequently cloudy, turbid, or opalescent or, at any rate, not crystal clear as are true solutions. But some colloid solutions are transparent, so the turbidity or clarity of a solution cannot be resorted to as a means of distinguishing a true from a colloidal solution Colloid solutions have, nevertheless, certain unique characteristics when, for example a beam of light is thrown from one side into a colloidal solution, the diameter of the particles, being greater than the length of the waves of light scatters the rays, and a bright opal escent cone is seen, known as Tyndall's phenomenon. The particles are also made dimly visible if a film of a colloidal solution is illuminated by a light directed upon it from one side and observed under the

<sup>&</sup>lt;sup>2</sup> One micron (abbrev μ) equals 1300 millimeter A millimicron (abbrev mμ) equals 1300 micron or 1300 500 of a millimeter A micromicron (abbrev μμ) is 1300 300 of a millimeter A micromicron (abbrev μμ) is 2.00 of a micron or 1300 500 of a millimeter A micromicron (abbrev μμ) is 2.00 of a micron or 1300 500 of a millimeter A micromicron (abbrev μμ) is 2.00 of a micron or 1300 500 of a millimeter A mill

Colloid solutions are divided into two main classes called emulsoids or hydropholic colloids (Gk hudor water + philo I love) and suspensiods or hydropholic colloids (Gk phobos fear) The emulsoids have a strong affinity for water and for this reason are held more firmly in solution than are the suspensiods. The colloidal solutions of living matter—such as protein solutions—are for the most part emul soids The suspensiods have a tendency to come out of solution and are in general, solutions of norganic substances

microscope The ultramicroscope is based upon this fact, by means of this instrument the presence of particles, invisible through an ordinary microscope, is revealed

We do not use the term solute in speaking of the particles of a colloid solution as we do of a substance which goes into true solution. The particles are called the internal or dispersed phase of the solution, the medium in which the particles are dispersed is known as the external or dispersion phase. A reversal of the two phases under certain conditions, such as a fall in temperature, is a characteristic of certain colloids (namely, glue, gelatin, and starch), the particles (internal phase) joining together and enclosing the medium (external phase).



18

Fig 18 Diagram to show the rearrangement of the particles when a sol changes to a gel

(See Fig 18) The change is quite evident to the naked eye, the material no longer flows but has become firm or jelly like The hydrosel (sol) has been converted to a hydrogel (or gel) as the colloid is now called A change of this nature occurs in the conversion of the plasma protein fibrinogen to fibrin when the blood coagulates (p 103)

Most colloids owing to the relatively large size of their particles, can be separated from the dispersing medium and from crystalloids by means of a suitable filter. The procedure is called ultrafiltration. Such a process takes place in the body. The membrane forming the capillary walls, for example, serves as a filter which, while holding the plasma proteins to a large extent within the circulation, permits water and crystalloids to pass freely through to the surrounding tissue cells (see Fig. 3.10 p. 66). Nor do the plasma colloids normally enter the urine they are prevented from escaping from the circulation by the interposition of a fine filter in the kidney, the water and crystalloids of the plasma alone passing into the renal tubules (p. 277).

Colloids by means of dialysis can also be separated from any crystalloids which may be associated with them Colloid particles are unable to pass through certain membranes, such as sausage skin, frog s skin cellophane, etc., which are permeable to the smaller molecules

of water and crystalloids. If, therefore, a colloid solution containing salt is placed in a small sac made of sausage skin or cellophane and immersed in distilled water, the molecules of salt escape into the surrounding water and molecules of water pass into the colloid solution. This process continues until the concentration of salt is the same on the two sides of the membrane. If the distilled water is renewed repeatedly, the colloid solution is eventually completely freed of the crystalloid.

Fig. 1.9 Brownian movement. The successive positions of a particle of a colloid as observed by means of the ultramicroscope.



Brownian movements. This is an astonishing phenomenon first described about one hundred years ago by a British botanist, Robert Brown. When a colloid solution is observed with the ultramicroscope, random, erratic, zigzag movements of the particles are seen (Fig. 1.9). The movements occur ceaselessly and with great rapidity, the particles appearing to dart here and there like midges or a great number of rubber balls struck by invisible hands. The movements are not an inherent property of the colloid particles themselves, but are caused by the molecules of water and other molecules of the external phase coming repeatedly into near collision with them (see Kinetic theory, p. 243). It appears that actual contact of the smaller, rapidly moving molecules with the larger ones does not occur, it is prevented by mutual repulsion.

The particles of many colloids when in aqueous solution are electrically charged, carrying negative  $(OH^-)$  or positive  $(H^+)$  ions (depending upon the kind of colloid) adsorbed (see below) from the surrounding slightly ionized water (p. 109). Since like charges of electricity repel one another, the particles of a given colloid having the same sign of charge (+ or -) are prevented from gathering into large clumps; they are thus kept in a state of uniform dispersion. This fact, plus the Brownian movements just described, keeps the particles in a state of permanent suspension; they show little or no tendency to settle to the bottom of the containing vessel.

Adsorption. The property of any material to attract and hold molecules—solid, liquid, or gaseous—to its surface is called adsorption. The molecules or particles do not penetrate into the substance or

# 20 ELEMENTARY PHYSICAL & CHEMICAL PRINCIPLES

structure of the adsorbing body, adsorption, therefore, is to be distinguished from absorption. Many colloids tend to be adsorbed on the surface of other bodies. They, in turn, exhibit the property of according smaller particles to a pronounced degree which, owing to the

small size of the individual particles, offer in the aggregate an immense surface to still smaller particles in the dispersion phase. For this reason, finely ground charcoal and other colloids are most effective clarifying, decolorizing, or deodorizing agents the molecules of the offending material being thus removed from solution or from air

# Protoplasm. The Cell. The Tissues

Protoplasm is the material basis of all forms of life. The substance of the animal body-muscles, brain, kidney, liver, etc.-is composed largely of this "life stuff." Yet the exact chemical constitution of living protoplasm is unknown, for any of the means which might be employed to analyze it cause its death. Irreversible physical and chemical changes then occur, and we are no longer dealing with the substance whose composition we had set out to determine. The most that can be said with regard to the chemistry of protoplasm is that it consists of a watery solution of proteins, together with smaller amounts of carbohydrates (glycogen and glucose), lipids (fatlike materials), and inorganic salts (of potassium, calcium, sodium, magnesium, etc.). The word protoplasm (literally, the first thing formed) is itself no more than a generic term, for there are innumerable types of this material, its constitution varying in the different forms of life as well as in the different tissues of any individual. Furthermore, it is the seat of innumerable chemical reactions, oxidations and reductions, decompositions and syntheses; its chemical nature is therefore changing ceaselessly. Upon these changes the varied phenomena of life depend.

Physically, living protoplasm is a hydrosol or hydrogel belonging to the class of colloids already described as emulsoids. It is somewhat foamluke in appearance under powerful magnification; but it is questionable whether this is a natural property of protoplasm or merely caused by the technique used in preparing it for examination.

# The Physiological Properties of Protoplasm

The various physical and chemical processes of plant and animal life are simply the manifestations of the properties of protoplasm. It is these properties which we recognize as distinguishing the living from the nonliving world.

The protoplasm of animals possesses four fundamental properties (1) irritability or excitability (2) conductivity (3) the power to convert the potential energy of food material into other forms of energy—thermal mechanical chemical or electrical and (4) growth

Irritability or excitability. These terms refer to the ability to respond to a stimulus. A stimulus may be defined as some change in the environment occurring at a sufficiently rapid rate. The annoba for example when stimulated as by the touch of a stiff hair or by heating the water in which it is immersed shows that it is alive by a movement




Fig 21 An amoeba

and a change in form of its minute body (Fig 2.1) Also the excised muscle of the frog shows its irritability so long as it remains alive by shortening (contracting) when stimulated in one or another of several ways (mechanically thermally chemically or electrically)

Conductivity. The effect caused by a stimulus is not confined to the region in the immediate neighborhood of the stimulated point but spreads throughout the mass of protoplasm. Thus when one end of a muscle is stimulated the contractile process travels to the other end. When the stimulus ceases the part of the muscle to which the stimulus was applied is the first to resume the resting state. The opposite end of the muscle is the last to become inactive. This phenomenon of the stimulated muscle illustrates the property of conductivity. Conductivity.

tivity reacnes its nighest development in nervous tissue. The disturbance set up in a nerve fiber by a stimulus is transmitted to the farthest end of the fiber, which in some instances is several feet long. In a warm-blooded animal, the disturbance or impulse, as it is called, is transmitted at the rate of some 300 feet per second—i.e., at about the velocity of a revolver bullet.

Metabolism. The ability which protoplasm possesses to liberate and convert to its own uses the energy stored in food material-is-referred to as metabolism (see also p. 351). The term embraces all those chemical reactions occurring in the tissues and from which energy is derived for the performance of muscular work or for sustaining the vital processes—e.g., the contractions of the heart, the activity of the nerve centers, the generation of body heat, and the manufacture of the digestive and other essential secretions (e.g., those of the ductless glands). Upon chemical reactions also depends the growth of the young animal. In the digestive tract, the food materials are broken down into simpler compounds which are absorbed by the blood and rebuilt to form body tissue.

# The Cell

In very simple forms of life—e g, the amoeba (Fig. 2.1)—a single small mass of protoplasm, called a cell, composes the organism's entire body. Such organisms are called unicellular. Other forms of life whose bodies are constituted of many such microscopic blocks of protoplasm are called multicellular. Higher animals are constructed of immense multitudes of cells grouped together into various patterns to form the different tissues of the body.

Though, as we shall soon see (p. 39), cells vary greatly in size and shape, the majority possess certain characteristics in common. Each possesses a smaller ovoid or globular body called the nucleus which in a typical cell lies near its center (Fig. 2.2) The protoplasm surrounding the nucleus and constituting, in most instances, the main bulk of the cell is called the cytoplasm.

The nucleus. The nuclear substance or caryoplasm is somewhat more viscous than the surrounding cytoplasm. The most important and essential nuclear material is in the form of granules and is called chromatin (Gk. chroma, color) which stains deeply with basic

<sup>1</sup> It is doubtful whether, in the living cell, the chromatin is in the form of granules. The latter are thought by some histologists to be the result of the reagents yased in preparing the cell for microscopical examination. It is probable that in the untreated cell the chromatin is in the form of threads.

dyes such as hematoxylin. The granules appear to be disposed along a netlike framework in the spaces of which is an almost structureless viscous substance called the nuclear sap. Chromatin material is rich in nucleoprotein, which is a simple protein combined with nucleic acid (Chap. 9). The nuclei of most cells contain one or two small round bodies called nucleoil (sing. nucleolus) which stain with acid dyes such as eosin Some of the masses of chromatin are conspicuous for their large size, and are known as caryosomes.




Fig. 2.2 The cell.

The function of the nucleus is closely related in some way to the metabolic activity of the cell, and the hereditary characters of the latter are bound up with the chromatin material, which breaks up into a definite number of separate pieces called *chromosomes* in the first stage of cell division (see Mitosis, p 25).

The cytoplasm. The cytoplasm stains with acid dyes so that with double staining the cell nucleus and the surrounding cytoplasm are strongly contrasted. There are six types of structure within the cytoplasm which will now be described: (1) mitochondria, (2) the centrosphere with its centriole, (3) Golgi apparatus or network, (4) fibrils, (5) vacuoles, and (6) inclusion bodies. The first four of these are known collectively as organelles or organoids; they are living structures—integral parts of the cytoplasm.

(1) Mutochondria (Gk. mitos, a thread + chondros, a granule). These are short threadlike, rod-shaped or granular structures scat-


tered throughout the cytoplasm (Fig. 2.2). The function of the mitochondria is uncertain but they appear to be associated with cellular metabolic processes.

- (2) The centrosphere or central body. This is a small sphere or oval of protoplasm lying close to one pole of the nucleus. It contains two small, sharply staining granules called centrioles. These latter in the resting cell (i.e., a cell not undergoing mitosis) are closely associated with cell reproduction; during mitosis they are centers from which fine lines are seen to radiate, forming the so-called asters.
- (3) Golgi apparatus or network. This was first described in 1898 by the celebrated Italian histologist Camillo Golgi. It consists of a meshwork of fibrils in the neighborhood of the centrosphere. Little is known of the function served by the Golgi apparatus, but in gland cells it appears to be closely associated with their secretory activity.
- (4) Fibrils. Very fine filaments are to be found in most cells, in which they form an infinitely fine interlacement throughout the cytoplasm. They are prominent elements in nerve and muscle cells, in which they run along more or less parallel lines. In some tissues they pass from cell to cell forming protoplasmic bridges.
- (5) Vacuoles. In certain cells, round or oval and apparently empty spaces known as vacuoles are present. Sometimes they form part of a system of fine canals. The wall of the vacuole consists of a definite membrane which, in certain unicellular forms, is capable of rhythmic contraction, the vacuole absorbing fluid from the surrounding protoplasm and discharging it to the exterior. Such a contracting vacuole serves in these forms as an excretory organ.
- (6) Inclusion bodies. These, as their name implies, are not integral parts of the cytoplasm; they are such particulate matter as granules of carbohydrate, representing the food in storage form, pigment (in skin, retina, etc.), and secretory granules—e.g., the zymogens of the salivary and pancreatic celts (p. 325).

The cytoplasm at the circumference of the cell is condensed to form what is usually referred to as the cell membrane. This acts as a semi-permeable membrane (p. 65), and also serves to stiffen the contour of the cell and to give it definite shape. The view is widely held that the cell membrane is composed mainly of lipid (fatlike) material.

Multiplication of cells—mitosis and amitosis. The body cells of developing and growing tissues multiply by a process called mitosis or caryocinesis. In this method of cell reproduction, complicated changes occur in the chromatin, preparatory to the division of the nucleus and cytoplasm and the formation of two new cells. The latter are called daughter cells.





Though mitosis is a continuous process, it is customary to describe it in four stages; prophase, metaphase, anaphase, and telophase.

Prophase. In the first stage or prophase, the chromatin loses its granular appearance and forms a number of bent or twisted and somewhat tangled filaments called spireme threads. The centrosphere divides into two equal parts, each of which contains a centriole. The newly formed centrospheres move apart, but remain connected by delicate protoplasmic strands which stretch between them. These strands are known as the mantle fibers and the fusiform structure which they form is referred to as the spindle. Protoplasmic filaments also radiate into the cytoplasm from the centrioles giving a starlike appearance to the latter, which are now called asters; the rays themselves are called astral rays. Toward the end of the prophase, the spireme threads break up into shorter and thicker sections. These are the chromosomes (Gk. chroma, color + soma, body). The nucleolus and the nuclear membrane have disappeared (A, B, and C, Fig. 2.3).

Metaphase. In the second stage or metaphase, the chromosomes have migrated to occupy the equatorial plane of the cell where they split lengthwise into two equal halves (D, Fig 2.3). The asters have by this time moved to opposite poles of the cell and the mantle fibers stretch to the corresponding chromosome halves; these latter are spoken of as homologous chromosomes. One member of each pair is composed of chromatin of the female parent (1e., contributed originally by the ovum, See Chap. 14), the other member represents the paternal (spermatozoon) contribution.


Anaphase. In the third stage or anaphase, the homologous chromosomes are completely separated and, as though moving along the mantle fibers, reach opposite poles of the cell The cell has become elongated (E, F, and G, Fig. 2.3).

Telophase. In the final stage or telophase, changes follow which are the reverse of those of the prophase. The chromosomes become longer and thinner, resembling the threads of the spireme; they are later joined by filaments to form a reticulum. The nucleus divides into two and the nuclear membrane and nucleolus are reformed in each new nucleus. The cytoplasm then divides, and two daughter cells are formed (H and I, Fig. 2.3).

It should be mentioned that the size, shape, and number of chromosomes vary with the species, but are always the same and characteristic for any given species of animal or plant. The chromosome number is 46 in man, 24 in the mouse, 4 in certain species of round worms, and 16 in the onion. It is also clear that, through the splitting of each chromosome into two, the nucleus of each daughter cell is assured of receiving the exact number of chromosomes characteristic of the species (see also Chap. 14).

In certain very primitive forms of life and occasionally, though usually under abnormal conditions, in the bodies of higher animals, reproduction occurs without any preliminary changes having occurred in the nucleus. This type of cell reproduction is called direct or amitotic division. In one type of amitosis, the cells divide into equal parts, half of the nucleus going to each new cell. This is called simple or binary fission (Fig. 2.4). In some forms (e.g., yeast cells), the cells reproduce by "pinching off" a part of their protoplasm, the newly

tissue cells, and the cells of the epidermis and the surface epithelium



28

Fig. 2.4 Simple or binary fission.

formed cell being much smaller than the original. This form of division is called "budding." Certain animal forms such as sponges also reproduce in this way (Fig. 2.5).

Sexual reproduction-i.e., the reproduction of an entirely new organism composed of many different tissues by the union of sex cells (spermatozoon and ovum)-will be described in Chapter 14.

With a few exceptions the cells composing the tissues of the adult bodies of higher animals do not multiply. A nerve cell, or a cell of the liver, heart, or muscle, for example, once it has been destroyed is not replaced by a cell of its kind. Its place is filled by fibrous (scar)

tissue. Blood cells, connective

of the mucous membranes, on the other hand, are replaced when they die by their own type of cell. But the adult cells of even these tissues do not multiply; a red blood cell or an epidermal cell, for example, once formed does not divide. The production of new cells of these tissues is a function of younger more primitive cells in the bone marrow or in the deeper layers of the epidermis, respectively. Were it not for the inability of the cells of most adult tissues-nervous, hepatic, cardiac, muscular, etc .-- to reproduce their kind, our bodies would, barring accidents, be immortal. Aging would be unknown; our organs and parts would be the seat of a ceaseless process of re-

juvenation. But it is inevitable that




Fig. 2.5 Budding of yeast cells.

with the passage of time the cells of vital structures show deterioration—undergo a progressive degeneration. This aging process commences at birth, and gradually, when the cells of some important structure are unable to perform their essential function, death occurs.

## The Tissues

Each of the various parts and organs of the body of a higher animal is highly specialized for the particular function which it is called upon to perform. Since any organ is a mass of cells, its specialization must depend upon the character of the cells of which it is composed. The amoeba performs all the fundamental functions—respiration, digestion, excretion, etc.—within the compass of a single cell. Like a solitary pioneer on the frontiers of civilization, it must do all those things which are necessary for its existence. In the multicellular animal, the cells have undergone differentiation into various types and, like the classes of individuals making up the population of a civilized community, each type, though highly proficient in the execution of one or other particular function, has lost the versatility of the primitive organism. The tissues of the body are divisible into four main types—epithelial, muscular, nervous, and connective—according to the type of cell of which each is composed. The structure of nervous tissue is described in Chapters 11 and 12.

The epithelial tissues Epithelial cells serve as a protective covering. They are arranged in a series of layers or strata to form the outer part of the skin (epidermis); they also line the digestive, respiratory, and urinary tracts Other types of epithelial cell have acquired secretory functions. The various external secretory glands (digestive, mammary, sweat, etc.), as well as some of the glands of internal secretion (thyroid, parathyroid, anterior pituitary), are composed of epithelial

ceils (Chap. 10).

In many situations the cells rest upon a very thin sheet of delicate, rather indistinct librils embedded in a cement substance. This is known as the basement membrane. In certain other structures—e.g., the mucosa of the digestive tract—the cells lie upon a much thicker layer of connective tissue called the membrana propria.

Epithelial cells are of several varieties. Some, the columnar type, are relatively long and narrow, appearing under the microscope as slender columns set side by side like the stakes in a palisade (Fig. 2.6). In some situations, as in the linings of the nose, trachea (windpipe), and bronchi, the columnar cells are surmounted by fine hairlike




Fig 26 Columnar epithelium




Fig 27 Ciliated columnar epithelium

structures called cilia (Fig 27) Ciliated epithelium also lines the Fallopian tubes, epididymis, Eustachian (auditory) tubes, and the middle ear The cilia, which are some 3 or 4 microns 'in length, show an incessant motion 'They bend quickly in one direction and then more slowly recover their original position. The movement is repeated at the rate of ten or more per second. The rate is slowed by cold and accelerated by warmth. Carbon dioxide, ether, chloroform, or alcohol vapor causes cessation of the movements. In the case of the respiratory

<sup>2</sup> A micron µ is 1000 millimeter; a millimeter is about it inch



Fig 28 Cuboidal epithelium



Fig 29 Squamous epithelium

passages, the quick movement is toward the exterior—that is, from below toward the larynx or from above toward the nostrils. The cilia do not beat in unison, on the contrary, each beats a little time before its neighbor placed nearer the exterior. Thus, waves or ripples appear to pass over a ciliated surface like a field of standing wheat stirred by a breeze. The advantage of such a motion is obvious it serves to sweep mucus and dust, or other small particles, from the respiratory passages. In the Fallopian tubes the cilia act to propel the ova to their destination (Chap. 14)




Fig 2 10 Stratified squamous epithelium,

32




Fig 2 11 Transitional epithelium lining empty urinary bladder

Epithelial cells, that are about as broad as they are long, are called cuboidal (Fig 28) Another type of epithelium is composed of thin platelike or scalelike cells, and is therefore termed squamous (squama - scale) (Fig 29) In some situations, as on the inner surface of the tympanic or drum membrane (Chap 13), the epithelium consists of a single layer of squamous cells lying edge to edge as in a pavement. In other places, as in the skin, cornea, and lining of the mouth and vagina, there are several strata of cells. The deepest ones are cuboidal or columnar, those covering the surface are squamous Between these two types is a series of layers whose cells decrease in thickness in each successive stratum nearer the surface. This type of epithelial tissue is termed stratified sauamous epithelium (Fig. 2 10)

the pelvis of tractified epithelium, called transitional, lines the pelvis of the Laurey, ureter, bladder, and part of the urethra Its surface cells are cuboidal rather than squamous. In the case of the bladder, this type of epithelium has the remarkable property of adjustment in relation to the state of the organ. In the distended bladder there are but three layers of cells, whereas in the empty viscus the number of layers is increased to six, the most superficial cells being rounded while the cells of the underlying layers are flask shaped or pear shaped (Figs 211 and 212)

The skin The skin consists of two main parts. The more superficial

of these is called the *epidermis* or *cuticle*; the deeper portion is known as the *dermis*, *corium*, or *true skin* (*cutis vera*). The epidermis is composed of stratified epithelium; distinguishing microscopical features at different depths enable it to be described in four layers which




Fig. 2 13 A section of skin cut vertically to the surface.

have been given special names. These layers from the surface inward are: (1) the stratum corneum or horny layer, (2) the stratum lucklum, (3) the stratum granulosum, and (4) the stratum germinativum (see Fig. 2.13).

The stratum corneum consists of clear scalelike cells which are very thin near the surface but increase in thickness in successively deeper layers. The cells of the horny layer contain a protein called keratin (p. 361) and are without nuclei; the most superficial cells are being continually shed and replaced from below. The stratum lucidum (clear layer) is a narrow glistening band in which the cells are very indistinctly marked off, and contain, as a rule, no nuclei.

### PROTOPLASM THE CELL THE TISSUES

The stratum granulosum is made up of two or three, sometimes more, layers of triangular, rhomboid, or spindle shaped cells, nu merous coarse darkly staining granules are to be seen in the cytoplasm The stratum germinativum (also known as the stratum mu cosum or the prickle layer) consists of a layer of columnar or cuboidal cells lying upon and anchored to the corium, and some four to




Fig 2.14 Vertical section through the skin showing appendages

six overlying layers of angular cells connected to each other by short protoplasmic spines or prickles. Multiplication of the cells of the skin takes place in this layer, especially in the most deeply situated cells, it is the only layer in which mitosis can be observed. It is from this fact that the layer derives its name. Most of the skin pigment is present in this layer (see below).

The dermis or corium is composed of loose connective tissue which contains the vessels and nerves of the skin. From its more superficial portion tonguelike processes—papillae of the skin—project into the overlying epidermis which appears to be molded over them

The skin contains numerous small tubular glands (sweat glands and sebaceous glands) which arising in the dermis pass vertically

outward through the epidermis to open upon the surface. These glands, together with the hairs and nails, are referred to as the appendages of the skin.

Beneath the dermis is a layer of connective tissue which usually contains much fat; it is called the *subcutaneous tissue*.

The hairs are developed from the epidermis but penetrate into the subcutaneous tissue. They are composed of an elastic horny material and have a definite structure. The shaft of the hair is that part projecting beyond the surface of the skin; the buried part is called the root. The lower end of the root shows a swelling known as the

hair bulb. The elongated socket enclosing the root of the hair is called the hair follicle: it is composed of several layers of epithelial cells surrounded by a connective tissue coat. Delicate strands of smooth muscle—the erectors of the hairs (arrectores pilorum; sing., arrector pilae)—arise in the outer part of the dermis and run obliquely inward to be attached to the hair follicles near their middle. These muscles cause the erection of the hairs of animals and the feathers of brids. They cause that




Fig. 2.15 The fingernail.

and the feathers of birds. They cause that roughening of the skin in man known as "goose flesh."

The nails. The human nails are horny curved plates firmly adherent to the dermis upon which they are molded. This part of the dermis is called the nail bed. The part of the nail embedded in the skin is called the root. The nails of man and the horns, claws, and hoofs of animals are modified epidermis (Fig. 2.15). The hard shiny exposed part of the nail is a thickened and greatly expanded stratum lucidum. The stratum corneum is represented only by the thin layer of cuticle at the base of the nail which tends to overlap the lunula—the half-moon-shaped pale area at the nail base. The nails grow in length by the multiplication of the cells composing the outer layers of the stratum germinativum.

The living color and the temperature of the skin depend upon the blood in a network of minute veins lying in the dermis immediately beneath the papillae, and called the subpapillary venous plexus. The blood in this plexus shows through the overlying epidermis. When the vessels are dilated, the skin is a deeper color and usually warmer

### PROTOPLASM. THE CELL, THE TISSUES

than when they are constricted. However, changes in skin temperature are not due so much to the caliber of the cutaneous vessels as to the rate of blood flow through them; that is, to the volume of blood in a unit of time that is drawn from the deeper, warmer parts of the body to the superficial vessels.

The hue of the skin—that is, its tendency to take on either a pink or a bluish tint—is dependent upon the degree to which the blood in the superficial vessels is oxygenated. If the blood contains the normal proportion of oxyhemoglobin, the skin has a pink or a scarlet hue; if it contains an abnormally high concentration of reduced hemoglobin, the skin will have a bluish tint. Therefore, in states in which the oxygenation of the blood in the lungs is interfered with as in pulmonary disease or strangulation, or the peripheral circulation is slowed as in heart disease, the skin is bluish (see cyanosis and anoxia, Chap. 6)

The permanent skin color—that is, the shade or tint of the skin independent of that caused by its blood supply—is due to fine granules of a pigment known as melanin present in certain cells of the stratum germinativum. In fair-skinned races, these cells, which are known as melanoblasts, are to be found only in the deepest cells of this layer but, in the darkest skins, they are present throughout the prickle layer and may even extend into the stratum granulosum.




Fig 2.16 Section of mucous membrane from the esophagus.

The skin carries out two essential functions: (1) to protect the underlying more delicate and sensitive tissues from mechanical, thermal, and chemical injuries, and from the inroads of bacteria (Chap. 5) or other pathogenic organisms; and (2) to aid in the regulation of the body temperature (p. 375). Small quantities of waste materials, especially urea, are excreted by the skin in the sweat.

This mode of elimination is probably of little importance in health, but is sometimes invoked in advanced kidney disease when the physician employs every means to promote the secretion of sweat

The skin is also an important medium through which the body

is supplied with vitamin D (Chap 9)

Mucous membrane The linings of the respiratory, digestive, and genitourinary tracts are called mucous membranes They consist of a layer or several layers of epithelial cells laid upon a basement membrane Beneath the latter lies a stratum of interlacing connec tive tissue fibers known as the membrana or lamina propria, which in many situations immediately overlies a relatively thick layer of loose connective tissue containing vessels and nerves, and small glands (Fig 2 16), this is called the submucosa In the esophagus,




Fig 2 17 Ciliated columnar epithelium, showing goblet

stomach, and intestines, but not in the respiratory and genitourinary tracts, a layer of smooth muscle known as the muscularis mucosae lies between the lamina propria and the submucosa The epithelial cells of mucous membranes differ in kind according to their location In the trachea and bronchi, for example, there are a number of cell layers, the most superficial being of the columnar ciliated type, but in the esophagus the cells are stratified squamous, and in the stomach and intestines they are simple columnar

Mucous membranes secrete a slimy, tenacious material called mucus This secretion, which contains a protein material (a glyco-

### PROTOPLASM THE CELL, THE TISSUES

38

protein, p 361) known as mucin, is formed within the cytoplasm of the epithelial cells. As the material accumulates the cell becomes distended and, finally bursting, discharges its contents. The mucus coats the epithelial surface, serving as a protection against injurious substances or to trap small foreign particles. From their flasklike appearance when filled with mucus, cells of this type are referred



to as goblet cells Though a certain proportion of the epithelial cells constantly secrete mucus the number is greatly increased by mechanical stimulation, or by infection or other harmful action. We are all familiar with the running nose of the common cold, and with the expectoration of mucus which accompanies inflammation of the trachea or bronch (Fig. 2.17).

Endothelium is usually classified as a type of epithelial tissue. It consists of extremely thin waterlike cells with wavy or jagged edges. (Fig. 2.18) which are laid edge to edge to form exquisitely delicate glistening membranes lining the chambers of the heart the blood vessels and the pericardial, pleural and abdominal cavities. The smallest lymphatic and blood vessels (capillaries) have transparent walls composed of a single layer of endothelial cells held together by a very thin layer of a cementing substance (see p. 50)

Glandular tusue The epithelial cells composing glands are arranged in various patterns (Fig 2 19) In one of the simplest types of gland the secreting cells form a single tube which opens durectly upon the surface (mucous membrane, skin, etc.) Such glands are called simple tubular. In other kinds of tubular glands, the deep part of the tube becomes branched In other glands again, the cells sur-

The term mesorhelium is now usually applied to the linings of the body cavities (pleural peritoneal etc.) the word endothelium being reserved for the linings of the heart blood vexels and lymphatics. But mesothelium and endothelium are similar in structure and origin.




Fig. 2.19 Diagrams showing four types of gland. I, simple tubular; 2, branched tubular; 3, simple alveolar; 4, compound alveolar.

round a central more or less spherical cavity into which the glandular secretion is received. The cavity with its wall of secreting cells is called an acunus, follicle, or alveolus (see Fig. 2.20). The channel or duct leading from the acinus may open directly upon the surface. But in large glands of this sort, the duct joins a similar one from a neighboring follicle to form a larger channel which unites in turn with others of the same size. Successive unions of this kind result in the formation of a rich rootlike system of ducts, the secretion being discharged to the surface ultimately through a single or a few large ducts. The salivary glands and the pancreas are glands of this latter

### PROTOPLASM. THE CELL, THE TISSUES

type, and, since the acini and the branching system of ducts bear a resemblance to a cluster of grapes, they are called acinar or racemose plands (L. acinus, a grape; racemus, a bunch of grapes).

In many glands, granules can be demonstrated within the cells; they vary in size, number, and distribution according to the secretory activity of the gland at the moment. These granules are the mother substance of the secretion; they constitute the material which, when the eland is stimulated to activity. is converted into the enzyme




Fig 2.20 Showing cross section of a racemose gland.

(p 288) or other essential component of the secretion. They are known in the former instance as zymogen granules. In an inactive or resting gland, the cells are loaded with granules, but when the gland is stimulated the latter move toward the cavity of the follicle, disappearing from the basal portions of the cells; with prolonged activity the cells are completely cleared of granules, the gland being then exhausted (see Fig. 2.21).

Muscular tissue. There are two main types of muscular tissue: struated or striped and unstructed or unstraped (also called plain or smooth). Skeletal or voluntary muscle—e.g., the muscles of the limbs and of other parts that carry out movements under the control of the will—is struated With the exception of that forming the heart, the involuntary muscle of the internal organs—e.g., stomach, intestines, bronchi and bronchioles, uterus, Fallopian tubes, etc—is nonstriated. The muscle fibers in the skin attached to the hair follicles, in the

walls of the blood vessels, and around the alveoli of various glands are also of the unstriated variety. The muscle of the heart though striated differs in important respects from voluntary muscle (p. 158). Muscle cells are long and slender and for this reason are usually referred to as muscle fibers. They are usually tapered at their ends which are applied side by side to form longer or shorter chains They vary very greatly in length, from a few microns—the length of cer-




Fig 2 21 Cross section of follicle of a racemose gland (eg, salivary) showing zymogen granules being discharged into alveolus

tain nonstrated fibers—to 35 or 40 mm as shown by the fibers of some skeletal muscles. In the large muscles of the skeleton, great numbers of such fibers are massed together into bundles. The bundles are bound together by connective tissue into larger masses. The contractile power of a muscle depends upon the combined effect of the innumerable fibers of which it is composed.

Upon close examination under the microscope a striated muscle

Upon close examination under the microscope a striated muscle fiber shows a series of alternating light and dark transverse bands. Each fiber is ensheathed by a delicate membrane called the sarco-lemma (Sarx, flesh [muscle] + lemma, husk), and running through its entire length are a great number of fine parallel filaments—the myofibrils. The latter are embedded in the cytoplasm, which is usually referred to as the sarcoplasm. Each myofibril is constituted of a number of alternating light and dark sections. They are called the J and the Q discs, respectively (see Fig. 2.22). A narrow dark line, known as the Z line or Krause's membrane, divides each light or J disc transversely into halves. The discs in all the myofibrils lie




Fig 2.22 A striated muscle fiber highly magnified A resting muscle two myofibrils teased out from end on right B contracting muscle C, cross section of a single muscle fiber

in line across the fibers—light to light and dark to dark. They are responsible for the crossbrinding of the fiber mentioned above. The myofibrils are the most minute parts of the muscle which can be seen with the ordinary microscope but even these delicate as they are, can be shown by means of the electron microscope to consist of a large number of fine threads of protein material called actomyosin (see p. 497). These infinitely fine threads are called micellae. They are the essential contractile elements of muscle fiber.

comere.

The striated muscle fiber contains one or more nuclei just be neath the sarcolemma; in a long fiber-several nuclei-may be seen. It having its nuclei placed near its outer boundary, the striated muscle fiber differs from most other cells, in which the nuclei are centrally situated. Each muscle fiber is separated from its neighbors and com-




Fig. 2.23 Cross section of a portion of a muscle.

pletely enveloped by a thin investment of connective tissue. This sheath which lies outside of the sarcolemma is called the endomysium. A connective tissue layer, the perimysium, also encloses groups of muscle fibers holding them in bundles. The envelope of connective tissue surrounding the entire muscle is called the epimysium (Fig. 2.23). See also page 479.

The striated muscle fiber as viewed under polarized light. Light particles (photons) vibrate in all planes at right angles to the direction of the beam of light. Certain crystals, such as a Nicol prism, have the property of intercepting the particles in all planes but one. The light is then said to be polarized. If, therefore, two Nicol prisms are set up in the path of the beam and at right angles to one another all particles are intercepted, and no light whatever gets through to the eye. The field is completely dark. If, however, a muscle fiber is placed between two such "crossed" prisms, the field is no longer black, but shows alternate light and dark bands corresponding to the striations seen by ordinary light, except that the light or J band is now dark and the dark or Q band is light (Fig. 2.24). This is




Fig 2.24 Muscle fiber as seen by polar ized light Resting muscle on left contracting muscle on right. Note that the Q and J discs become almost equal in thickness during contraction.

because the Q band contains some material, probably of a crystalline nature that is able to bend or twist part of the light transmitted through the first Nicol in such a way as to allow it to get through the second The Q band is therefore said to be doubly refractive or aniso tropic (Gk an not + isos equal + tropos turning—1e unequal refraction) The J dises are said to be isotropic (equal refraction)

Generally speaking unstrated or smooth muscle fibers are shorter than the striated variety and are usually spindle shaped (i.e., thicker in the middle than at the ends), they do not possess a sarcolemma the nucleus is more centrally situated, and the myofibrils are few and inconspicuous They show no regular cross markings (Fig. 2.25)

The connective tissues Connective tissue serves as a strengthen ing connecting and supporting material. It binds together masses of other types of cell and forms a supporting framework for various organs. Thus, it holds bundles of muscle fibers together, forms fibrous investments (capsules) for the kidney, liver, etc., and enters largely into the composition of the deeper layers of the skin. It also serves to fill in spaces between neighboring organs and parts, and



Fig 2.25 Smooth (nonstriated) muscle fibers




Fig. 2.26 Areolar tissue.

replaces tissue which has been destroyed by injury or disease (scar tissue). Connective tissue proper consists of a mass of long slender fibers embedded in a homogeneous jellylike matrix or ground substance. The ground substance predominates, cells being relatively few. The fibers are the elongated bodies or the long processes of the cells; nuclei are therefore sparsely scattered throughout the tissue. In many types of connective tissue, the fibers possess elastic properties, due to the presence of a protein known as elastin. The walls of the blood vessels are also well supplied with elastic fibers which are found in two layers. The larger arteries recoil after stretching almost as freely as a piece of rubber.

In some regions, such as in the walls of the trachea, bronchi and bronchioles, in the larynx, in the deeper layers of the skin, and in certain ligaments of the spinal column, the connective tissue is dense and composed largely of elastic fibers. The tendons of muscles, the ligaments of the majority of the joints, and the membranous cover-



Fig 2.27 Fibrous connective tissue.

### PROTOPLASM THE CELL THE TISSUES

ings of the internal organs are composed of bundles of strong, tough fibers which, though possessing little or no elastic property, are highly flexible These fibers contain a protein known as collagen and are therefore spoken of as the collagenous type of connective tissue fiber. The typical cell of the connective tissues is the fibroblast. From these the numerous fibers are derived. The fibroblast is a long, flat, spindle shaped cell with elongated processes or fibers. In tendons, membranes, liraments, etc., nuclei are relatively scarce, the sub-




Fig. 2.28 Adipose (fatty) tissue Clear spaces in the left hand figure are fat droplets in the figure on the right the fat has been stained black by osmic acid. The fat cells are shown in various stages of development.

stance of the tendon, membrane, or ligament being composed almost entirely, or at least predominantly, of fibers

Areolar tissue is the most generalized and widely distributed form of connective tissue. It is loose in texture and composed of an interlacement of collagenous and elastic fibers running in all directions a gelatinous ground substance, and a variable number of cells of different types scattered throughout. The meshes (areolae) between the fibers have suggested the name for this type of tissue. Areolar tissue is found beneath the skin (subcutaneous tissue), filling the spaces between muscles and the internal organs, and ensheathing the blood vessels.

Arcolar tissue is much more cellular and contains leucocytes, fat cells, and various reticulo endothelial elements (p 94), as well as the fibroblasts just mentioned (Figs 2 26 and 2 27)

Adipore or fatty tissue is a modified connective tissue Certain cells of arcolar tissue have the special ability to withdraw fat from

the blood stream and deposit it within their bodies. The fat collects within the cytoplasm as droplets which gradually increase in size until they fill most of the cell, the original cytoplasm and the nucleus being displaced to the periphery and flattened against the cell membrane (Fig 2 28). Adipose tissue consists of a mass of such fat laden cells. Other cells of the arcolar tissue have been crowded out while the fibers have largely disappeared. When the body loses weight, fatty tissue is one of the first to become reduced, it is drawn upon during starvation to furnish energy. The fat cells undergo a transformation. The fat droplets disappear the cells shrink and developing elongated processes assume the appearance of fibroblasts. Adipose tissue is developed chiefly in subcutaneous situations and in the arcolar tissue in relation to the abdominal organs.

Cartilage or gristle is also a modified connective tissue. There are three main types hyaline (glasslike) cartilage elastic cartilage and fibrocartilage.

Hyaline cartilage is translucent and of a bluish white color. It consists of large spherical or oval cells embedded in a nearly homogeneous ground substance. The cells are frequently found in pairs




Fig 2 29 A section of hyaline cartilage

or groups and are then flattened where they are in contact with one another (Fig. 2.29). A narrow zone immediately surrounding the cartilage cells stains more deeply and is thus marked off though not very sharply from the rest of the ground substance. It is referred to as the capsule. Hyaline cartilage is found covering the ends of the

### PROTOPIASM THE CELL THE TISSUES

bones inside joints (articular, cartilage), between the anterior ends of the ribs and the breast bone or sternum (costal cartilages), and in the nose, trachea, and bronch: The greater part of the skeleton of the embryo is first laid down in cartilage which undergoes gradual conversion into bone (ossification) during the development of the embryo and throughout the growth period of the young animal after birth. The process of ossification is not complete in the human subject until adult life.

Elastic cartilage is yellow in color It contains the same type of cell as that seen in hyaline cartilage, but the ground substance is not



Fig 2 30 Thigh bone (femur) from which the minerals have been removed by acid, it is then able to be tied in a knot

homogeneous, being reinforced by numerous elastic fibers. This type of cartilage is found in the external ear, epiglottis, and Eustachian tube.

Fibrocartilage is tough and dense. It is composed of collagenous fibers embedded in a homogeneous ground substance (matrix). This type of connective tissue is considered to hold a transitional position between hyaline cartilage and the dense connective tissue constituting ligaments and tendons. It is found in the form of discs be tween the vertebrae (intervertebral discs) and in the shoulder, knee, and hip joints. In each of the latter three situations, it forms a rim to the articular cartilage, thus serving to deepen the joint socket.

to the articular cartilage, thus serving to deepen the joint socket Bone or osseous lissue, another type of connective tissue, contains a large proportion of mineral matter, chiefly calcium and phosphorus (as tricalcium phosphate and calcium carbonate), but, also smaller amounts of magnesium and traces of fluorine, chlorine, and iron Some bones (e.g., those of the limbs) are developed by the deposition of minerals in cartilage. Others (e.g., the bones of the cranium) are formed by the mineralization of membraneous tissue.

But for the stiffening effect of these minerals, the bones would be no more rigid than many other tissues. Figure 2.30 shows a bone from which the minerals have been removed by treatment with acid.

Two types of osseous tissue enter into the construction of the long bones of the limbs. The shaft or diaphysis of a limb bone consists of a tube of hard compact bone, while the ends or epiphyses are composed of cancellous or spongy bone, covered by a shell of compact hone.

A section of compact bone from the shaft of the femur (thigh




Fig. 2.31 A, section of osseous tissue from the shaft of a teg bone, showing Haversian canals with concentric lamellae; B, enlarged drawing of a bone cell.

bone) is shown in Figure 2.31. The heavily mineralized substance is laid down in a series of concentric plates or lamellae around narrow channels, called Haversian canals. The latter run for the most part parallel to the long axis of the bone; they contain blood vessels. nerves, and soft connective tissue. The bone cells or osteocytes, which have oval bodies and numerous fiberlike processes, lie in small cavities-Howship's lacunae-between the lamellae. Fine canals (canaliculi) leading from the lacunae lodge the processes of the osteocytes. The canals of neighboring lacunae communicate with one another. The bony lamellae are thus pierced in all directions by a system of fine interconnected channels. The hollow center of the shaft of the bone is filled with a soft, fatty material called the yellow marrow.

The osteocytes are of two types—osteoblasts and osteoclasts.

### PROTOPLASM. THE CELL, THE TISSUES

50

Through the activity of the former type of cell, bony tissue is laid down and molded in the growing bone. The osteoclasts are responsible for the absorption of bony tissue. Thus, by the properly balanced actions of these two types of bone cell the growing bone is sculptured into its characteristic structure and shape.

Cancellous bone is made up of slender bars of osseous tissue called




Fig 2.32 Section of the upper end of the thigh bone (femur) showing cancellous bone. Note the heavier bars of bone along the lines of stress—i.e., through which the weight of the body is transmitted.

trabeculae which interlace with one another (Fig. 2 32), and give a spongelike pattern to the bony structure. The spaces between the trabeculae are filled with a soft, dark red tissue called the red marrow which is responsible for the manufacture of red blood cells, granular leucocytes, and probably also the blood platelets (Fig. 2.33). This substance consists of a mass of immature red/blood cells and granular leucocytes in various stages of their development, as well as the forerunners of the platelets—the megacaryocytes (Chap. 3)—and a few mature crythrocytes and leucocytes The ribs and bodies of the vertebrae as well as the ends of the long bones are composed of cancellous bone covered by a thin layer of compact bone. In the cranial bones, the spongy bone is also found saridwiched between two layers of dense bone.

Intercellular substances. Between the cells forming a solid tissue there is a semifluid viscous and structureless material resembling a soft jelly. In most tissues this gelatinous material is very scanty and serves as a glue or cement to hold the cells together and to permit water, dissolved substances, and even particulate matter to pass between the cells. Thus, material passes readily through the capillary wall from the blood into the surrounding fluid in the tissue spaces,




Fig. 2.33 Showing a section of red bone marrow EO, cosinophil leucocyte; ER, erythrocyte, LY, lymphocyte, ME. megakaryocyte, NE, neutrophil leucocyte; NO, normoblast; NU, extruded nucleus from normoblast; PR, primitive erythrocyte, PL, primitive leucocyte, R, reticulocyte, RE, reticulocyte with extruded nucleus; HE, earliest stage in development of the blood cells—ancestor of both red cells and granulocytes (hemocytoblast)

or in the reverse direction from the extravascular fluid into the blood stream. This material between the cells is called intercellular cement. In other tissues-e g., the connective tissues-the jellylike material is more abundant and forms a matrix in which cells and fibers are embedded. It is called intercellular ground substance. Chemically, it consists of hyaluronic acid. It is present as a homogeneous jellylike cell-free material in several situations, such as in the interior of the eyeball (the vitreous body), and as a thick covering for the umbilical cord, which has been known for many years as Wharton's jelly. Hyaluronic acid is liquified by an enzyme known as hyaluronidase, This enzyme is present in many tissues of the body. When injected subcutaneously, its liquifying action upon the ground substance of the subcutaneous tissues enables any injected fluid (e.g., physiological saline) to penetrate more widely from the point of injection. When certain tissue extracts (e.g., of testes) were observed to exert such an action, the factor responsible, whose chemical identity was then unknown, was referred to as the "spreading factor." Hyaluronidase is also produced by certain bacteria which are thus aided in spreading through an infected tissue.

# The Blood, Lymph, and Tissue Fluids; the Spleen and the Reticulo-endothelial **System**

The blood and tissue fluids of multicellular animals perform those functions which in unicellular forms are carried out through the . medium of their watery environment. Thus, the amoeba receives its oxygen through diffusion from the surrounding water. Carbon dioxide produced within the cell diffuses outward through the cell membrane. Respiration, the absorption of oxygen and the elimination of carbon dioxide, is therefore a relatively simple process for the amoeba. The processes of nutrition and excretion are accomplished in a manner equally primitive. Food materials pass through the cell membrane either in solution or as fine particles, and waste products pass into the surrounding medium. Other requirements of the unicellular organism, such as the maintenance of an optimum temperature and the proper degree of moisture, are dependent upon the immediate environment.

The elemental needs of each ceil in a multicellular form, from the most primitive type to the highest vertebrate, are the same as those of the unicellular organism; yet in the evolution of higher forms, the cells constituting their bodies have become far removed from immediate contact with the outside world. No longer can the exchange of respiratory gases, the acquisition of nutriment, and the excretion of waste products be carried out in the direct and simple manner practiced by the unicellular forms. The more primitive of the multicellular types overcame the difficulty by the development of canal systems which opened upon their exteriors and through which the water flowed freely, in and out, bringing oxygen and nutriment to the more deeply lying cells and bearing carbon dioxide and other excretory products away. This, the first attempt at a circulation, was an open one. As higher forms evolved, the circulation became closed—the waters of the environment no longer flowed through the body. Yet, the vessels of this closed circulatory system were filled with a liquid which took the place of and fulfilled the duties of the watery environment of the more primitive types. The blood and other body fluids may be looked upon as that environment which has been enclosed within the bodies of the higher forms, but which has undergone certain modifications in composition to meet the requirements of the more specialized cells of various types of which their bodies are composed.

In health, the chemical constitutions of the blood, and body fluids generally, vary with respect to both gases and solids in solution, within but very narrow limits. The mechanisms maintaining this relative constancy—the chief of which are the lungs, kidneys, skin, bowels, and ductless (endocrine) glands-interact in the most complex manner and carry out their functions with the greatest precision. They respond with the utmost delicacy to any condition which might tend to cause any departure from the normal state. This relatively unvarying composition of the internal environment has been named homeostasis by the late Professor Cannon of Harvard.

Functions of blood and body finids. The functions of the blood ad body fluids are summarized as follows:


- 1. Respiratory. The transport of oxygen from the air in the lungs the tissues, and of carbon dioxide from the tissues to the lungs.
- 2. Nutritive. The conveyance of food materials, glucose, amino cids, and fats from the alimentary canal to the tissues.
- 3. Excretory. The removal of the waste products of metabolism: g., urea, uric acid, creatinine, etc.
- 4. Maintenance of the water content of the tissues. Though the lood itself is contained within definite channels-arteries, capiliries, and veins (p. 121)-a constant interchange of fluid takes lace across the walls of the finest vessels (p. 65). It is the fluid lying

outside the blood vessels-the interstitial or tissue fluid-and not the blood itself which comes into actual contact with the cells of the solid tissues Such fluid is comparable to the watery environment of uni cellular forms of life and may be rightly looked upon as the cellular or internal environment of higher forms or, as the great French physiologist of the nineteenth century, Claude Bernard, termed it, the milieu interne Interstitial fluid closely resembles the blood plasma in composition and is identical in this respect to lymph (p 98) Through the medium of the interstitial fluid, the final stage in the passage of oxygen and food material to the cells, and the first stage of the journey of carbon dioxide from the tissues to the lungs and of other waste products to the kidneys is made

5 Regulation of the body temperature (see p 376)

6 Protective and regulative The blood, tissue fluids, and lymph contain certain chemical substances of a complex nature, antitoxins, lysins, and other antibodies which are the basis of the body's defense against bacteria and injurious agents of various kinds. The blood is also the vehicle by which the hormones of the different ductless glands (Chap 10) are carried to all parts of the body

The distribution of the body water. The total quantity of water in the body is around 70 percent of the body's weight. Part of this water, about 20 percent of the body weight lies outside cells and is



divisions of the body water

therefore called extracellular water. It comprises the water of the blood plas ma, of the lymph of the interstitual or tissue fluid, and the small amounts of fluid in the pleural and abdominal cavities and in the pericardial sac (p 115) The water within the tissue cells, the intracellular water, is much more abundant, amounting to about 50 percent of the weight of the body The partitions of the body water are shown in Figure 31 and the water content of the several types of tissue are given in Table 3-1 It may cause some surprise that the water content of the blood is no greater than that of some of the so called

solid tissues

In health, the volume of body water is kept remarkably constant Any tendency toward a reduction in volume arouses the sensation of thirst, and a balance is struck between the intake and the output

70

Table 3-1

White matter

	••	
Adipose tissue		20
Blood (cells 65, plasma 92)		80
Bone (marrow free)	25-	30
Connective tissue		60
Kidney		80
Liver		70
Muscle (striated)		75
Nervous tissue		
Gray matter		85

Percentage of water in several types of tissue

of water The amount of water taken into the body daily, or over any considerable period, must equal the amount lost from the body (i.e., the water output) Otherwise the body would lose or gain weight, the tissues would contain less (i.e., become dehydrated) or more than the normal amount of water

The water *intake* comprises water drunk as such or as other beverages, as well as the water in semisolid and solid food 'The *output* of water takes place through the skin, the expired air (which is saturated with water), and in the urine and feces

A balance sheet of the intake and output of water for an averagesized man, at moderate air temperature and humidity, is given in Table 3-2.

Table 3-2
The water balance
(24-hour period in cc)

Intake		Output	
Solid and semisolid food	1200	Skin	350
Water of metabolism 1	300	Expired air	500
Drinks (water, milk, coffee,		Urine	1500
tea, etc )	1000	Feces .	150
	2500		2500

Meats, bread and the so-called solid foods, other than those consisting largely of the state of the solid solid

The amount of water lost through the skin surface and lungs amounts to about one third of the total water output Water is lost through the skin not only in sweat, but as a result of the diffusion outward of water from the most tissues beneath the skin and its evaporation from the body surface. Even though no secretion of sweat occurs, as in the rare instances when sweat glands are lacking, water is continually being lost from the body in this way as from any most object. The water lost through skin and lungs varies, of course, with the temperature and humidity of the air. The greatest loss occurs in a hot, dry atmosphere. A rise in air humidity reduces evaporation from the body surface and consequently the amount of water lost in this way is also reduced. A rise in body temperature increases water lost both by increasing the rate of respiration and by increasing the evaporation from the body surface. In fever, therefore, excessive

water loss (dehydration), unless guarded against, is likely to occur. The physiological mechanisms controlling the body's water balance are of the utmost complexity and delicacy, they can only be touched upon lightly here. The composition of the blood (its osmotic pressure), the antiduretic hormone of the pituitary body (p. 438), the sense of thirst, and the activity of the kidneys are the chief factors concerned. Any tendency toward a reduction in body water is followed by a rise in the osmotic pressure of the blood which stimulates the output of urine, at the same time the sensation of thirst signals to consciousness the necessity for drinking and thus replenishing the body's store of water. Generally speaking, water can be retained in the body only as an isotonic solution (p. 67), that is, one with an osmotic pressure identical with that of the fluids within the tissue cells. Therefore, any dilution of the blood in electrolytes, or any undue concentration, is counteracted as just indicated, by increasing the water output or intake, respectively.

## Composition of the Blood

The blood is a highly complex fluid in which are suspended solid elements—the blood cells or corpuscles. Though blood is fluid, it must be classed as a tissue. As a matter of fact, the water content of blood is not greatly higher than that of most of the so-called solid tissues. If a tube is filled with blood and rotated rapidly in an instrument known as a centrifuge, the cells are thrown down to the bottom of the tube. The blood is thus separated into two portions—a packed

mass of cells which constitute around 45 (44 to 47) percent of the volume of the specimen and an almost clear, faintly yellow fluid called the plasma, which makes up the remaining 55 (53 to 56) percent. To obtain a precise measurement of the proportion of red cells to plasma, an instrument called a hematocrit is employed (see Pl. 1 and Fig. 3.7). The plasma contains proteins as well as many organic and inorganic substances in solution—nutritive and waste materials, antibodies, hormones, and other compounds of an unknown or imperfectly known chemical constitution. The specific gravity of whole blood is about 1.055; that of plasma about 1.027. The cells of blood are of three types—the red corpuscles or erythrocytes, the white corpuscles or leucocytes, and the platelets or thrombocytes.

The composition of blood is summarized as follows:

## Whole blood

### A. Cells.

- (1) Red corpuscles or erythrocytes
  - (2) White corpuscles or leucocytes
  - (3) Platelets or thombocytes

### B. Plasma.

- (1) Water 90 to 92 percent
- (2) Solids, 8 to 10 percent
  - (a) Proteins, 7 percent Serum albumin, serum globulin, and fibrinogen
  - (b) Inorganic constituents, 0.9 percent Sodium, calcium, potassium, magnesium, phosphorus, etc
  - (c) Organic constituents (other than protein) Nonprotein nitrogenous substances (urea, uric acid, xanthine, hypoxanthine, creatine and creatinine, ammonia, and aminoacids), neutral fats, phospholipids, cholesterol, glucose
- (3) Respiratory gases, oxygen and carbon dioxide
- (4) Internal secretions, antibodies, and various enzymes.

The plasma proteins. The total protein concentration of the plasma is around 7 percent. The plasma proteins are of three types—serum albumin (4 percent), serum globulin (2.7 percent), and fibrinogen (0.3 percent).\* The origin of the serum albumin is not known with certainty, but the experimental evidence points to the liver as the most probable source; there is little doubt that fibrinogen is produced in the liver. The serum globulin is derived from the lymphocytes. It was thought at one time that plasma protein was food protein

<sup>2</sup> Plasma from which the fibringen has been removed through clotting is called serum.

which was being transported from the digestive tract to the tissues It is now known, however, the food protein is not absorbed as such, but must first be broken down into its constituent amino acids (p 360)

By a special method (electrophoresis) the serum globulin can be split into three fractions called alpha, beta, and gamma globulins

The proteins of the plasma serve several important functions (a) they exert an osmotic pressure amounting to from 25 to 30 mm of mercury, which is a factor of first importance in the regulation of the blood volume (p 63), and in the exerction of urine (p 276), (b) they give viscosity to the blood, and thus aid to some degree in the maintenance of the blood pressure (p 147), (c) the fibrinogen, but not the other kinds, plays an essential role in the coagulation of the blood (p 103), and (d) the gamma fraction of the serum globulin (gamma globulin) is associated with the production of immune bodies (pp 101 and 215) The concentration of this protein is increased during the process of immunization

Most of the organic constituents of the plasma other than protein represent waste products of metabolism (e.g., urea, uric acid, etc.) together with nutritive materials (e.g., amino acids, glucose, and fats) absorbed from the intestinal tract. Of the inorganic constituents of plasma, sodium chloride is in highest concentration. The plasma also contains potassium, calcium magnesium sodium bicarbonate, and minute amounts of iodine and iron. Phosphorus is present in both inorganic and organic forms. The concentration of inorganic phosphorus is about 3 mg and that of the organic form about 40 mg per 100 cc of blood. Potassium is in relatively high concentration in the cells of the blood and in the cells of the solid tissues, but is in low concentration in the plasma and the body fluids, but is in low concentration in the blood plasma and the body fluids, but is in low concentration in the blood plasma and the body fluids, but is in low concentration in the blood cells and in the cells of the solid tissues. In some species this mineral is absent from the erythrocytes (see Table 3.3)

Table 3-3

Inorganic constituents of plasma, red cells, and whole blood
Milligrams per 100 cc, average values

<del></del>	Sodium	Potas sum	Cal-	Magne sium	Chlo- rine	Iodine	Iron
Plasma	340	20	10	27	370	0 001	0 1
Cells	0	420	0	60	190		100
Whole blood	160	200	5	40	250		50

The proportions of the inorganic elements in plasma, blood cells, and whole blood are given in Table 3-3. The concentrations of minerals in blood cells and plasma may be taken as representative of their proportions in the cells and fluids of the body generally. Serious results follow if, as in adrenal insufficiency (p. 425), these relationships are disturbed.

# The Red Blood Cells or Erythrocytes

The human erythrocytes of man and those of most higher animals are circular disc-shaped cells possessing no nucleus. They have a mean diameter of 7.2 microns (0.0072 millimeters), and a thickness of



Fig 3.2. Showing the diameters of a red blood cell.

about 22 microns (Fig. 3.2). The central portion of the cell is much thinner than the circumference which, therefore, appears as a rim around a central depression. This construction gives the cell a bi-concave contour or a roughly dumbbell outline when viewed edgewise. In shed blood, the red cells show a tendency to adhere to one another by their flat surfaces. This arrangement, which resembles a stack of



Fig 3 3 Showing erythrocytes in rouleaux.

coins toppled over (Fig. 3.3), is spoken of as a rouleau and the process as rouleaux formation. The cell is bounded by a membrane composed of protein in association with the lipid materials, lecithin and cholesterol. The chief function of the red cells is respiratory; namely, to carry oxygen from the lungs to the capillaries (p. 249) of the various tissues, and mainly in an indirect way, to transport carbon diox-




Fig 34 Hemocytometer for counting the red blood cells 1. pipette for diluting blood sample 2 glass slide provided with two platforms on each of which ruled squares marked by +, and shown highly magnified in 4, are engraved Each square has an area of 1/400 sq mm 3 stide in cross section with cover glass in position The skin near the base of the thumb nail is pricked with a sterile needle or a fragment of glass and blood drawn into the pipette as far as the mark 0.5 A suitable diluting fluid is then sucked up as far as the mark 101 The blood is thoroughly mixed with the diluting fluid (the glass bead in the dilated part of the pipette facilitates maxing) The fluid in the stem of the pipette is first expelled a drop of the blood diluted I part in 200 is then expressed on to the surface of each platform of the glass slide and a cover glass laid on A trench bounding each platform on three sides provides for any overflow The depth of the film of diluted blood be

tween a platform and a cover glass is 1/10 mm. The volume of diluted blood marked off by one square is therefore 1/4000 or mm. The blood cells in a num ber of squares are counted and the average taken. The concentration of cells in cubic millimeter of blood is then readily calculated. If the average number of cells in each square is 7, then the total number in a cubic millimeter is—4000 × 7 × 200 = 5 600 000.

4000 × 7 × 200 = 5 600 000
(cu mm (average (dilu (cells per fluid over cells tuon) cu mm)

1 square) per square)
The white cells are counted in a similar manner but the blood is diluted 1 in 20 by a fluid which destroys the red cells a similar pipette but with appropriate graduation is employed

ide from the tissues to the lungs (pp 111 and 252). The ability of the cell to carry oxygen is dependent upon its containing a remarkable pigment called hemoglobin (p 68). It is to this pigment that the characteristic color of blood is due

The number of red cells in a cubic millimeter of blood is usually stated to be 4,500 000 for women and 5,000,000 for men. As a mat ter of fact the values in health are somewhat higher than these, and

in a robust young male 6,000,000 is not an unusual figure. The number shows some slight variation during 24 hours, being lowest in the early morning, but increasing gradually throughout the day. The number of red cells is somewhat higher (by half a million or so per cubic millimeter) in newborn infants than in older children or adults.

The number of red cells in a cubic millimeter of a given specimen of blood is determined by counting the cells beneath the microscope. The instrument used for such a purpose is called a hemocytometer (see Fig. 3.4). The actual procedure followed in the enumeration of the red and white cells is described in the legend By knowing the number of red cells per cubic millimeter (i.e., their concentration) as well as the quantity of blood in the body (p 63), an approximate estimate of the total number of cells in the circulation may be arrived at. The figure is about thirty-five trillion (35,000,000,000,000,00).

Variations in the number of red cells. The red cell count (i.e., the number of red cells in each cubic millimeter of the blood) is increased: (a) during muscular exercise, (b) at high altitudes, and (c) as a result of a rise in environmental temperature. Also, a loss of water or of plasma from the blood (anhydremia, p. 83), although obviously not increasing the total number of circulating cells, will raise their concentration, and will therefore increase the red cell count.

The rise in the red cell concentration which occurs during muscular exercise, and at high environmental temperatures, is due mainly to contraction of the spleen which serves as a reservoir for red cells (p. 96). Contraction of the spleen also occurs when a person breathes




Fig. 3.5 A graph of the retheulocyte response to low tension of oxygen in the atmosphere (high altitude). (After Barcroft.)

62

rarefied air (i.e., air having a low pressure of oxygen) either in a closed chamber, in which the atmospheric pressure is reduced arti ficially, or in airplane or mountain ascents. An extra supply of red cells is thus discharged into the circulation. If the breathing of rarefied air is continued over a long period, as by those living in mountainous regions the red bone marrow—the tissue wherein the red cells are manufactured (p 50)-is stimulated its bulk is considerably in creased and the red cell count is permanently raised A count of 8 000 000 is not unusual in mountain dwellers. When the red bone marrow is suddenly stimulated by a low oxygen tension as when a person who has been accustomed to live at a low altitude ascends to a great height (over 10 000 feet), the cells that are discharged into the general circulation are in a somewhat immature stage of their de velopment These cells, which normally are less than 1 percent of the red cells in the general blood stream may increase to constitute 5 percent or more of the total number of red cells These slightly im mature cells are known as reticulated cells or reticulocytes (p. 70, see also Pl 2 and Fig 35)

Blood whose concentration in red cells has been raised either by contraction of the spleen or by the greater manufacture and discharge of corpuscles from the red bone marrow, is capable of carrying a




Fig 3.6 Some abnormal types of erythrocyte found in the blood in

greater load of oxygen from the lungs to the tissues This is a dis tinct advantage during muscular exercise or when the pressure of oxygen in the atmosphere is low In the latter instance the high red cell count helps to compen sate for the low oxygen pressure of the atmosphere

When the temperature of the environment is raised the dis charge of blood from the spleen serves the useful purpose, through augmenting the volume of circu lating blood of increasing the capacity of the body's heat radiating system and thus aiding in the

regulation of the body temperature (Pl 6A) Abnormalities in the form and structure of the erythrocytes Nu cleated cells containing hemoglobin and of the same size as erythro-

cytes are known as normoblasts (see p 70), they are normal con

stituents of the red bone marrow, but are only seen in the general circulation in anemia or other abnormality of the blood. Also, the presence in the circulation of reticulocytes persistently over 1 percent is abnormal, or at least unusual. In disorders of the blood, red cells in very early stages of their development and containing little or no hemoglobin appear in the circulation, as well as deformed erythrocytes known as poikilocytes. These latter include mulberry, shrunken. hourglass, and other bizarre shapes. Shrinkage of the erythrocytes with the production of eminences on their surfaces, like the mulberry forms just mentioned, can be induced by immersing the cells in a hypertonic saline solution (p. 68), when, as a result of the higher salt concentration in the surrounding fluid as compared with that inside the cell, osmotic force causes the transfer of fluid from the cell. Such shrinkage and deformation of the cell is called crenation (Fig. 3.6); the process whereby it is produced is the reverse of that causing hemolysis (see p. 84).

Cells below or above the normal size are characteristic of certain anemias (p. 72).

## The Volume of the Blood

The total quantity of blood in the body of an average-sized man (70 kilograms) is around 6 liters (over 6 quarts), or about one eleventh of the total weight of the body. It can be measured in the living sub-




Fig. 3.7 Hematocrit. Blood to which sodium oxalate or other anticoagulant agent has been added is drawn into the graduated tubes which are then placed in the holder and rotated in a centrifuge at a rate of 3000 revolutions per minute. The blood is thus separated into a red (cells) and a straw-colored portion (plasma), the proportions of each being easily determined by means of the graduations on the tubes.

ject by injecting a solution of a harmless dye into a vein. Vital red is the dye most commonly used for this purpose. A few minutes after its injection the dye becomes thoroughly mixed with the circulating blood A sample of blood is then withdrawn from a vein by means of a syringe, and the plasma separated from the cells by centrifuging The plasma, which is now colored pink by the dye, is compared with a standard solution containing the dye in known dilution The extent to which the sample of injected dye has been diluted in the blood stream is thus arrived at From this the total volume of the plasma is calculated. But it is desired to know the volume of the whole blood—1 e, of cells and plasma. The next step is, therefore, to determine the proportion of cells to plasma in a sample of the subject's blood. This is done by centrifuging a sample of blood in an instrument known as an hematocru (see Fig. 3.7). If the blood has the normal proportions of cells and plasma—namely, 45 percent and 55 percent, respectively—and the volume of the plasma as indicated by the dilution of the dye in the circulation is 3.4 liters (3400 cc), then the total blood volume is \$\frac{44}{2} \times 100 \leftarrow 6.1 liters.

Another method of more recent development in which the volume of the whole blood is measured employs a radioactive isotope of iron (Fe<sup>34</sup>) Erythrocytes from a healthy donor belonging to a compatible blood group (p 78) are 'tagged' with the isotope by injecting the latter into the donor some 24 hours previously A sample of his blood is withdrawn at the end of this period, its radioactivity is determined, and it is then injected into the subject whose blood volume is to be estimated. After time has been allowed for thorough mixing in the circulation, a sample of blood is drawn and a determination made by means of a Geiger counter of the quantity of radioactive iron that it contains, the dilution of the injected blood in the recipient's blood is the nealculated and the blood volume estimated.

Regulation of the blood volume. The passage of materials through membranes. The volume of the blood in health remains remarkably constant There are two main factors in its regulation (a) the osmotic pressure of the proteins of the plasma and tissue fluids, and (b) the hydraulic or hydrostatic pressure of the blood in the capillaries and of the fluids surrounding them

Osmotic pressure may be best explained by citing an example. If an aqueous solution of cane sugar or sodium chloride is placed in a small closed sac made of a suitable membrane and immersed in water, and if the membrance is permeable to water but impermeable to the molecules of the dissolved substance, then water will pass into the sac but the sugar or salt molecules will not pass out (that is, into the surrounding water). The pressure within the sac, therefore, rises, the walls of the sac become distended and may rupture. The force created

in this way is spoken of as the osmotic pressure. The dissolved material appears to "attract" or "draw" the water through the membrane. A membrane which permits the passage of water, but bars the passage of some water-soluble substance is said to be semipermeable with respect to that substance (Fig. 3.8). Now, the membrane forming the walls of the capillaries (which connect the small vessels on the arterial side of the circulation with the small veins, see p. 124) is semipermeable only in so far as the plasma proteins are concerned. It permits the passage of water and the relatively small molecules of sugar, urea, sodium chloride, and other crystalloids, but hinders the passage of the relatively very large molecules of the plasma proteins. Only very small quantities of the latter pass through the capillary wall. The proteins of the plasma, therefore, exert a force which lustrating osmotic pres-"draws" water from the surrounding tissue sure A sac formed of spaces, and tends to prevent the passage of water out of the vessels. The crystalloids (sugar, end of a tube The large sodium chloride, etc.) on the contrary exert dots represent sugar little or no osmotic pressure in so far as the plasma and the tissue fluids are concerned. The osmotic force exerted by the proteins of the the pores of the memplasma equals a pressure of from 25 to 30 mm of mercury (mm Hg).\* In order to overcome this osmotic pressure and drive water and dissolved salts through the capillary wall, an opposing hydraulic pressure must be applied This glass tube,




Fig 3.8 Diagram a semipermeable membrane is fastened to one molecules, the small ones, water molecules, Water passes through brane as indicated by the arrow. The pressure within the sac rises as shown by the height of the fluid column in the

is provided by the pressure of blood in the capillaries. The capillary blood pressure must, of course, be greater than the protein osmotic pressure. Such a relationship between osmotic and hydraulic pressures exists at the arterial end of the capillary. But a gradual fall in blood pressure occurs from the arterial end where it amounts to a little over 30 mm Hg, to the venous end where it is around 12 mm Hg. In the

<sup>3</sup> The pressure exerted by the plasma proteins is very low as osmotic pressures go. A solution of cane sugar or a solution of an electrolyte, such as sodium chloride. when separated from water by a semipermeable membrane (impermeable to the sugar or salt but permeable to water) would exert an osmotic pressure hundreds of times greater than that exerted by the blood plasma.




Fig 39 Illustrating the exchange of fluid between blood capillary, tissue spaces, and lymph vessel

venous end of the capillary. the osmotic pressure, therefore, exceeds the hydraulic pressure by several millimeters of mercury, and fluid is "drawn" back into the vessels (Fig 3.9). Thus, through the passage of fluid from one part of the capillary and its return to the circulation at another part, oxygen and nutritive materials are conveyed from the blood to the tissue cells, and carbon dioxide and other waste products from the cells to the blood. The osmotic and hydrostatic pressures

of the tissue fluids surrounding the capillaries are also factors which must be considered in the transference of fluid across the capillary membrane. The osmotic pressure of these fluids serves to counteract in part the osmotic pressure of the plasma. The tissue fluids also exert a certain, though small, hydrostatic pressure which offsets somewhat the driving force of the capillary blood pressure.

In order to visualize the passage of molecules or ions through a membrane, which appears to be a continuous sheet, it is supposed that the membrane is pierced by numberless "pores" or "channels," and, acting like a sieve, permits molecules below a certain size to pass but bars the escape of larger particles Such a mechanical conception, though it may not express the precise truth, does, neverthe-




Fig 3 10 Diagram to illustrate the retention of large molecules (plasma proteins) within the circulation, and the escape of smaller ones (sodium, chloride, glucose, urea, amino acids, etc.) The relative dimensions of the molecules and of the blood and tissue cells are represented only in a very general way

less, explain why a membrane, such as that forming the capillary walls, permits the free escape of the small molecules of water, sugar, sodium chloride, etc., while barring the transmission of the much larger molecules of the plasma proteins (see Fig 3 10). When, how ever, we come to consider other membranes, such as the one enveloping the erythrocyte, any structure of a sievelike nature fails to explain the free passage of some ions and the barring of other smaller ones. With the exception of the hydrogen ion the membrane of the red cell

Table 3-4

Blood		Tissue fluid	
Hydrostatic pressure 30 mm Hg		Hydrostatic pressure 8 mm Hg	
Net or effective hydrostatic pressure 22 mm Hg	y wall	<u> </u>	
Osmotic pressure 25 mm Hg	Capıllary	Osmotic pressure 10 mm Hg	
Net or effective osmotic pressure		<u> </u>	

Therefore, force driving fluid from capillaries = 7 mm Hg

is almost impermeable to cations, such as those of potassium  $(K^+)$ , sodium  $Na^+$ ), and calcium  $(Ca^{++})$ . As we have already seen, the interior of the blood cells and also the tissue cells are rich in potassium but poor or quite lacking in sodium, or calcium, or in both The blood plasma on the other hand, and the body fluids generally, have a relatively high concentration of sodium and a moderate content of calcium Anions, such as  $HCO_z^-$  (of carbonic acid) and  $Cl^-$ , pass freely across the boundary of the red corpuscles. The membrane thus holds potassium within the cells while preventing the entrance of sodium and calcium. These facts cannot be explained on the basis of size, for the  $HCO_z$  ion is larger presumably than the potassium, sodium, or calcium ion

Except for small variations which occur from time to time as a result of ionic interchanges the osmotic pressures of both the plasma and the fluid within the red cell are equal. The contents of the cell and the plasma are in osmotic equilibrium, or isotonic, which is the customary term. If distilled water be added to the plasma, or the cells

be placed in a salt solution of lower osmotic pressure than that of the interior of the erythrocytes—that is, the cells are now surrounded by a hypotonic solution—water is "attracted" or 'drawn" into the cells which then increase in volume and lose their hemoglobin (p 84). On the other hand, when, as described elsewhere, the erythrocytes are immersed in a fluid having a higher salt concentration than the plasma—that is, in a hypertonic solution—water is lost from the cells which become shrunken or crenated (p 63).

A homely illustration of osmotic effects is afforded by the behavior of an unskinned fruit such as a peach when it is placed in water (the water being hypotonic with respect to the fluids within the peach skin which acts as a semipermeable membrane) or when it is placed in a strong solution of sugar (which is hypertonic) In the first instance, water is absorbed by the peach which then swells and may burst its skin. In the second instance, the peach shrinks because water is 'drawn from it by the concentrated sugar solution in which it is im mersed.

### Hemoglobin

Hemoglobin, the coloring matter of the blood, belongs to the class of conjugated proteins, its molecule being composed of a pigment group called heme or hematin and a protein known as globin. The heme, in turn, is constituted of a pigment called porphyrin combined with iron.

Porphyrins are very widely distributed throughout animal and vege table life. The green coloring matter of plants known as chlorophyll, and pigments in the plumage of certain birds are porphyrin compounds. The brown pigment in the shell of the hen's egg is a porphyrin very closely allied to the porphyrin in hemoglobin. Porphyrins possess the property of combining with various metals—e.g., copper, cobalt magnesium (in chlorophyll), silver, nickel, iron, etc. Such. compounds are grouped as a class under the term metalloporphyrins. Heme is a member of this class, the metal in this instance being iron. Hemoglobin is therefore a porphyrin iron globin compound.

The constitution of the hemoglobin molecule may be shown concisely as follows

porphyrin + iron = heme (or hematin) heme (or hematin) + globin = hemoglobin

It has been mentioned that hemoglobin serves as a carrier of oxygen from the lungs to the tissues. This function is dependent upon the unique property, which the pigment possesses, of forming a very loose or unstable compound with oxygen. The hemoglobin molecule contains four atoms of iron, each of which can combine with a molecule

tains four atoms of iron, each of which can combine with a molecule of oxygen. But a true oxide of iron is not formed; the hemoglobin is therefore said to be oxygenated, not oxidized. The oxygenation of hemoglobin may be represented thus: Hb<sub>1</sub> + 4O<sub>2</sub> = Hb<sub>2</sub>O<sub>3</sub>.

The affinity of hemoglobin for oxygen is remarkable. When exposed to air it combines rapidly with oxygen—the compound being then called oxyhemoglobin. If the oxyhemoglobin is then exposed to then called oxyhemoglobin. If the oxyhemoglobin is then exposed to an atmosphere in which the oxygen pressure is low, the compound readily decomposes, oxygen being liberated. That is, the oxyhemoglobin is reduced; the compound is then called reduced hemoglobin. The ease with which hemoglobin gives up its oxygen, as the oxygen pressure to which it is exposed is lowered, shows the loose way in which the oxygen and iron are combined. In a true iron oxide, as we know, oxygen is in firm combination with the metal and is not released when the oxide is exposed to a low pressure of oxygen.

The absorption of oxygen by the blood. The quantity of oxygen which blood will absorpt is some sixty times greater than that which

which blood will absorb is some sixty times greater than that which can be absorbed by an equivalent volume of water. For example, if 100 cc. of water are exposed to an atmosphere containing the same percentage of oxygen as is present in the air of the lungs, about  $\frac{1}{2}$  co of the gas will be absorbed. Yet 100 cc of blood exposed to the same atmosphere will absorb nearly 20 cc of oxygen. The difference is due entirely to the hemoglobin. The total amount of blood in the human body (5 to 6 liters) will hold from 1000 to 1200 cc of oxygen. This quantity of oxygen is used by the tissues in 5 minutes or so during rest, and in a fraction of a minute during strenuous muscular exercise. In the absence of hemoglobin, the entire duty for the carriage of In the absence of hemoglobin, the entire duty for the carriage of oxygen would devolve upon the plasma (which is mostly water), and in order that this should be able to absorb and carry the quantity of gas necessary to satisfy the requirements of the tissues, its volume would have to be some sixty times greater than it is. The circulating 'fluid instead of being 5 or 6 liters, or about one eleventh of the body weight, would need to be more than 350 liters—over five times the bulk of the solid tissues!

The volume of oxygen which blood will take up when its hemo-globin is fully saturated is called the oxygen capacity of the blood. As just mentioned, this depends almost entirely upon the hemoglobin. The oxygen capacity of a sample of blood is therefore directly pro-portional to the quantity of hemoglobin which it contains. Each gram

of hemoglobin takes up a maximum of 1 34 cc of oxygen Now, each 100 cubic centimeters of normal human blood contains some 15 grams of hemoglobin So then (15 × 1 34 —) 20 cc is the oxygen capacity of 100 cc of normal human blood If, as in anemia, the hemoglobin content is below normal, the oxygen-carrying capacity of the blood is correspondingly reduced Further details of the role played by hemoglobin in supplying oxygen to the tissues—i e, its respiratory function—will be given in Chapter 6.

# The Formation and History of the Red Cells

70

The red cells are manufactured in the red marrow of the bonesskull, ribs, vertebrae, and to much less extent except in children, in the ends of the long bones of the limbs. The term hemopoiesis is applied to the process of blood formation-ie, to the production of all types of blood cell When it is wished only to refer to the manufacture of red cells, the term erythropoiesis is employed. The red cells pass through several stages of development before they are discharged from the marrow into the general circulation. At one of the earliest stages leading to its maturation, the cell is quite large, has no hemoglobin, and possesses a nucleus. The cells of this stage are called megaloblasts (Gk megas, megal, large + blastos, a germ or sprout). A little later, the cell acquires hemoglobin, becomes reduced in size, and is then called an erythroblast. The stage of the normoblast follows (Pl 2) The cell at this stage is about the size of a mature red cell, it contains its full complement of hemoglobin, but still possesses a nucleus In the final stage of maturation, the nucleus is expelled The cell is now called a reticulocyte from the fact that when suitably stained its cytoplasm shows a fine reticulated or filigree pattern. Many cells at this stage of development, as well as normoblasts, may be seen in normal red marrow, but cells of the earliest stage, the megaloblasts, are very scarce, and erythroblasts are not plentiful The reticulocytes are discharged into the general blood stream, normally, they constitute only about 0.5 percent of the total red cells in circulation (Fig 35, p 61) As mentioned earlier, when the bone marrow is stimulated, as by the low oxygen tension of the atmosphere which exists at high altitudes or by iron or liver administration in anemia (p 74), large numbers of reticulocytes appear in the general circulation They are also greatly increased in certain types of anemia

Within a few hours after their arrival in the general blood stream, the reticulocytes lose their reticulated pattern and are then mature red cells or erythrocytes. In health, normoblasts, or any cells younger than these, do not leave the bone marrow. But in certain types of anemia, red cells in any stage of development may be found in the general circulation.

The life of the red cell. The average life of the erythrocyte is around 110 days (100 to 124 days according to various estimates). There are no special means provided for its destruction. It simply wears out as a result of the stresses and strains to which it is subjected, and breaks up in the blood stream Small fragments hemoconia or blood dust as they are called, may often be detected during the microscopic examination of a specimen of normal blood. The fragments are finally disposed of by the spleen, which contains large mobile cells which engulf them

It has been estimated that in health something like a million cells per second disappear in this way and, of course, the same number must be formed afresh by the bone marrow. The number of red cells in the body at any moment, therefore represents the balance struck between the red cell wastage and the production of new red cells by the bone marrow.

Factors in the regeneration of the blood. The red cell, as we have seen, consists of a framework (stroma) composed of protein and lipid materials, and a complex pigment—hemoglobin The well-nourished body possesses adequate supplies of building materials for the manufacture of the cell stroma, and rarely is anemia due to any lack of the material required for the formation of this part of the cell Not a great deal is known with regard to the mechanism of hemoglobin manufacture. It might be thought that green foods since they are rich in the pigment chlorophyll (which, as mentioned on p 68, is related chemically to heme), would supply elements necessary for hemoglobin synthesis. Yet it appears that chlorophyll is not utilized to any important extent for this purpose. Nor are the porphyrin and "the iron in hemoglobin when taken in food utilized for the formation of hemoglobin. The heme cannot apparently be split into its constituents by the digestive enzymes (see p 288), and neither heme nor hemoglobin can be used intact for the production of fresh hemoglobin. The globin part of the molecule can, however, be utilized and the feeding of hemoglobin hastens the regeneration of blood pigment in certain forms of anemia. Not only is the iron in heme unavailable but the metal in other organic combinations is not utilized or is utilized.

very poorly Iron in *morganic* form (e.g., ferrous carbonate or ferrous sulphate) is absolutely necessary for normal erythropoiesis A diet deficient in iron leads to anemia (p. 73), as does also the lack of first-class protein, the amino acids of which are required for the production of globin. The porphyrin part of the hemoglobin molecule (p. 68) can be synthesized by the body from substances furnished in the food or from body tissies.

The ability of dogs on various diets to regenerate hemoglobin following hemorrhage was studied by Dr Whipple and his associates of Rochester, N Y (see Table 3.5)

Table 3.5

The influence of diet upon hemographic production

the mineral of the plant in the production					
Hemoglobin produced grams in 2 week period					
3					
3					
10					
15					
25					
45					
80					
80					
70					
95					

Of all materials investigated liver was found to cause the most rapid restoration of the hemoglobin Kidney and chicken gizzard were also highly effective As we shall see presently, these experi ments led to the discovery of a highly effective means of treating a very grave type of anemia in the human subject

It has been discovered within comparatively recent years that copper and cobalt though not themselves constituents of the hemoglobia molecule, are also necessary in some unexplained way, for the normal synthesis of the blood prement

#### Anemia

When the concentration of hemoglobin is below normal, the condition is called anemia. There are several varieties and grades of anemia. In very severe types, the hemoglobin may be as low as 10 percent of the normal quantity; i.e., it may be 1.5 grams per 100 cc of blood instead of 15 grams, as in health. The reduction in the hemoglobin may be the result either of there being fewer red cells, as in the most severe types of anemia, or of there being less hemoglobin in each cell, the total number of cells not being very greatly reduced. In some forms of anemia, both of these factors are present.

It has been mentioned that the number of red cells in the blood in health represents the balance struck between the loss of red cells through wear and tear in the circulation, and the production of new cells by the bone marrow. Anemia may result, therefore, from either increased destruction (or loss from the body) of red cells, or from diminished production. The anemias then may be divided into these two main categories.

A. Anemias due to blood loss or increased destruction of red cells.

1. Posthemorrhagic. In this type, a large amount of blood may be lost from the circulation either suddenly (acute hemorrhage) as a result of the opening of a large vessel—e.g., in accidental wounds, duodenal ulcer, etc.—or by repeated small hemorrhages (chronic hemorrhage).

2. Hemolytic. In this type, an abnormal number of red cells undergo destruction in the blood stream. Certain poisons—e.g., lead, arsenical preparations, etc.—cause destruction of the cells. In another type of hemolytic anemia, there appears to be some inherent defect in the erythrocytes themselves. They are more fragile and, in consequence, disintegrate more readily than usual.

B. Anemias due to defective blood formation. This group embraces anemias caused by:

- 1. Iron deficiency. Should the diet contain inadequate amounts of iron, anemia develops. In this type, the red cells, as a rule, are not greatly reduced in number, but they are smaller than the normal, and the concentration of hemoglobin in each cell is low. The pale color of the crythrocytes is usually quite evident upon microscopic examination. The qualifying terms microcytic (Gk. micros, small + cytos, a cell) and hypochromic (Gk. hypo, low + chromos, color) are frequently employed in referring to anemias of this class. Infants after the age of 6 months are likely to suffer from this type of anemia, because milk and other foods taken at this time are low in iron content, and the store of iron with which the baby came into the world has become exhausted.
  - Protein deficiency. A diet lacking in adequate amounts of protein of good quality is conducive to the development of anemia, for the amino acids necessary for the synthesis of the globin portion of

the hemoglobin molecule are absent or scarce. Food containing protem of high quality as in meat, liver, milk, eggs, etc , is just as important as iron for hemoglobin synthesis (see p 71)

3 Vitamin deficiency A severe dietary lack of vitamin C or of certain factors of the vitamin B complex, especially pteroylgiutamic (folic) acid, is sometimes a cause of anemia


Anemias due to deficiencies in the diet are usually referred to as nutritional anemias

4 Lack of the specific antianemic factor. Anemia due to this cause is commonly referred to as pernicious anemia because, until a cure was discovered in 1926 it resisted all forms of treatment and invariably caused death. The hemoglobin may fall to a very low percentage of the normal, due to the great reduction in the number of red cells Each cell actually contains a little more hemoglobin than does a healthy cell The cells are considerably larger than normal, and for this reason are called macrocytes (Gk macros, large + cytos, a cell) In pernicious anemia, very large cells known as megaloblasts are also seen, often in large numbers. They are very primitive cells, and re semble but are not identical with very immature crythrocytes of the same name found in small numbers in normal bone marrow. This normal bone marrow cell, as well as the megaloblasts of pernicious anemia, contains no hemoglobin and in appearance is not unlike the large lymphocyte (p 86)

The observations of Whipple and his associates, already mentioned (p 72), suggested to two physicians of Boston—Drs Minot and Murphy—that liver might be of value in the treatment of per-nicious anemia. The spectacular success which followed the feeding of liver to victims of this disease is now well known. As a result of this work of epochal importance, pernicious anemia was deleted, almost at once, from the list of incurably fatal diseases

It appeared that the liver normally discharged into the blood stream a principle essential for the maintenance of the blood forming function of the bone marrow. It was therefore concluded that the failure of the liver to supply this essential principle was the cause of per-nicious anemia. It has since been discovered that this essential substance is a vitamin of the vitamin B complex. It has been designated vitamin B<sub>1x</sub> and is present in many foods as well as in liver Extremely minute doses of this material given by injection to a patient with pernicious anemia arrest the disease almost immediately. But there is more to this interesting story. If pernicious anemia is caused by a lack of vitamin B<sub>12</sub> and this vitamin is present in most foods of any good diet, including food taken by those who develop pernicious anemia, it may well be asked, "Why then does the disease ever occur?" (See Fig. 3.11.)

Fig 3 11 Diagram to show the normal mechanism governing red cell production Black ducs, vitamin B<sub>12</sub> (extrinsic factor), ritippled ducs, enzyme (intrinsic factor).



The answer is that the primary cause of pernicious anemia is to be found in a defect in the mucosa of the stomach. The patient with pernicious anemia lacks an enzyme normally found in the gastric mucosa which is necessary for the absorption of adequate amounts of B12. When relatively very large doses of the vitamin are given by mouth to the pernicious anemia patient, as in a potent liver extract, a small part passes through the intestinal wall into the circulation and is capable of arresting the disease. Ordinary foods, however, do not contain the vitamin in such large amounts. But, as mentioned before, only a very minute dose of pure vitamin B12 (a mere fraction of a milligram) given parenterally is effective. Also, in the presence of a normal gastric mucosa, the amount of the vitamin in the diet is quite sufficient to prevent any lack of vitamin B12. Pernicious anemia is, therefore, essentially a vitamin deficiency disease. Powdered hog's stomach, since it contains the enzyme essential for the absorption of vitamin B12 from the food, also arrests the disease.4

4 The gastine enzyme has not been identified but is not one of the known digestive enzymer, such as pepsin The presence of such a principle was originally postulated by Dr. W. B. Castle of Boston and generally known as Castle's intrinsic factor. According to the theory which, until lately, was very generally accepted, the intrinsic factor acted upon an extrinsic factor in the food to produce a principle in the liver called the hematinic principle. The latter was the essential substance required for normal blood formation through an action exerted upon the bone marrow. There was much speculation as to the nature of the extrainsic factor, though some glumpse of the truth was gained, for it was shought to be a factor of the vitamin B complex or at least closely associated with the latter. The extransic factor is now identified as vitamin Ba, and as the so-called hematine principle as well.

#### THE BLOOD, LYMPH, AND TISSUE FLUIDS

Another factor of the vitamin B complex, quite distinct from B<sub>11</sub>, has been found within recent years to exert a curative effect in pernecious anemia. It is present in green vegetables, liver, kindey, and other foods. It was first named folic acid, but is now generally referred to by its chemical name piero/glutamic acid. The action of this factor differs in certain respects from that of vitamin B<sub>1</sub>. It has no power, for example, to prevent or arrest the degeneration of the nerve tracts in the spinal cord and the resulting parallysis, which is a very serious complication of permicious anemia. The essential physiological relationship between this factor and vitamin B<sub>1</sub>, is not altogether clear

5 Toxic agents which depress the function of the bone marrow Certain chemical poisons (e.g., benzene compounds, radium salts, or toxic substances) produced in disease (e.g., nephritis and various infections) depress or destroy the function of the bone marrow In some instances, the bone marrow is almost functionless, is much reduced in amount, and few red cells are produced to replace those which have been destroyed Circulating red cells as a consequence, fall to a very low level. The term aplastic is given to this type of anemia

#### Hemorrhage

76

Hemorrhage may be defined as the loss of blood as a whole (1 e, of plasma and cells) from the blood vessels The blood may escape from the body, as in accidents causing injury to an artery or vein—external hemorrhage. On the other hand, the blood may pass into the surrounding tissues, such as the brain or lungs or into one of the hollow viscera, such as the stomach or intestine. This is called internal hemorrhage. However, the general effects are practically the same whether the hemorrhage is external or internal. If the quantity of blood lost is large, and especially if it occurs suddenly, the subject suffers severely from lack of oxygen as a consequence of the reduction in the number of crythrocytes. There is a fall in blood pressure, as a result of the reduction in volume of circulating fluid, and the heartbeat increases in rate.

Of the physiological adjustments which ensue to safeguard the body against the dangers of blood loss, some are effected almost immediately after the hemorrhage, others not for some time. The first requisite, of course, is that the bleeding be staunched. This is accomplished through the clotting of the blood (p. 102), the contraction of the walls of the divided vessels, and the sticking together of the inner coat of the vessel. In this way the leak in the circulating system may be repaired. The capacity of the circulating system must also be reduced, in order to conform to the reduced volume of the blood. This is brought about through nervous reflexes. Nerve impulses are transmitted from centers in the brain and cord to the small blood vessels (arterioles, p. 122) in parts of the body, such as the skin, muscles, and intestines, whose functions are not immediately essential to life. The caliber of the vessels in these structures is narrowed. The blood pressure is therefore raised (p 153), and an adequate supply of blood to the heart and the vital centers in the brain thus maintained. The spleen responds to hemorrhage (p. 96) by contraction and the discharge of a quantity of blood into the circulation; thus the blood volume is at once, in part at least, restored. The blood volume is also augmented by the passage of fluid-mainly water and salts—into the vessels from the tissue spaces. This process commences almost immediately after the blood has been shed and, provided that the hemorrhage is not very severe, restores the blood volume to normal within a remarkably short time. The withdrawal of fluid from the tissues causes the subject to suffer from thirst, and the administration of fluids at this time is of value in hastening the return of the blood volume to its normal value. A longer time is required for the replacement of the plasma proteins and a longer time still for the complete restoration of the blood cells by the bone marrow. The time following the hemorrhage, at which the blood cells are brought back to their normal number, varies with the nature of the diet (p. 72), the quantity of blood which has been lost, and the recuperative powers of the individual subject. After the withdrawal of a pint of blood from a healthy person for transfusion purposes, from 6 to 8 weeks clapse before the red cell count returns to normal.


#### Transfusion .

When more than 40 percent of the blood is lost over a short period of time the body is usually unable to repair the loss unaided. Some artificial means of replacing the lost fluid must be resorted to. The intravenous injection of blood, plasma, or serum, or of some artificial solution with the object of restoring the blood volume is called transtusion.

Transfusion with whole blood. Blood obtained from another person (usually referred to as the donor) is the ideal transfusion fluid,

Strictly speaking, the word transfusion refers to the use of whole blood, serum, or plasma, and infusion to the use of other fluids.

for it is capable not only of restoring the blood volume but of furnish ing crythrocytes as well. The transfused cells survive for a considerable length of time (probably for from 100 to 125 days). Blood may not be taken indiscriminately from any person and transfused into any other. Great care must be exercised in the choice of the person (the donor) from whom the blood is taken, for the blood of one person may result in the death of another into whose veins it is injected. The patient who receives the blood is called the recurrent. Death may



78

Fig 3 12 Showing the method of determining the group to which a sample of blood belongs black discs = agglutnation, white = no agglutnation of it neither test serum causes agglutnation of the corpuscies of the blood being tested the latter must belong to group O If the corpuscies are agglutnated by B but not by A serum the blood belongs to group A If agglutnation occurs with A but not with B serum the blood is of group B II both sera cause agglutnation the unknown blood is placed in group AB

occur even though donor and recipient are closely related Such an untoward result is due to the patient's plasma containing a substance called an agglutinin while the donor's erythrocytes contain a complementary substance called an agglutinogen. When unsuitable blood is transfused, the donor's erythrocytes become gathered together into clumps—agglutination (Pl 3A, and Fig 3 12). Such clumps of cells block the small vessels, later the cells disintegrate and liberate their hemoglobin. The pigment in the process of excretion by the kidneys obstructs and damages the urinary tubules, and the patient dies as a result of the suppression of urine. Types of blood which, when mixed, behave in this way are termed incompatible. It has been discovered that the entire human population of the earth—of no matter what race—can be divided into four groups according to the reactions of their bloods when mixed together. The groups are designated, respectively, by the capital tetters, O, A, B, and AB, Table 3 6 shows the

reactions between plasma (or serum) and the corpuscles of the various groups according to the classification worked out by Jansky

It will be noted that the serum of group AB (vertical row on extreme right) does not cause agglutination of the corpuscles of any group That is, the red cells of any donor, it would be expected, could be injected with safety into a subject belonging to group AB A person belonging to group AB has therefore been called the "universal

Table 3-6 Serum

		0	A	В	AB
Corpuscles	0	-			-
	A	+	- :	+	_
	В	+	+	-	-
	AB	+	+	+	-

+ = agglutination, - = no agglutination

recipient' It is also evident from the table that the corpuscles of group O are not agglutinated by any serum (top horizontal row) A person of this group has therefore been called the "universal donor Serious reactions and even death may result, however, when the donor belongs to group O or the recipient to group AB The terms "universal donor" and "universal recipient' are therefore misleading The donor should when possible be of the same group as that to which the recipient belongs, there is then no fear of incompatibility in so far as the main blood groups are concerned 7 Or better still, the compatibility of the two samples of blood should be tested before the transfu sion is made. The clumping of the corpuscles which occurs when they

7 This precaution will not however exclude the danger of incompatibility with respect to certain other factors especially the Rh factor (see p 80)

<sup>6</sup> Previously the blood groups were designated by Roman numerals 1 II III and IV The capital letters used today refer to the agglutinogen or agglutinogens present in the corpuscles O means that the cells contain neither of the two main agglutinogens A that they contain one (A) B that they contain the other (B) and AB that they contain both In designating the two types of agglutinins (antibodies) in serum the Greek letters o  $\alpha$  and  $\beta$  are employed o means the absence of both  $\alpha$ ,  $\beta$  and  $\alpha\beta$ mean respectively that one or other or both are present Therefore to give the com plete characteristic of a given type of blood in respect to its agglutinogens (corpuscles) and agglutinins (serum) we would write O αβ (no agglutinogen both agglutinins) Aβ (agglutinogen A agglutinin β) Bα (agglutinogen B agglutinin α) and ABo (both aguittinger in a aguittinin f) but (aguittinger in aguittinger in a aguittinger in and the serum of AB group will not aguittinated because they contain no aguittinger and the serum of AB group will not aguittinate any cells because it contains no

are mixed with an incompatible serum is clearly seen under the microscope. Even to the naked eye the masses of agglutinated corpuscles may be seen as minute particles, like grains of cayenne pepper, floating upon the surface of the clear serum \*

The agglutination of corpuscles when incompatible bloods are mixed is in the nature of an immune reaction (Chap 5) resembling that whereby bacteria are agglutinated, and thus rendered innocuous by a specific substance (antibody) in the blood of a person infected with the same type of microorganism However, an antibody active against bacteria is as a rule, developed only after the blood has been invaded by the microorganisms, whereas the antibody to the agglutinogen in the corpuscles of an incompatible blood—that is, the agglutinin—is present naturally as an inherited characteristic. In this system the agglutinogen of the corpuscles is the antigen (Chap 5)

The blood characteristics are inherited according to Mendelian laws, and just as the offspring of a brown eyed and a blue-eyed parent may have either brown or blue eyes, so it is impossible to predict the blood characters of a child from those of its parents

Transfusion of blood is employed in other conditions besides hemorrhage, namely, in surgical shock extensive burns, in certain severe infections, and in extreme malnutrition of infants

The Rh factor Until a few years ago (1940), the four blood groups were believed to be the only ones that it was necessary to take into account when choosing a donor for transfusion, yet deaths following transfusion occurred from time to time for which no satisfactory explanation could be given In so far as the O, A, B, and AB groups were concerned the blood transfused in these fatal cases should have been compatible with that of the patient. When a survey was made of a number of these fatalities, it was discovered that they had all occurred either (1) in persons who had been transfused at some earlier time, that is, the fatality followed a second transfusion, or (2) in wamen who were pregnant or had given birth to a child Research workers interested in the subject recalled that Landsteiner and Weiner of the Rockefeller Institute had discovered that the in

e In order to determine to which of the four groups a particular blood sample begins the blood is tested with serum from blood groups A and B Samples of sera from these groups are available in small sealed valls which in all large hospitals are kept in the refrigerator ready for use If the corpuscles of the blood sample are agglutanted by A test serum but not by B it belongs to group B If agglutination occurs with B serum but not with A it belongs to group B If agglutination follows musting with both A and B sera the sample belongs to group AB and if neither A nor B serum causes agglutination the unknown serum belongs to group O (see Fig. 312)

jection of the blood of the Rhesus monkey into guinea pigs caused the formation in the blood of the latter of an agglutinin which caused the agglutination of the monkey's corpuscles and also of human corpuscles. The Rhesus corpuscles contain a substance—an agglutinogen—which, when injected into an animal such as the guinea pig whose blood does not 'itself contain it, stimulates the production in the recipient animal of an antibody—that is, of an agglutinin, Now, about 85 percent of members of the white race contain such an agglutinogen,




Fig 3.13 Diagram showing reactions caused by the Rh factor. Q = Rh-; ② = Rh+ (agglutinogen), Y = anti-Rh (agglutinin). A, reaction following a second injection of Rh+ blood into an Rh− patient. B, reaction caused by an Rh+ fetus in an Rh− mother.

or Rh factor, as it has been named (after the Rhesus monkey); the remaining 15 percent do not. In other words, the corpuscles of 85 percent of persons of the white race, in that they contain the agglutinogen, resemble Rhesus corpuscles, the corpuscles of the remaining 15 percent, in that the agglutinogen is absent, resemble guinea pig corpuscles. Blood which contains the Rh factor is termed Rh-positive; blood which lacks it is called Rh-negative. When the blood of a person who is Rh-positive is injected into the circulation of one whose blood is Rh-negative, the antibody—agglutinin or anti-Rh factor—is produced. Such transfusion merely causes the production of the agglutinin; no reaction follows. But, if a second, similar transfusion is given, 10 days or more (even up to years) after the first, a severe, often fatal, reaction follows, due to the action of the accumulated agglutinin or anti-Rh factor. The fatal reactions following an initial

Certain races—e.g., North American Indians, Japanese, Chinese, and Negroes of pure African stock—are 99 to 100 percent Rh-positive.

A2

transfusion in women who had borne a child are explained on an essentially similar basis. But in such cases the blood of the unborn child contains the agglutinogen (inherited from the father). That is, in all cases in which such a reaction occurs, the blood of the fetus is Rh-positive while that of the mother is Rh-negative. Evidently red cells containing the agglutinogen, or fragments of such cells, have escaped from the fetal to the maternal blood through the delicate membranes of the placenta separating the two circulations, and have stimulated the tissue cells to produce the anti-Rh factor in the mother's blood (Fig. 3.13).

Though an Rh-negative mother bearing an Rh-positive fetus suffers no ill effects unless transfused with incompatible blood, the fetus is often very seriously affected by the anti-Rh factor of the mother which enters the fetal circulation and exerts a destructive (hemolytic) action upon the red cells of the child, both before and after birth. The blood disease so called is known as hemolytic disease of the new born, or erythroblastosis foetalis; it is invariably fatal unless treated by large transfusions of Rh-negative blood

Transfusion with plasma or serum. Within recent years it has been generally recognized that in many cases of hemorrhage it is unnecessary to transfuse with whole blood The body possesses an immense reserve store of red cells in spleen, liver, and skin which can be drawn upon in such an emergency. Death from hemorrhage is, as a rule, due not to the reduction in the oxygen-carrying capacity of the blood but to the loss of the bulk of circulating fluid and the resulting fall in blood pressure which this entails Plasma or serum has therefore come into very wide use as a transfusion fluid There are certain very definite and great advantages to be gained by the use of these blood derivatives. In the first place, plasma or serum when collected from a number of donors and mixed together (pooled) does not cause agglutination of the transfused patient's corpuscles. That is, it is universally compatible Another great advantage of plasma or serum over whole blood is that either can be dried and stored in bottles, and thus be ready at hand for almost immediate use. All that is then required is the addition of sterile distilled water in an amount equal to that which had been removed by evaporation in the drying process In the preparation of the dried product, the plasma or serum collected from a number of donors is separated from the cells by centrifuging and decanted off. The cells, as a rule, are discarded. The liquid part of the blood is frozen and then dried in the frozen state by subjecting it to a high vacuum (6 mm Hg or less). This method, the desivac

process, is now employed in the preparation of <u>dried serum</u> (or plasma) on a large scale for the armed forces in Britain, Canada, the United States, and other countries.

The albumin of human plasma, dried and powdered, is also available. It is put up in sealed containers and ready for use after the addition of distilled water and the necessary blood salts.

Transfusion materials of nonhuman origin. The search for a suitable substitute for blood or its derivatives, plasma or serum, as a transfusion material has engaged the attention of scientists for a number of years. Provided that the material resembles plasma or serum in its physical properties and is capable of restoring and maintaining the blood volume for a reasonable length of time and thus of raising the blood pressure from a dangerously low level, the chemical composition of the transfusion fluid is of secondary importance. But the requirements of such an artificial transfusion material are many:

(a) The molecule of the dissolved substance must be of such a size that the fluid will not leave the vessels too freely. (b) The solution must exert an osmotic pressure and possess a viscosity approaching as closely as possible that of whole blood; these qualifications depend upon molecular size and shape. (c) It should be as nearly as possible isotonic with the contents of the erythrocytes (see p. 68) (d) It must, of course, be innocuous in every respect. In addition, it should be readily available, preferably cheap, and capable of being quickly and easily prepared for intravenous administration. Provided it is suitable in the respects just listed, there appears to be no valid objection to the use of some fluid other than blood or serum to fill the vessels immediately after hemorrhage. Nevertheless, for some reason not altogether clear whole blood appears to be superior to any other transfusion fluid in the treatment of shock or severe hemorrhage.

Among the solutions of nonhuman origin which have been used in the past or are under investigation and trial at the present time are: (a) saline (0.9 percent solution of sodium chloride), (b) a 6 percent solution of gum acacia, and (c) a 6 percent solution of isinglass.

Saline solution (0.9 percent of sodium chloride) is of little value in hemorrhage or shock, for the simple reason that it is not held within the vessels. The molecule of sodium chloride is so small that it escapes freely through the capillary walls into the tissue spaces. When, however, the water of the blood alone is reduced (anhydremia), saline is of benefit.

Gum acacia was introduced during World War I for the treatment of hemorrhage resulting from wounds, and was highly successful. The

#### THE BLOOD, LYMPH, AND TISSUE FLUIDS

RA.

reason for its success lay in the fact that its molecule is large and does not readily escape from the circulation. In the diution used (6 percent of gum in 0.9 percent salney, this substance exerts an osmotic pressure about equal to that of the plasma proteins. The fluid is therefore retained within the vessels, the blood pressure is elevated, and the circulation to vital structures maintained Gum acacia has come into disfavor, however, since the discovery that it damages the liver.

Isinglass is a protein very closely allied in its chemical and physical properties to gelatin. It is prepared from the swim bladders of fish (hake, cod, sturgeon, etc.) As as 6 percent solution in 0.9 percent




Fig 3 14 Showing the effect on an animal's blood pressure of the transfusion of isinglass solution. The animal's blood pressure had been lowered by hemorrhage to the very low level of 20 mm Hg at the start of the transfusion. The animal survived but undoubtedly would have died had it not been transfusion.

saline, it exerts an osmotic pressure comparable with that of the plasma proteins. It has proved successful in the treatment of acute hemorrhage in animals and in patients. It does not cause a foreign protein (antigenic) reaction or any other unfavorable effect (Fig. 3.14). As artificial blood substitutes, gelatin and dextran also have their advocates. The latter is a complex polysaccharide.

#### Hemolysis or the Laking of Blood

The plasma contain little or no hemoglobin. When normal blood is centrifuged, the red cells are thrown down and the supernatant plasma may then be seen to be almost colorless. But under the action of certain agents the hemoglobin is not retained by the crythrocytes, but escapes and colors the plasma This process is termed hemolysis or the laking of the blood The substance inducing the hemolysis is called a hemolysin or a hemolytic agent. Among the various agents

which cause hemolysis are: (a) hypotonic solutions; (b) chloroform, ether, and benzine, and certain other fat solvents; (c) bacterial poisons (toxins); (d) the venom of certain snakes; and (e) specific hemolysins. The depth of color given to the plasma varies from a faint pink to a deep red, according to the degree to which the erythrocytes have been attacked by the particular hemolysin.

Hypotonic solutions (p 68) act by disturbing the osmotic equilibrium between the interior of the red cell and the surrounding plasma. When, for example, distilled water is added to blood, the salts of the plasma are diluted and the osmotic pressure of the plasma is reduced below that within the cells Water is therefore "attracted" into the cell which, as a consequence, increases in volume and, finally rupturing, iberates its coloring matter. When the cell is completely hemolyzed his way, its colorless framework may still be seen. Such decolorzed cells are referred to as "ghosts."

The fragility test. If normal red cells are immersed in a series of typotonic solutions of sodium chloride graded in small differences of salt concentration, they commence to hemolyse when the percentage of salt is approximately 0.42. Hemolysis is complete when the salt concentration is approximately 0.36 percent. In hemolytic forms of anemia, the red cells are less resistant—that is, more fragile—than normal to hyoptonic solutions and commence to hemolyse at a salt concentration of about 0.5 percent, and are completely hemolysed at around 0.4 percent. In pernicious anemia, on the contrary, they show an increased resistance to hemolysis; the hemoglobin commences to escape from the cells at a salt concentration of 0.38 percent and is complete at 0.30 percent.

Chloroform, ether, etc., cause hemolysis, apparently, by dissolving the hpid materials composing the cell membrane and stroma. The manner in which bacterial and many other of the organic hemolytic agents act is not clear.

Specific hemolysins constitute a class of immune substances. For example, if an animal is given a series of daily injections of blood of another species, a substance is gradually developed in the serum of the recipient animal which has the power to hemolyse the erythrocytes of that species to which the donor animal belongs, but is not hemolytic for the erythrocytes of other species. (See Chap. 5.)

# The White Cells or Leucocytes


The leucocytes, unlike the red cells, possess a nucleus but no hemoglobin or other coloring matter. They number only about 8000



Fig 3 15 A film of normal blood illustrating the relative numbers of red and white cells and platelets A field as large aga n as that shown would probably contain no other white cell.

(6000 to 10000) per cu mm That is the red cells outnumber the white by about 600 to 1 (Fig. 3.15). The white cells are of two main types (a) Those without granules in the cytoplasm and with an un lobed nucleus these are called agranular leucocytes (b) Those with granules in the cytoplasm and a nucleus possessing two or more lobes these are termed granular leucocytes granulocytes or polymorphonu clear leucocytes (see Pl 2 and Fig 3 16)

The agranular leucocytes are of three varieties-the large and small lymphocytes and the monocytes. The first two types closely re semble one another except in size the small lymphocyte is about 8 microns in diameter the large from 10 to 12 microns. The small lymphocytes constitute about 25 percent of the white cells the large lymphocytes about 3 percent The monocytes are larger than the large lymphocytes measuring about 15 microns in diameter and their nuclei which are usually indented on one side or horseshoe shaped do not stain so deeply as those of the large lymphocytes Though there is some doubt as to the nature and origin of the mono



cytes, most observers believe that they belong to the reticulo-endothelial system of cells (p. 94). They may become very actively phagocytic. As a rule, they constitute no more than from 2 to 4 percent of the white cells.

The granulocytes are also of three types—the eosinophils, the basophils, and the neutrophils. The cytoplasm of the cosinophils is packed with coarse granules which stain with acid dyes (e.g., the red dye, cosin); the nucleus has, as a rule, but two lobes. Normally, these cells constitute only 2 or 3 percent of the leucocyte population.

The granules of the basophils stain with basic dyes (e.g., methylene blue). Their nuclei are also bilobed Basophils are scarce (not more than 0.5 percent) in normal blood.

The neutrophils are about 10 microns in diameter, and contain numerous fine granules which stain with neutral dyes, or with a mixture of an acid and a basic dye (e.g., eosin and methylene blue). When so treated, the granules are colored violet. The majority of the neutrophil leucocytes possess nuclei which are divided into from




Fig. 3.17 Showing stages in the life of a polymorphonuclear leucocyte (see text).

two to five lobes (Fig. 3.17). The number of lobes depends upon the age of the cell, the oldest ones having five or more, while the youngest (which are very few in number) show only a suggestion of lobulation. The neutrophils are the most numerous type of white cell, constituting from 65 to 70 percent of the total count.

The granulocytes are formed in the red bone marrow, whereas the large and small lymphocytes are produced in the lymph nodes and lymphoid tissue, generally, throughout the body (see p. 98). The monocytes are generally believed to be elements of the reticuloendothelial system. (See also p. 94.)

The functions of the leucocytes. The neutrophilic polymorpho-

The functions of the leucocytes. The neutrophilic polymorphonuclear leucocytes, together with the monocytes and other reticuloendothelial elements, constitute probably the most important means that the body possesses for its defense against invading microorganisms. The ability of these cells to attack bacteria depends upon their motility, and a proclivity for the ingestion of solid particles



Fig 3 18 Drawing of a neutrophil at half minute intervals to show motility and phagocytosis. The dots represent a group of bacteria.

(Fig 3 18) The latter action, which was first demonstrated by Metchnikoff, is termed phagocytosis (Gk phago, I cat) These two varieties of white blood cell are free lances among the body cells; they wander from place to place through the tissues, for practically no part of the body is barred to them They insinuate a process (pseudopodium), improvised at the moment from their cell protoplasm, through one of the joints in the endothelium of the capillary wall (Fig 3 19) Then, by causing the semifluid substance of the cell body to stream into the protoplasmic protrusion, they pass out of the blood vessels. By this process of diapedesis, as it is called, myriads of the white corruscles may pass out of the vessels in a remarkably




Fig 3 19 libustrating the passage of polymorphonuclear leucocytes from the circulation by diapedesis L, lymphocyte, N, neutrophil leucocyte.

short time. Reaching a point where the bacteria have entered the body and established a beach head, they surround the threatened area and proceed to destroy the invaders. If, for example, an actively inflamed region should be examined under the microscope, masses of neutrophils would be seen, and many of these would be observed to hold bacteria imprisoned within their bodies. As many as fifteen or twenty organisms may be seen at times within a single cell. It has been shown that the germs are ingested alive, and remain so for a time within the leucocyte.

time within the leucocyte.

When a tissue such as the mesentery or web of a frog, in which the capillaries are clearly visible beneath the microscope, is examined in the living state a short time after a suspension of bacteria has been injected into it, the small vessels leading to the sist of inoculation are found swarming with neutrophils. In the tissues round about, the amoeboid cells are seen moving somewhat ponderously hither and thither to engulf the offending bacteria. The monocytes, though much less numerous, also join in the general attack and show their phagocytic propensities to a marked degree. After the first flooding of the tissues with neutrophils and monocytes, numbers of the latter come to rest and together with other reticulo-endothelial elements of the tissues surround and isolate the infected area from the neighboring tissues surround and isolate the infected area from the neighboring tissues surround and isolate the infected area from the heighboring healthy tissues. Until this is accomplished the danger of the infection becoming more widely spread always exists. In their struggle against bacteria, equipped as these are with powerful toxins, many of the white cells are killed. These collect within the infected area together with exuded plasma, liquefied tissue cells, and a few red cells that have escaped through the injured walls of the capillaries. This mate-The circumscribing wall and its semifluid contents constitutes pus, and the so-called pus cells are dead leucocytes. The circumscribing wall and its semifluid contents constitute an abscess. A "boil" or furuncle is a small abscess of the skin and subcutaneous tissue which commences as an infection within a hair follicle. By the action of the phagocytes, aided by a protein-digesting ferment (protease) which they elaborate, the overlying structures whether connective tissue, mucosa, or skin are, in part, removed piecemeal. The wall of the abscess cavity over a small area of skin or mucous membrane becomes gradually eroded to the thinness of paper. The abscess is then said to "point"; it bursts soon afterward and discharges its content of pus.

Not only the bacteria themselves but any infected foreign material, such as a rose thorn, a wood splinter, bullet, or shell fragment, is loosened or eroded and, when possible, removed by the phago-

cytes (Fig. 3.20). Sterile foreign material-e.g, a catgut suture, or a blood clot-is broken down, and necrotic tissue is separated from the living by phagocytosis and by enzymes present in the blood tis-sues. Devitalized bone, though not removed in its entirety, unless it is of very small size, is, nevertheless, eroded and separated from the living tissue by the leucocytes. The disappearance of effete organs,




Fig 3 20 Showing migration of leucocytes (polymorphonuclears) to an infected area in the skin.

such as the tail and gills of the metamorphosing tadpole or the creep-ing muscles of insect larvae, is effected in a similar manner.

Of the functions of the other varieties of granulocytes-the eosinophils and basophils-little is definitely known. They are not mark-edly motile, and are not phagocytic. The eosinophils are increased above the normal of 2 or 3 percent in certain affections, notably, asthma, infestations with various types of parasitic worms, and in allergic conditions. The number of eosinophils in the circulation is strongly influenced by the adrenocorticotrophic hormone of the hypophysis (Chap 10), through its stimulating effect upon the produc-tion and liberation of adrenal cortical hormones. When the pituitary hormone or an extract of the adrenal cortex is injected into the body, a marked reduction in the number of circulating eosinophil leucocytes occurs. Indeed, the eosinopenic response constitutes the most sensitive test for the presence of ACTH in any biological fluid.

The lymphocytes are not amoeboid—that is, they do not progress

by the protrusion of pseudopodia. They are capable of a certain slow

progression, however, as a result of spasmodic movements of the cell nucleus. They have little phagocytic power and from all accounts

possess no proteolytic ferment.

The role played by the lymphocytes has been a subject of speculation for a number of years. It has long been recognized that these blood cells were concerned in chronic inflammatory conditions, for a great migration of small lymphocytes to the chronically inflamed tissues (small cell infiltration) was a common observation, but, until tissues (small cell infiltration) was a common observation, but, until recently, little was known definitely of the functions of these white cells. Modern research has revealed the important duty which they perform in the processes of immunity (p. 215). They give origin to one of the proteins—the globulin—of the blood plasma. The antibodies of paramount importance in establishing resistance to bacterial infections are closely associated with plasma gamma globulin. In experiments upon animals, it has been found that during an infective state the lymph nodes and other collections of lymphoid tis sue show a great destruction of lymphocytes They break down, their substance undergoing disintegration to furnish plasma globulin. The adrenal cortex and the pituitary gland, through its adrenocorticotrophic hormone (p 435), exert a controlling influence upon lymphocyte dissolution and globulin production and, consequently, upon the production of immune substances (antibodies).

Variations in the number of leucocytes. A rise in the number of white cells of the blood is called leucocytosis. The increase may be due to any one of the various cell types. More specific terms are frequently employed, such as, lymphocytosis, eosinophila, and neutro-philia for an increase in the number of lymphocytes, eosinophils, and neutrophils, respectively. The neutrophil count is raised markedly by acute septic infections. Among physiological conditions which cause a moderate increase in this type of white cell are muscular cause a moderate increase in this type of white cell are muscular exercise, pregnancy, and injections of adrenaline. Lymphocytosis occurs in certain chronic infections; eosinophilia is seen in asthma, in some skin diseases, and in diseases caused by parasites (e.g., intestinal worms). In order to determine which type of white cell is responsible for the leucocytosis, it is necessary to perform a differential count. This consists in examining a stained smear of blood under the microscope and counting the numbers of the different types in a total of several hundred cells. The percentages of the various types are then calculated. In some instances the proportions of the different types are altered, though no change occurs in the total white cell count.

A grave disease of the blood marked by an enormous, one might call it a malignant, increase of leucocytes is called *leukemia*. There are several different forms of the disease depending upon which type —granulocyte, small or large lymphocyte, or monocyte—is increased in number. The white cell count in this condition may reach 250,000 or more per cu mm, and many cells in the earlier stages of develop-ment are seen in the circulation. The red cells are usually reduced in number: i.e. there is an associated anemia.

Leucopenia is the term applied to a reduction in the number of Leucopenia is the term applied to a reduction in the number of leucocytes below the normal; it occurs in certain infectious diseases, notably pneumonia and typhoid fever. Certain drugs (e.g., amidopyrine) used for the relief of headache, or pain in other parts of the body, are believed to be responsible for a fatal disease in which the granulocytes are reduced to a very low level Resistance to infection is greatly lowered as a consequence. Scarcity of one or other type of leucocyte alone is called lymphopenia, neutropenia, or eosinopenia, respectively

The production and maturation of the leucocytes. The granuloottes, like the red cells, pass through several stages of development in the bone marrow (see Pl 2), and it is only cells in the final stage which normally reach the general circulation. The cells of the earliest stages are nongranular and show no division of the nucleus into lobes. These cells are called myeloblasts. Subsequent stages are marked by the appearance of granules which increase in number as maturation progresses and undergo differentiation into the three types, but the nucleus as yet shows no lobulation. These cells are called myelocytes. In the final stage—i.e., just before the cell is discharged from the marrow into the general circulation—constriction of the nucleus occurs at one point, but no definite division into lobes is evident. As already mentioned, the lobulation becomes pronounced and the number of lobes increases as the cell ages in the circulation. The life of a granulocyte is about 21 days.

The lymphocytes are produced not in the bone marrow, but in the

lymphoid tissues (e.g., lymph nodes, spleen, etc.).

Physiological factors concerned in acute inflammation. It is evident from what has been said in the preceding section that the functions of the neutrophil leucocytes are closely bound up with the physiological processes associated with acute inflammation. A substance is to be found in pus or other inflammatory exudate, known as the leucocytosis promoting factor (Menkin), which attracts the neu-

trophils to the injured tissues. A second chemical factor discovered by Menkin and named leucotaxin increases the permeability of the walls of the smallest blood vessels (capillaries, p 124), and a third, histamine, dilates the capillaries as well as increasing the permea bility of their walls. In inflammatory reactions, the capillaries and the vessels next in size, namely, the arterioles, are dilated as well through axon reflexes (p 520), the fine nerve endings being stimulated by the injurious agent, whether this is of a chemical (e g, bac terial toxin) or a mechanical nature. These effects upon the minute blood vessels cause the passage of an excessive quantity of fluid (plasma) from the circulation into the surrounding tissues, which re sults in swelling of the inflamed part. The dilated vessels bring more blood to the affected region which causes reddening and a local rise in temperature The swelling redness and heat-three of the cardi nal signs of inflammation-are thus accounted for by the vascular reactions The other signs of inflammation—namely pain and loss of function—naturally follow, the former being due to the involvement of the sensory nerves in the inflammatory process as well as to the pressure upon the nerves by the accumulated fluid. The temporary loss of function of a movable part is due in many instances, simply to the pain which results from any attempt at movement. This is na ture's way of encouraging recovery, by enforcing rest. In other in stances the loss of function—which may be permanent—is due to the inflammation having implicated the special structures upon which the particular function depends (See also p 233)

#### The Blood Platelets or Thrombocytes

These are small bodies about a quarter of the diameter of a red blood corpuscle. They do not possess a nucleus but their protoplasm con tains distinct granules (Fig. 3.21). Their origin is not known with certainty. The most widely held opinion concerning the origin of the platelets is that they are fragments of protoplasm broken or separated in some way from giant cells in the red bone marrow. These latter cells, known as megacaryocytes are of relatively enormous size (Fig. 3.22) and contain several nuclei massed together near their centers. Some authorities have suggested that the platelets are the remnants of disintegrated erythrocytes.

The platelets number about 250 000 per cu cm. They play an important part in the coagulation of the blood (p. 102)




Fig. 3.21 Blood platelets. A red blood cell shown for size comparison.

Fig. 3.22 A megacaryocyte. Red cell shown for size comparison.

#### The Reticulo-Endothelial System

This is a system of primitive cells normally present in the general connective tissue, lungs, spleen, liver, lymph nodes, bone marrow, and other situations. Many varieties exist, several of which possess phagocytic properties, and some of very large size are actively motile. One type of reticulo-endothelial cell, namely, the monocyte, has al-



Fig. 3.23 Macrophages containing phagocytosed carbon grains (left) and a red blood corpuscle (right).

ready been mentioned as being present in blood. Reticulo-endothelial cells of the spleen are responsible for the final disposal of the fragments of red cells which they engulf, the hemoglobin being freed of iron and converted to bile pigment (p. 329). The reticulo-endothelial cells in the blood sinuses of the liver are known as Kupffer cells. These, as well as those in bone marrow and the general connective tissues, also possess the ability to convert into bile pigment the hemoglobin liberated from disintegrated red cells. The reticulo-endothelial cells of the general connective tissues are of exceptionally large size. They play a prominent role in the body's defense against invasion by microorganisms. These cells are actively motile and phagocytic, ingesting bacteria and any other foreign material which may be introduced into the tissues. For example, after the injection of India ink (which consists of a suspension of carbon grains) into an animal, these macrophages. They are thus made conspicuous beneath the microscope (Fig. 3.23). In inflammatory conditions, they gather in large numbers around and within the inflamed part. In the lung, macrophages are active in the removal of foreign particles (e.g., dust, carbon particles) which have been carried into the alveoli by the inspired air.

The reticulo-endothelial elements in the neuroglia of the central nervous system are called *microglia*. They differ little from those in the general connective tissues.

# The Spleen

The spleen is an organ about the size of the fist situated in the abdomen behind the stomach and above, but to some extent overlapping, the left kidney. It contains, under ordinary circumstances, a relatively large amount of blood which is held in spaces (sinuses) lined by cells belonging to the reticulo-endothelial system. The blood is delivered into the substance of the spleen (splenic pulp) by small arteries. It then percolates into the sinuses through gaps between the cells forming the latter's walls (see Pl. 3B). The sinuses are drained by veins through which the blood is conveyed to the portal vein and thence to the liver. Dotted throughout the spleen, like islands, and surrounded by the pulp, are small lighter-colored areas composed of lymphoid tissue. These are the corpuscles of Malpighi (an Italian anatomist of

<sup>10</sup> The smaller phagocytic cells of the blood, namely, the neutrophils, are sometimes referred to as microphages.

the seventeenth century who first described them) Each of these areas is pierced near its center by a small blood vessel The capsule of the spleen contains smooth muscle, smooth muscle also penetrates its substance

The spleen is known to possess three important functions

1. Blood reservoir. It serves as a reservoir for blood which can be drawn upon to augment the blood volume when the need arises "1 The blood held by the spleen has a higher concentration of red cells than the blood of the general circulation, so that, when the organ contracts, not only is the total volume of circulating blood increased, but the





Fig 3 24 Changes in volume of spleen as a result of emotional excitement (after Bacroft) Sketch on left, R rest C, dog sees cat The numbers represent the relative sizes of the dog spleen Sketch on might, — rest, — smells eat, hears cat, — sees cat, — chases cat

number of cells per cubic millimeter of blood is also raised The several conditions which cause the spleen to contract are a rise in envir onmental temperature, strenuous muscular exercise, emotional excitement (fright, rage, etc.), carbon monoxide poisoning, hemorrhage, and rarefied atmospheres. The immediate stimulus to splenic contraction in the last three conditions is a low oxygen tension in the blood, but its great sensitivity to nervous influence is illustrated by the ob servation that even a sudden sound (e g, the slamming of a door) will cause it to contract (see Fig 3 24) The spleen also shows spon taneous rhythmical contractions at the rate of about two per minute (Fig 3 25) The small changes in blood volume which such move ments induce cause corresponding variations in the blood pressure The value of the organ in certain emergencies associated with a need for an additional supply of red cells is evident from an experiment performed by Professor Barcroft of Cambridge University upon guinea pigs. Two groups of animals were exposed to carbon monoxide gas. The concentration of gas in the atmosphere and the length of exposure were the same for both groups. The animals of one group,

it Experiments upon animals indicate that as much as 25 percent of the normal blood volume may be added to the circulation by contraction of the spleen

however, had had their spleens removed; these animals died, whereas those of the other group which had not been operated upon survived.

2. Destruction of damaged red cells. This function depends upon the presence of the large phagocytic cells (macrophages) already mentioned under the section on the reticulo-endothelial system. Healthy red cells appear to be immune to attack from these scavenger cells, but cells which have outlived their usefulness, or those whose structure is in some way abnormal, are readily disposed of. Fragments of red cells which have broken up in the blood stream are also ingested by the macrophages of the spleen.







Fig. 3.25 Showing effect of splenic contractions on arterial blood pressure, S.V. = splenic volume. B P = general arterial blood pressure. T = time, 5 sec B = baseline at 40 mm Hg Spleen in plethysmograph; a, before clamping splenic vessels, b, after clamping vessels, c, after removal of clamp. (From Barcroft and Misimaru)

3. The formation of lymphocytes. The lymphoid tissue of the spleen, as does similar tissue in other situations, lymph nodes, etc., manfactures lymphocytes. In embryonic life, erythrocytes and granulocytes are also formed in the spleen, but its ability to produce these types of blood cells ceases a little time before birth.

The spleen probably possesses other functions which have not yet been discovered. It exercises certain functions (e.g., bile pigment production and antibody formation) by virtue of the large numbers of reticulo-endothelial elements which it contains. It has not been demonstrated to possess an endocrine function (p. 403). That it is an important organ and guards the welfare of its possessor in several ways is undoubted. Nevertheless, it is not essential to life under ordinary circumstances, for when excised it is scarcely missed. Excision of the spleen (splenectomy) is practiced for certain hemorrhagic diseases in which the platelets are reduced in number, presumably as a result of some abnormal action which it exerts upon these blood elements. Splenectomy is also performed for a type of hemolytic anemia in which the red cells are more fragile than normal. The spleen appears to be in some way responsible for the increased fragility of the

cells, for, following its removal, the anemia is usually cured or greatly improved

Although no ill effects have ever been observed in man following removal of the spleen, yet from the results of experiments upon animals it is probable that in certain emergencies (e.g., hemorrhage, gas poisoning, or surgical shock) a person deprived of his spleen would not have quite as good a chance of survival as would one with his spleen intact.

#### The Lymph and the Lymphatic System

The formation and composition of lymph. The tissue fluid, as explained elsewhere, is derived from the blood plasma. A large part of the fluid which leaves the arterial end of the capillary is returned to the circulation through the capillary near its venous end But the tissue spaces, as well as being in relation to the blood capillaries, are also drained by a fine network of capillary vessels called *lymphatics*. Whereas the fluid reabsorbed by the blood capillaries is an aqueous solution of crystalloids (inorganic salts, sugar, urea, etc.), the fluid which passes into the lymph vessels contains some plasma protein as well Small colloidal particles, such as those of India ink and other dyes which may have entered the tissue spaces from the blood, are also taken up by the lymphatics Such particles do not return to the circu lation through the walls of the blood capillaries. The fluid in the lymphatic vessels is called lymph, but it is almost identical in composition with the tissue fluid Lymph contains the same constituents as are found in blood plasma, though in different concentrations. It con tains the three plasma proteins, but the total protein concentration is between 3 and 4 percent as compared with around 7 percent for plasma The calcium and total phosphorus are both lower than in the plasma, the other constituents (e.g., sodium, potassium, magnesium, chlorine, sugar, urea, etc ) are in about the same concentrations in the two fluids Lymph contains numerous lymphocytes and an odd granulocyte, but very few or no red cells, under normal circumstances The lymphocyte count varies considerably in different specimens of lymph from 1000 to 20,000 per cu mm

The lymph vessels (lymphatics). The lymphatic system commences at the periphery in the network of fine vessels just mentioned. Such vessels are found in the skin (see Fig. 3.26) and subcuttaneous tissues, in the fasciac of muscles, and in the connective tissue of the abdominal and thoracic viscera. Those in the villi of the small intestine are

called lacteals (L lac, milk) Through them a large proportion of the fat of the food is absorbed into the circulation, their name having been suggested by the milky appearance of their contents (usually referred to as the chyle) after a meal of fat The lymph capillaries join to form larger vessels which ulti mately form two main trunksthe thoracic duct and the right lymphatic duct The former receives all the lymph of the body except that from the right side of the head, neck and thoracie wall, right arm right lung, right side of the heart, and upper surface of the liver Lymphatics of these parts drain into the right lymphatic duct Each of these large vessels opens into the subclavian vein of the corresponding side of the body (see Fig 3 26) A section of a lymph ves sel is shown in Figure 3 27

The lymph nodes (or glands) At certain strategic points in the course of the medium sized lymph vessels are situated small ovoid or round structures composed of lymphoid tissue. These are the lymph nodes (Fig. 3.28). These bodies are composed of a network of delicate sepacked with masses of lymphocy.




Fig 3 26 On left above thorace duct (a) opening into left subclavian vein (b) (c) internal jugular vein on right above the main lymphatic vessels of the upper limb (d) lymph nodes Below, main lymphatics of the lower limb (f) lymph nodes in the groin (g) nodes (dotted outline) at the back of the knee

posed of a network of delicate septa, the spaces between which are packed with masses of lymphocytes Lymph vessels, usually four or five in number, enter the convexity of the node, they are called affer ent vessels. The lymph leaves the node by a single efferent vessel which emerges from the concavity or hilum of the node. The lymph in its course from the tissue spaces to the point where it is returned to the




Fig 3 27 A lymph vessel

blood must pass through the lymph nodes, which act, in a sense, as filters

In the upper part of the body, lymph nodes are situated in the subcutaneous tissue on the inside of the front of the elbow joint (Fig 3 26), in the armpit, behind the ear, and running down either side of the neck Those in the armpit and at the elbow drain lymph from the hand, those of the neck receive lymph from the head and throat A group of lymph nodes can be felt as hard shotlike objects in the groin, they receive lymph from the external genital organs and lower lumbs Another group of lymph nodes, situated behind the knee, drain lymph from the foot and leg Lymph nodes are also situated more deeply in the tissues of the neck and limbs, as well as along the course




Fig 3.28 Diagram of lymph node, sectioned to show internal structure, one area free of lymphocytes to show lymph spaces

of the lymph vessels draining the viscera. When the hand or foot is infected, the lymph glands at the elbow and armpit in the one instance, and at the knee or in the groin in the other, may become inflamed and swollen as a result of bacteria or their products which have been carried upward by the lymph current. In septic conditions of the scalp, throat, or ear, the nodes receiving lymph from the infected area become similarly involved. Within the nodes, bacteria are attacked by leucocytes and other phagocytic cells. The lymph nodes must therefore be looked upon as important elements of defense against the invasion of the blood by microorganisms traveling along the lymphatic channels. They constitute a second defense line, the phagocytic cells at the primary site of infection bearing the first shock of the attack.

Other functions of the lymph nodes. Besides acting as filters and thus playing a prominent role in the defenses of the body against the inroads of bacteria, the lymph nodes exercise two other highly important functions: (a) the production of lymphocytes, and (b) the manufacture of plasma protein (globulin). The nodes form plasma globulin through the disintegration of the lymphocytes of which they are largely composed. This protein (gamma globulin) of the plasma is closely associated with the immune bodies (antibodies) through which bacteria and their toxic products are destroyed and antidoted The disintegration of the lymphocytes within the lymph nodes and the production of antibodies appear to be controlled by the adrenocorticotrophic hormone of the pituitary body (Chap. 10). This immunizing function of the lymph nodes constitutes a third line of defense against infective agents.

These functions are not confined to the lymph nodes; the nodes possess such functions because they are composed of lymphoid tissue. The manufacture of plasma globulin and of lymphocytes are functions of lymphoid tissue in general—in the spleen, tonsils, thymus, Peyer's patches of the intestinal wall, etc.

Edema. An increase in the quantity of tissue fluid to the point where it causes a readily detectable increase in volume of a part is called edema. It is most frequently seen in the skin and subcutaneous tissues, about the ankles, or in the face below the eyes, as a symptom of heart or of kidney disease. When the edema is well marked, the skin appears "puffy," and, when one presses it with a finger, a dent or pit is left which takes a little time to level up again.

Edema is due to an imbalance of those factors regulating the interchange of fluids between the vessels and the tissue spaces (see p. 65). It may result, therefore, from any of the following causes: (a) increased capillary pressure, as in heart disease; (b) reduced plasma osmotic pressure, as in chronic kidney disease; (c) increased permeability of the capillary wall, as in acute kidney disease or as a result of certain poisons (e.g., histamine); (d) obstruction of the lymph channels (lymphatic edema).

Edema may be quite local and form small, well-defined, raised, pale areas in the skin Such occur in susceptible persons after eating certain foods (e.g., shellfish, strawberries, celery, etc.). The condition is called uricaria, and known popularly as hives.

## The Coagulation of Blood and Lymph

It blood is collected in a test tube it will be found after 5 or 6 minutes to have set into a jelly. The tube may be inverted, but the blood, which is now said to have clotted or coagulated, does not flow. When a section of this clot is examined under the microscope, it is found to be composed of a tangled mesh of very delicate fibrils among which are entrapped, as in a net, crythrocytes, leucocytes, and many fragmented platelets. The filaments are composed of fibrin, an insoluble gel (p 18) form of fibrinogen produced during the clotting process.




Fig. 3.29 Showing red cells, platelets, and fibrin threads in a blood clot.

They may be seen in many places to radiate from centers formed of platelets. If the clot is allowed to stand for a while, it undergoes shrinkage, and, as it shrinks, expresses from its meshes a clear, faintly straw-colored fluid This clear fluid is the serum (Pl. 1). The serum remains fluid indefinitely; it is quite incapable of clotting, for it contains no

fibrinogen Plasma separated from the blood cells by centrifuging, clots in a way similar to that of whole blood and expresses the clear serum The clot is white, since it contains no cells, but except for this difference it is identical with that formed in whole blood. The clotting process is essentially, therefore, a phenomenon of the plasma Lymph also clots, though somewhat more slowly and less firmly than does blood or plasma.

The clotting mechanism. Four substances are necessary for the coagulation of blood prothrombin thromboplastin calcium and fibrinogen Prothrombin gives rise to thrombin, an enzyme Fibrinogen, prothrombin, and calcium are present in circulating blood Thromboplastin (a lipid or fatlike compound containing phosphorus) is widely distributed throughout the tissues the lung and brain being especially rich in this factor. It is absent or present in only small quantities in blood plasma. When blood is shed, thromboplastin is liberated from injured tissue and probably also from the leucocytes of the blood itself. The thromboplastin, acting upon the prothrombin in the presence of calcium, in an ionized form converts it to the active thrombin. Thrombin acts in turn upon the soluble protein fibrinogen, convert ing it to insoluble fibrin which as mentioned above, is deposited as fine threads to form the framework of the clot (Fig. 3.29)

The foregoing is a description of the clotting mechanism in the simplest possible terms. The chief factors are summarized in the following scheme.

The mechanism of blood coagulation has been found to be most complex and several details have been omitted from the above scheme one of the more important of these concerns a globulin in the plasma which accelerates the conversion of prothrombin to thrombin This factor has been named plasma accelerator globulin or briefly plasma AC globulin

Blood does not clot in the living body because there is not sufficient free thromboplastin to convert the mactive prothrombin into the active thrombin Clotting within the vessels is further guarded against by the presence in the blood plasma of a substance which neutralizes the action of any thrombin which might be present This antagonist to thrombin is called the normal antithrombin of plasma. When, however, blood escapes from a cut vessel and flows from a wound, throm-boplastin is liberated from the damaged tissue cells and also from disintegrated platelets. Thus, the clotting process is immediately initiated.

Anticoagulants. Substances that prevent the coagulation of blood are called anticoagulants Among such are hiridin and heparin, dicoumarin, sodium or potassium oxalate, fluorides, citrates, certain neutral salts, e.g., sodium and magnesium sulphates. The oxalates act by precipitating the calcium of the blood which, as mentioned above, is required for the formation of thrombin. Fluorides and citrates form soluble compounds with the calcium, but the latter, in so far as the clotting process is concerned, is in an inactive form (un-ionized). Hruidin is a material secreted by the mouth glands of the leech. It prevents the blood of the parasite's host from coagulating while it is being sucked. Hirudin is prepared commercially for use, chiefly in physiological experiments, heparin, however, has largely taken its place.

Heparin (Gk hepar, liver) is an anticoagulant of special interest, since it is found in mammalian liver, lung, muscle, and other tissues. It is very powerfully anticoagulant, 1 mg of a highly purified preparation being capable of preventing for 24 hours the coagulation of 500 cc of blood, if kept at a temperature of about 50° F

Dicoumarin is a poison formed in spoiled clover silage. This material has been responsible in the past for a fatal hemorrhage disease of farm animals that have been fed on spoiled clover. Dicoumarin acts by reducing the prothrombin concentration of the blood to such a low level that clotting cannot occur. Prothrombin is produced in the liver, and it is probable that the poison acts upon this organ and interferes in some way with prothrombin production. This anticoagulant is now widely used in the treatment of coronary thrombosis, and other forms of intravascular clotting, for the purpose of preventing the spread of the clotting process.

A low prothrombin concentration of the blood with a tendency to bleed is associated in some animal species with a dietary lack of a vitamin found in certain green foods (e.g., alfalfa, spinach, cabbage, etc.) This is known as vitamin K, or the antihemorrhagic vitamin (see p. 397)

The concentration of prothrombin in the blood of the newborn child is frequently much below the normal level, due apparently to a lack of vitamin K in the mother's diet. It is unusual, however, for this

to be a cause of bleeding, because when the baby is fed, the bacteria introduced into the intestine with the food are capable of synthesizing the antihemorrhagic vitamin. Sometimes, however, the prothrombin level of the infant's blood is so low at birth that it does not rise to normal before a grave hemorrhagic state—hemorrhagic disease of the new born—supervenes. It is treated by the intravenous administration of synthetic vitamin K.

The essential importance of prothrombin in the clotting mechanism is clearly evident from the foregoing account of its presence in low concentration. We have also seen that after ionized calcium has been removed from the blood by oxalate, or other means, clotting is, entirely prevented, since prothrombin cannot be activated to thrombin.

The indispensability of the two other primary factors in the clotting process, namely, thromboplastin and fibrinogen, may be illustrated by the following observations. Thromboplastin, as mentioned above, is derived from tissue cells and from damaged cells in the blood itself. If the blood is drawn by means of a syringe directly from a blood vessel into a clean glass container made smooth by a coating of liquid paraffin or of a solution of silicone, the cells remain largely intact. For this reason, and because the blood does not come into contact with the tissues, it receives a minimum quantity of thromboplastin and, as a consequence, clots much more slowly than usual. The experiment of the "living test tube" also demonstrates the indispensability of thromboplastin for the clotting of blood. When a short section of the jugular vein in the living animal is isolated by ligatures from the rest of the vessel and removed, the blood within it, since it remains undisturbed and in contact with the smooth and uninjured lining of the vein's wall, does not clot for many hours.

Defibrinated blood is blood from which the fibrinogen has been removed as fibrin; it is incapable of forming a clot. Defibrination is carried out by whipping the blood with a bundle of thin twigs or wires. As clotting occurs, the fibrin collects upon the twigs and can in this way be removed, leaving only the serum and cells.

The mechanism underlying the clotting of the blood is chemical in nature, and, like other chemical reactions, is slowed by lowering the temperature and hastened by warming Blood placed in the refrigerator immediately after it has been shed may remain fluid for 2 or 3 hours.

Means used to hasten coagulation. The coagulation of blood as it issues from a wound can be accelerated by several measures. Heat, powder dusted on the wound, tissue extracts (containing thrombo-

plastin), a preparation of thrombin, and various chemicals (known from their action in this regard as styptics or hemostatics) accelerate the clotting process. Among the latter are alum, ferric chloride, zinc chloride, and silver nitrate Such substances arrest bleeding in two ways, they hasten clotting of the blood through the precipitation of the plasma proteins which, by forming a sticky mass, mechanically impede the flow of blood and thus favor platelet disintegration, they also cause constriction of the walls of the small bleeding vessels. Cer tain physiological conditions hasten coagulation. Blood shed after muscular exercise or emotional excitement clots more quickly than usual owing to the discharge of adrenaline into the circulation (p 420) Adrenaline (epinephrine) injected into the body also causes a sample of blood drawn shortly after the injection, to clot more quickly, but no such effect upon the clotting mechanism results from adding adrenaline to blood outside the body 12 or by applying it to a bleeding vessel In the latter instance, however, adrenaline tends to stop the bleeding by constricting the small vessels with which it comes into contact. Thus, superficial bleeding or slow oozing of blood is very quickly arrested. The arrest of bleeding either by hastening coagula tion of the blood or by constriction of the bleeding vessels is called hemostasis

Coagulation time and bleeding time. The coagulation time is defined as the length of time in minutes, measured from the moment the blood sample is collected to the appearance of the first definite sign of clotting Several methods for determining this interval have been devised. The simplest is that of Howell, which consists in collecting the blood in a small test tube coated inside with liquid paraffin and gently tilting the tube every 20 seconds or so until the blood is no longer liquid—that is, when the line of its surface no longer changes freely with tilting

The bleeding time is measured by pricking the base of the finger nail and when a drop of blood appears, dabbing it lightly from time to time with a small piece of filter paper. The time from the appearance of the drop of blood to when the filter paper is no longer stained is taken as the bleeding time.

<sup>1°</sup> The fact that adrenaline histens the congulation of the blood if injected into the body but not when added to blood in a test tube strongly suggests that it acts not directly upon the clotting mechanism but only through some other a ructure, possibly a ductiess gland causing it to secrete a principle which exerts a direct congulant action.

## Hemorrhagic States Due to Failure of the Clotting Mechanism


In rare instances a deficiency of fibrinogen is the cause of the blood failing to form a firm clot, with the result that severe hemorrhage may follow a wound which in a normal person would be of little con cern It is a very common practice to give calcium in the belief that it will promote coagulation of the blood in hemorrhagic states. It appears, however, that the blood is never so low in calcium that clotting fails Contrary to what might be expected, a hemorrhagic tendency is never due to a low concentration of calcium in the blood Certain serious effects (e g, tetany) ensue before the blood calcium is reduced to the point where the clotting mechanism is interfered with The common use of calcium to promote the coagulation of the blood in hemorrhagic states is, therefore, without any scientific foundation

A depression of the prothrombin of the blood (hypoprothrombinemia) is one of the commonest defects of the clotting mechanism The tendency to bleed in obstructive jaundice and, as mentioned on page 104, in the hemorrhagic state of cattle fed upon spoiled clover, as well as in the hemorrhagic disease of the new born, is due to this cause

In obstructive jaundice, bile, which is an essential for the absorption of vitamin K, does not reach the intestine. The body fails, therefore, to secure the vitamin required for the manufacture of prothrombin

Hemophilia. Hemophilia is a disease in which the blood takes an abnormally long time to clot The blood of a normal person clots within 5 or 6 minutes after its withdrawal from the body. In hemo philia the blood may remain fluid for an hour or more There is grave danger, therefore, of a person afflicted with this disease bleeding to death, even from a trivial wound Hemophilia is hereditary but it occurs almost solely in males and is transmitted only by females. This sex linked type of heredity is also seen in the case of color blindness

The manner in which the disease is carried from generation to generation may be illustrated by an example. If a man suffering from hemophilia-commonly known as a bleeder-marries a normal woman, no offspring of the union will show the disease Some of the daughters, nevertheless, but not the sons will when they marry, transmit the disease to some of their sons, but not to the daughters The latter, however, can again transmit it The disease, therefore, skips a generation, a bleeder inheriting it from his mother's male forebears.



B means a bleeder; O means not a bleeder, T means a transmitter but not a bleeder.

What part of the clotting mechanism is at fault in this disease is uncertain, but there appears to be some defect in the production of an active thromboplastin in the plasma. The coagulation time in hemophilia is greatly prolonged. Bleeding due to this disease is treated most successfully by blood transfusions.

Purpura. Purpura is another hemorrhagic disease, in which blood leaks from the small vessels of the skin and mucous membranes. When the hemorrhages are from the cutaneous vessels, the skin is discolored by small purplish patches. Some defect in the capillary walls and a reduction in the number of platelets appear to be the chief factors concerned in the production of the disease. The coagulation time is within normal limits but the bleeding time is prolonged. One severe type of purpura is apparently due to some destructive action of the spleen upon the platelets. Removal of the spleen is followed in most instances by an increase in the platelet count and a cessation of the hemorrhages.

# The Regulation of the Reaction of the Blood and Other Body Fluids

The acidity or alkalinity of a solution is dependent upon its concentrations in hydrogen ions (H<sup>+</sup>) and hydroxyl ions (OH<sup>-</sup>), that is, upon the degree to which the molecules in the solution are dissociated into their constituent ions. A tenth normal solution of hydrochloric acid (H<sup>+</sup>Cl<sup>-</sup>), for example, contains a high concentration of hydrogen ions, whereas the hydrogen ion concentration of a tenth normal solution of acctic acid is very much lower (see Table 1-3, p. 13). The acid properties of hydrochloric acid are, therefore, much greater than those of acetic acid. Similarly, a tenth normal solution of sodium hydroxide (Na<sup>+</sup>OH<sup>-</sup>) has a much higher concentration in hydroxyl

ions than has a tenth normal solution of sodium bicarbonate. The former solution is therefore more alkaline. It must be emphasized that it is the concentration of the hydrogen ion—i.e., the extent to which the molecule is dissociated—and not the total number of hydrogen atoms which determines the acid reaction. Thus, HCl though it contains only one hydrogen atom in its molecule undergoes almost complete dissociation into H+ and Cl- and is, in consequence, a much stronger acid than acetic, whose molecules dissociate to a relatively small extent. Water undergoes dissociation to a very slight degree into H+ and OH-. The ionized hydrogen in a liter of water is 0.0000001 gram, or more conveniently expressed,  $1 \times 10^{-7}$  gram. We therefore say that the H ion concentration of water is  $1 \times 10^{-7}$  gram per liter. The symbol  $^{-7}$  to the right of the figure 10 is termed the negative exponent or index, and means that to express the value to the right of the decimal fraction the figure 1 must be placed 7 places to the right of the decimal point (0 0000001). Expressed as a common fraction the 10<sup>-7</sup> would be 10000000 gram per liter.

Other examples of this system of notation are:

10<sup>-1</sup> means 0.1 or  $\frac{1}{10}$ 10<sup>-2</sup> " 0.01 or  $\frac{1}{100}$ 10<sup>-3</sup> " 0.001 or  $\frac{1}{1000}$ , and so on


The concentration of OH ions in water is also  $1\times 10^{-7}$  gram per liter. That is, the H+ ions and OH- ions are in equal concentration; water is, therefore, neutral in reaction. Now the product of the concentrations of H ions and OH ions in water is  $1 \times 10^{-14} [(1 \times 10^{-7})]$  $\times$  (1  $\times$  10<sup>-1</sup>) = 1  $\times$  10<sup>-14</sup>]. Furthermore, in any aqueous solution, whether acid, neutral, or alkaline, the product of the concentrations of H<sup>+</sup> and OH<sup>-</sup> ions is constant at 1 × 10<sup>-14</sup>. This means that in an acid solution the H+ ion concentration increases, whereas the OHion concentration is reduced reciprocally. It is evident, then, that in order to express the reaction of a solution it is not necessary to give the concentrations of both H+ and OH- ions, but only of the H+ ion. The reaction of water or any other neutral solution is indicated, therefore, by stating that it has an H+ ion concentration (usually designated cH) of  $1\times 10^{-7}$ . The hydrogen ion concentrations of alkaline solutions are less than  $1\times 10^{-7}$ —that is,  $1\times 10^{-4}$ ,  $1\times 10^{-4}$ ,  $1 \times 10^{-1}$ , and so on, according to the degree of alkalinity. The hydrogen ion concentrations of acid solution are more than  $1 \times 10^{-7}$ ; i.e., 1 × 10<sup>-4</sup>, 1 × 10<sup>-4</sup>, 1 × 10<sup>-4</sup>, etc.

To summarize: neutral solutions have 0.0000001 gram of ionized hydrogen per liter—i e., a cH of  $1 \times 10^{-7}$ .

Alkaline solutions have a lower H<sup>+</sup> ion concentration—i.e., from  $1 \times 10^{-7}$  to  $1 \times 10^{-14}$ .

Acid solutions have a higher  $H^+$  ion concentration—i.e., from  $1 \times 10^{-7}$  to  $1 \times 10^{-2}$ .

Within recent times a more convenient method of expressing the hydrogen ion concentration has been introduced, in which the symbol pH, instead of cH, is used. Thus the expression cH × 10<sup>-3</sup>, indicating a neutral solution, becomes pH 7. Alkaline solutions have pH's ranging from pH 7 to pH 14; acid solutions from pH 7 to pH 1. It will be observed that the lower the pH number, the higher is the hydrogen ion concentration—i.e., the less alkaline or the more acid is



the solution (Fig. 3.30) More advanced texts should be consulted if the reader wishes to understand the mathematical basis for this method of expressing the hydrogen ion concentration in terms of pH.

Arterial blood has a pH of between 7.39 and 7.45. Venous blood is very slightly less alkaline, having a pH between 7.37 and 7.42; the change is due to carbon dioxide and small quantities of lactic acid which the blood absorbs in passing through the tissues. Though large quantities of acid (carbonic, lactic, sulfuric, hydrochloric, etc.) are constantly being produced in the body as a result of various metabolic processes, the reaction of the blood and other body fluids remains remarkably constant around the figures just given. The three important mechanisms responsible for the maintenance of this constancy of reaction are:

- 1. The elimination of CO2 by the lungs (p 247).
- 2. The excretion of acid by the kidney [the urine is decidedly acid in reaction (p. 275)]
  - 3. The buffer systems of the blood and other body fluids

Buffers may be defined as chemical substances which when pres-

ent in a solution permit no change, or only a slight one, to occur in that solution when acid (or alkali) is added to it. An amount of hydrochloric acid which if added to water would cause a pronounced increase in acidity would, when added to blood or any solution buffered against acid, produce no appreciable change in reaction. It is from this action in buffering the shock, one might say, of the added acid that these substances derive their name. For example, when hydrochloric acid is poured into a solution of sodium bicarbonate the very weak carbonic acid together with sodium chloride, a neutral salt, is produced. Furthermore, the carbonic acid is volatile; it escapes into the atmosphere. The result is that little change in the reaction of the solution occurs, even though it has received relatively large amounts of the strong acid. The sodium bicarbonate acts as a buffer.

The chief buffer of the plasma and tissue fluids is sodium bicarbonate. A fixed acid such as lactic, formed during muscular activity, reacts with the bicarbonate as shown in the following equation.

The sodium bicarbonate of the plasma thus serves as a store of alkali available for the protection of the body against poisoning with the acid products of metabolism. It is therefore called the alkali reserve.

Other buffers of the blood are the plasma proteins (which form proteinates), phosphates, and the hemoglobin within the red cell. Hemoglobin is a weak acid; it is combined in the red blood cells with alkali (potassium) to form potassium hemoglobinate (KHb). This alkali acts to buffer carbonic acid entering the blood from the tissues. The contents of the erythrocytes, however, are separated from the plasma by a semipermeable membrane which will not permit the passage of K+, Na+, or Ca++ ions but allows the free passage of Cl-, HCO<sub>1</sub>-, and other negative ions, as well as the H+ ion and CO<sub>2</sub>. When, therefore, the carbon dioxide produced in the tissues enters

112

the plasma, the following interchange of ions occurs across the crythrocyte boundary. The carbon dioxide (CO<sub>2</sub>) diffuses freely into the cell and, through the action of an enzyme called carbonic anhydrase, combines rapidly with H<sub>2</sub>O to form carbonic acid (H<sub>2</sub>CO<sub>2</sub>). The carbonic acid, being a stronger acid than reduced hemoglobin, seizes the base combined with the pigment to form potassium bicar-




Fig 3.31 Showing main processes occurring in the blood during the output of carbon dioxide and uptake of oxygen in the lung (After Roughton, modified)

bonate (KHCO<sub>2</sub>) which dissociates into K<sup>+</sup> and HCO<sub>3</sub><sup>-</sup> ions. The membrane of the cell, as just mentioned, is permeable to HCO<sub>4</sub> ions but not to K<sup>+</sup> ions HCO<sub>3</sub><sup>-</sup> ions diffuse across the cell boundary into the plasma; the K<sup>+</sup> ions are retained In order to maintain the balance between negative and positive ions on the two sides of the cell membrane, Cl<sup>-</sup> ions derived from the Na<sup>+</sup>Cl<sup>-</sup> of the plasma pass into the cell in exchange for the HCO<sub>3</sub><sup>-</sup> ions which pass out. In the plasma, the HCO<sub>3</sub><sup>-</sup> ions unite with the Na<sup>+</sup> ions, which had been coupled with Cl<sup>-</sup> ions, thus forming sodium bicarbonate (Na<sup>+</sup>HCO<sub>3</sub><sup>-</sup>). The mechanism just described is termed the chloride shift (see Fig. 3.31). This mechanism is responsible for the phenomenon, now well recognized, that an increase in the carbon dioxide of the plasma causes a rise in the alkali reserve (sodium bicarbonate). In the lungs, ionic interchanges in the reverse order occur. The hemoglobin when it be-

comes oxygenated is a very much stronger acid than is reduced hemo globin or carbonic acid. It, therefore, recaptures the alkali (K\*) bound to carbonic acid. The carbonic andydrase, whose action is now reversed, a decomposes the carbonic acid into carbon dioxide (CO<sub>2</sub>) and water (H<sub>2</sub>O). The carbon dioxide diffuses out of the cells, as they traverse the capillaries of the lungs, and is exhaled. The concentration of HCO<sub>2</sub>—ions within the cells is reduced as a result of these reactions HCO<sub>4</sub>—ions pass from the plasma into the cells and Cl—ions are transferred from the cells to the plasma. The sodium bicarbonate of the plasma is reduced, the sodium chloride increased, by the interchange

Acidosis and alkalosis. The proportion of carbonic acid to sodium bicarbonate in the plasma is about 1 to 20, thus

$$\frac{\mathrm{H_2CO_3}}{\mathrm{NaHCO_3}} = \frac{1}{20}$$

It is upon this ratio that the reaction of the blood largely depends Any shift in the ratio causes a change in blood reaction If, for example, the ratio should increase, say, to  $\frac{1}{25}$ , the blood would be more alkaline If the ratio should be reduced to  $\frac{1}{15}$ , the blood would be less alkaline In health, the ratio remains virtually constant. The several mechanisms mentioned above ensure this constancy. If, for example, there should be a tendency for the ratio to rise, as a result of an increased production of carbon dioxide, the bicarbonate of the plasma increases through the chloride shift mechanism. If, on the other hand, the bicarbonate is reduced owing to the accumulation of fixed acids (see equations, p. 111), more carbon dioxide is excreted by the lunes and the ratio thus maintained at its normal value.

In certain diseases, e.g., diabetes and kidney disease (nephritis), excessive acid production or reduced elimination of acid occurs, the bicarbonate of the plasma becomes markedly reduced, and a change in the ratio may result. The blood becomes less alkaline and the alkali reserve is diminished. The condition is then spoken of as acidosis. The term acidosis is also applied to the condition in which the plasma bicarbonate is reduced but there is no change in the H-CO\_NaHCO\_s ratio and, consequently, no change in blood reaction. This is spoken of as compensated acidosis. When, as in the first mentioned instance, the acidosis is accompanied by a change in blood reaction it is said to be uncompensated.

It should be emphasized that the H2CO2/NaHCO2 ratio does not

12 The action of most enzymes is reversible

#### 114 THE BLOOD, LYMPH, AND TISSUE FLUIDS

vary from its normal value of \$\frac{1}{20}\$ unless serious disease exists. There are two possible exceptions to this statement. First, in very strenuous muscular exercise the large quantities of carbon dioxide and lactic

acid produced might result in a slight and temporary reduction in the alkalimity of the blood Secondly, forced breathing (p 255) if prolonged may, through the excessive loss of carbon dioxide, cause the blood to become temporarily slightly more alkaline

It should be remembered moreover, that the blood practically never, except in the last stages of some fatal disease, becomes actu ally acid in reaction-that is, with a pH less than 7, and rarely does it even approach the neutral point. The range of blood reaction compatible with life is from about pH 7 8 to 6 8. It is quite evident then, that the term acidosis is frequently used-especially by the advertisers of quack remedies-without any clear understanding of its meaning

Alkalosis is the corresponding term used to denote an increase in sodium bicarbonate with or without a change in the H.CO./NaHCO. ratio 14 As in the case of acidosis it is called compensated or uncompensated respectively, according to whether or not a change in the ratio and so, of course in the blood reaction has occurred Alkalosis may result from the ingestion of large quantities of sodium bicar

bonate, from persistent vomiting when hydrochloric acid is removed in the gastric juice, or from forced breathing when, as already men tioned unusually large quantities of carbon dioxide are eliminated 14 It has been the custom to use the terms acidosis and alkalosis to denote changes in the alkali reserve regardless of whether or not an accompanying change occurred in pH But with a view to more precise definitions it has been suggested that the

above terms be reserved for changes in alkalı reserve in which the pH is normal and that acidemia and alkalemia be used to denote changes in alkali reserve with a fall or a rise respectively in pH.

# The Circulation of the Blood

# General Description of the Circulatory System

The blood circulates through the body in a completely closed system consisting of a pump—the heart—and a network of tubes—the blood vessels.

The heart. The heart is a hollow muscular organ, roughly con shaped, situated near the center of the thoracic cavity, and in closs relation to the lungs. About a third of its bulk lies to the right and two thirds to the left of the mid-line of the body. The heart is divided by partitions or septa into four chambers—the right and left auricle. (or atria) and the right and left ventrucles. The auricles are situated above, posteriorly, and to the right; the ventrucles are situated below more anteriorly, and to the left. The broad upper part of the hear formed by the auricles is called the base. Its somewhat pointed lower part formed by the left ventricle is called the apex; this is directed downward, forward, and to the left (Fig. 4.1).

The heart is enclosed within a delicate two-layered membrane called the sepous pericardium; each layer consists of a single stratum of endothelial cells. The inner layer, which is in contact with and adherent to the heart muscle, is called the visceral layer or epicardium. The outer layer is separated from the inner by a thin film of fluid and is known as the parietal layer. The potential space between the two layers is known as the pericardial sac. A tougher fibrous membrane—the fibrous pericardium—surrounds the serous pericardium and extends upward to blend with the coats of the great vessels; it is firmly

<sup>1</sup> Auricle, though an older anatomical term, is still most frequently used in

physiology.

311

attached in front to the posterior aspect of the sternum and below to the central part of the upper surface of the diaphragm

The right and left halves of the heart—the right auricle (or atrium) and ventricle, on the one hand and the left auricle (or atrium) and ventricle, on the other—are not in communication. The partition in terposed between the two auricles is called the interauricular septum.




Fig 4.1 Showing the position of the heart in relation to the chest wall and lungs. The black area indicates that part of the heart uncovered by the lungs.

that separating the two ventricles from one another is named the interventricular septim. The auricle of each side communicates with the corresponding ventricle through an opening the auriculo (or atrio-) ventricular orifice guarded by the auriculo (or atrio-) ventricular valve. These valves open toward the ventricles permitting blood to enter from the auricles but close the instant that the ventricle con tracts thus the passage of the blood backward into the auricles is prevented. The auriculo- (atrio-) ventricular valve on the left side is also known as the mitral valve from its fancied resemblance to a bishops miter or tall cap that on the right side as the trictuspid valve (see also p. 174) because it consists of three triangular leaves or cusps

The muscular walls of the auricles are thin compared with the muscle of the ventricles. The left ventricular wall has a wall about double the thickness of that of the right ventricle, since the work which it must perform is much greater than that performed by the




Fig 4.2 Cross section of heart through the ventricles.

right ventricle (Fig. 4.2). The heart chambers are lined by a delicate endothelial membrane called the endocardium Springing from the wall of each ventricular cavity are two small muscular pillars, somewhat conical in shape and known as the papillary muscles (musculi papillares). From the apices of the latter, slender cords, the chordae tendineae, arise and are attached to the ventricular aspects of the auriculoventricular valves (Fig. 4.3). Acting like guy ropes, they support the delicate leaflets of the valves against the pressure developed in the ventricle during its contraction

The arrangement of the fibers of the ventricles is very complex. They are disposed in a superficial and a deep series of layers. Of the superficial layers, some course from the upper and anterior part of the right ventricle downward and to the left, looping around the apex, in the form of a vortex (Fig. 4.4). Passing upward and inward they end in the papillary muscles of the left ventricle. Other superficial groups of fibers pass diagonally across the posterior surface of the heart from right to left. Of the deep layers, some pass from the papillary muscles of one ventricle to those of the other, encircling both ventricles in a scroll-like or S-shaped manner. Other groups encircle the left ventricle alone.

The right side of the heart contains venous blood; that is, blood which has passed through the tissues and given up a proportion of its

load of oxygen (p. 247). The left side contains bright red (arterial blood—blood which has been oxygenated in the lungs. The venous blood is delivered to the right auticle by two large veins—the superior vena cara which drains blood from the upper part of the body (head,




Fig. 4.3 Showing the heart and great blood vessels; interiors of right auricle and ventricle exposed. The right auriculoventricular valve (tricuspid) is shown.

neck, arms, and the interior of the thoracic wall), and the interior vena cava which conveys blood from the lower part of the body (lower limbs and abdominal organs) The blood flows from the right auricle through the auriculoventricular orifice, into the right ventricle, from which it is discharged during ventricular contraction into the pulmonary ariery and thence through the vessels of the lungs. After passing through the capillaries of the lungs, where it gives up carbon dioxide, and absorbs, oxygen, the oxygenated blood is conveyed by the four pulmonary veins (two from each lung) to the left auricle. It



Fig 44 Dissection of heart muscle to show the course of the fibers (Redrawn, after Mall)

then passes through the left auriculoventricular orifice into the left ventricle. When the ventricular muscle contracts, the blood is pumped into the aoria—the great artery which arises from the upper part of the left ventricle.

Thus, the blood circulates through the lungs from the right ventricle to the left auricle, and through the rest of the body from the left ventricle to the right auricle (Pl 4) The course through the lungs is called the pulmonary or lesser circulation, that through all other parts of the body, the systemic or greater circulation. The right ventricles



Fig 4.5 William Har-

tricle is therefore the pump for the pulmonary circulation, the left ventricle serves a corresponding function for the systemic or greater circulation <u>Both ventricles contract at the same instant</u>

Though it is now a matter of general knowledge that the blood circulates in this way, the truth lay hidden until William Harvey, an English physician of the segenteenth century (Fig 45), made the discovery By his clear reasoning from a number of observations and simple experiments, he convinced those who were not blinded by pre




Fig 46 From an engraving in Harvey's Exercitatio anatomica de motil cordis et sanguinis in animalibus illustrating one of his demon strations (See text)

conceived notions that the blood must take the course outlined above Harvey pointed out that, owing to the structure and arrangement of the valves of the heart, the blood must course through it only in one direction. He showed that when the veins of a limb were compressed the blood was dammed back on the side of the block farther from the heart (Fig. 4.6). These facts could mean only that the blood made a circuit of the body. An examination of the valves of the veins which were so placed that they would allow blood to flow toward the heart, but not in the reverse direction, supported this conclusion. In 1628, he published his great work Evercitatio anatomica de motu cordis et sanguinis in animalibus (Anatomical Exercise on the Motion of the Heart and Blood in Animals). Before Harvey's day the dogmas of Galen, a Greek physician who practiced medicine in Rome in the

second century of the Christian era, dominated all medical thought. His views on physiology were accepted without question as the infal-lible pronouncements of a god. According to Galen, the food material lible pronouncements of a god. According to Galen, the food material absorbed from the intestines was conveyed by the portal vein to the liver where it was converted into blood. This newly formed and somewhat crude type of blood then passed to the right side of the heart where it was purified; the impurities, finding vent through the pulmonary artery and lungs, were exhaled in the breath. The blood formed in the liver did not circulate, but was conceived as simply ebbing and flowing in the veins; from it the tissues derived their nutriment. A small part of the blood in the right heart was believed, however, to find its way through invisible pores in the septum to the left side where, mixing with air received from the lungs through the pulmonary weins it was transformed into a more refined type of blood.

side where, mixing with air received from the lungs through the pulmonary veins, it was transformed into a more refined type of blood. This arterial blood, as the result of the action upon it of a vital essence—the pneuma—received from the air, possessed a life-giving principle which Galen called the vital spirit.

This brief description of ancient and medieval physiology will give the reader an appreciation and idea of the fundamental nature and immense importance of Harvey's discovery. Galen was a great and wise physician and experimentalist of his time and it can be seen that in his theory of the pneuma, by which he sought to explain the difference between venous and arterial blood, he was groping for an essential life-sustaining principle in air which we now identify as oxygen. Indeed, with the clear vision of genus, he remarked, "When we know what supports a flame we shall know the cause of the body's heat."

Oxygen was not discovered and its importance to living processes recognized until Lavoisier's work in the eighteenth century.

The blood vessels. There are four main types of blood vessels arteries, arterioles, capillaries, and veins.

arteries, arterioles, capillaries, and veins.

The arteries. These are of various sizes; the largest in the body are the aorta-which is the sole outlet for the blood from the left ventricle —and the pulmonary artery, which receives the contents of the right ventricle. In man, each of these vessels is a little over an inch in diamventrice. In main, each of mess vessels a fatte over all first in diameter. Traced peripherally, the channels of the arterial system divide and subdivide extensively like the limbs and twigs of a tree. The aorta ascends from its origin for a short distance and then, forming an arch, descends in front of the vertebral column through the thorax and abdomen. At the level of the fourth lumbar vertebra it divides into two vessels—the right and left common iliac arteries. Each common iliac divides into two branches—the external and internal iliac arter-




Fig 47 Dagram of man vessels of the body Veins and pulmonary artery in hatched shading arteries stippled I right ventricle Ia right auricle left ventricle 3 arch of aorta 4 innominate artery 5 right subcla vian artery 6 left common caroud ar tery 7 right common carotid artery 8 r ght subclavian artery 9 pulmonary artery 10 superior vena cava 11 jugular vein 12 left subclay an vein 13 abdominal aorta 14 branch es of celiac axis artery 15 hepatic veins 16 inferior vena cava 17 renal arteries and ve ns 18 superior mesen teric artery 19 inferior mesenteric artery 20 common iliac veins 21 common il ac artery A A adrenal glands D D daphragm K, k kid

ies. The former carries blood to the lower limbs after entering the thigh it is called the *femoral artery*. The internal iliac (or *hypogastric*) artery is distributed to the pelvis.

The thoracic and abdominal viscera are supplied by branches that spring from the aorta in its course through these cavities. Three large arteries spring from the arch of the aorta the left common carotid the left subclavian and the innominate arteries. The last is a short trunk which divides into the right common carotid and the right sub-

clavian arteries. The common carotid arteries are the main arterial trunks to the head. They ascend on either side of the neck and at the trunks to the need. They accend on either side of the need and at the upper border of the thyroid cartilage divide into the external and internal caronal arteries. The latter enters the crannal cavity where it divides into branches to supply the brain with blood. The subclavian artery conveys blood to the upper limb, at the root of the limb it is called the axillary artery, and in the arm the brachial artery (see Figs 4 3 and 4 7). It will be noted that the common carotid and subclavian arteries on the right side spring from the innominate, whereas the left common carotid and subclavian arise directly from the aorta

The foregoing is but the barest outline of the arterial system.

The walls of the arteries are thick and strong they contain a large proportion of elastic tissue and a smaller amount of smooth muscle. They are lined by endothelium which is called the internal coat or




Fig. 4.8. Cross sections of a small artery and vein showing the three coats. Note that though the vein is larger it has a thinner wall.

tunica intima. The middle coat or tunica media is made up of elastic and muscular tissues. The outer coat, composed of connective tissue, is called the tunica adventitia (see Fig. 4.8). The proportions of smooth muscle and elastic tissue in the arterial wall vary with the size of the artery Though elastic tissue enters largely into the construction of the largest vessels (e g, the pulmonary artery and aorta) muscular

tissue is relatively small in amount. In the medium and smallest arterial branches, especially in the latter, the smooth muscle is more abundant, the elastic tissue being correspondingly reduced. The walls of the arteries, except those of the smallest (1 mm or less in diameter), are themselves furnished with minute blood vessels. These vessels, called vaxa vasorum (vessels of vessels), ramify in the tunica adventitia and outer layer of the tunica media. The small arteries, as well as the vasa vasorum, are supplied with nerves through which their calihers are controlled.

The arterioles. Each of the smallest arterial branches divides into a number of still finer vessels, called arterioles. The arterioles are about 0.2 mm in diameter, being just visible to the naked eye. Although they possess an endothehal lining and a thin layer of elastic tissue, their walls are composed mainly of smooth muscle whose fibers form encircling rings. The arterioles are supplied with nerves which, by bringing about contraction or relaxation of the rings of smooth muscle, control their calibers. Thus, the quantity of blood passing through this part of the circulatory system may be altered from time to time as occasion requires (see pp. 195-6).

The capillaries. An atteriole when traced peripherally is found to break up into a number of extremely narrow tubes, the capillaries.



Fig 4.9 A capillary, its wall is composed of a single layer of endothelial cells.

These vessels (Fig. 49) are from ½ to 1 mm long, and have a caliber which in many instances is no greater, and may be less, than the diameter of a red blood corpusele (7 to 8 microns) The wall of a capillary is composed of a single layer of endothelial cells—a membrane of exquisite thinness which, as mentioned elsewhere (p 66), permits the free passage of water and crystalloids between its cells, but is only slightly permeable to the plasma proteins In amphibians, but not in mammals, there are to be seen, lying close to or in contact with the capillary wall, peculiar cells whose slender processes form a lattice-




Fig. 4.10 Showing Rouget cells on the wall of an amphibian capillary.

work embracing the vessel (Fig. 4.10). They are called Rouget cells. The capillaries, though constituting such an inconspicuous part of the circulation, are nevertheless the raison d'être for the rest of the circulatory system. The heart is the pump and the relatively thick-walled arteries and veins simply conduits which carry blood to and from the capillaries; in the capillary region the object of the circulation—namely, to convey oxygen and nutritive materials to the tissues and to carry carbon dioxide and waste products away—is fulfilled. The walls of vessels larger than the capillaries are impervious to gases and liquids.

The veins. Like the arteries, the veins have walls composed of three coats, but the tunica media is poorly developed as compared with an artery of about the same size. The wall of a vein is, consequently, much thinner than that of its companion artery, but its caliber is considerably larger (see Fig. 4.8). The larger veins of the abdomen and lower limbs possess valves which open in the direction in which the blood is flowing—that is, toward the heart (Fig. 4.11). The very small veins succeeding the capillaries are generally referred to as venules.




Fig. 4.11 A vein of the lower limb opened to show valve. The arrow indicates the direction of the blood flow.

### Hemodynamic Principles

In order that certain features of the circulation may be understood, some of the physical laws governing the flow of liquids through tubes must be briefly reviewed. The more important of these laws may be illustrated by means of a model

In Figure 4 12 is shown a piece of apparatus made to represent the systemic circulation and consisting of two horizontal glass tubes, AA and BB, from each of which a series of vertical side tubes of smaller caliber, T, T T, and T' T', T', T', are given off. The two systems of tubes are connected by finer tubes of elastic rubber. The hand bulb, H, is so valved as to direct a flow of fluid as indicated by the




Fig 4.12 Circulation scheme A model composed of glass and rubber tubes and a hand bulb. H to illustrate certain facts in the circulation of the blood (Description in text.)

arrows. It represents the left ventricle of the heart; the tube, AA, represents the aorta, and the vertical tubes, T, T, and T, represent some of its main branches. The system on the right corresponds to the venous system, in which BB are the larger veins (venae cavae) opening into the right auricle, Au, and the vertical tubes T', T', and T', their larger tributaries. The smaller rubber tubes, C, connecting the two systems represent the minute vessels, arterioles, capillaries, and venules. When the bulb, H, is alternately compressed and released at a suitable rate, it acts as a pump and, drawing fluid from the part of the system representing the auricle, Au, and the large veins. BB. forces it along AA. The fluid rises in the vertical side tubes, T, and its height in any tube indicates the pressure in centimeters of water at the point in the horizontal tube where that tube is given off. At every beat of the pump, the fluid in each side tube will be seen to fluctuate above, and then fall below, a certain mean level." It will also be observed that the pressure (height) of the fluid in the side tubes falls progressively from tube to tube, as indicated by the slanting lines, R-R and O-O. This slope of pressure is due to the loss of energy caused by the frictional resistance offered to the flow of fluid along the horizontal tubes; for with a given amout of energy (transmitted to the fluid through the hand bulb), the greater the proportion of that energy which is expended in driving the fluid against resistance along the horizontal tubes, the smaller will be the proportion exhibited as pressure. The greater the area of the frictional surface over which the fluid flows-that is, the longer the distance along the horizontal tubes-the greater will be the energy expended in overcoming frictional resistance and dissipated as heat. Hence, the pressure in the second tube is less than in the first, and the pressure in the third tube is less, again, than in the second; and, if there were a long series of tubes given off from the horizontal tube, the pressure in each would be lower than that in the one before it to the end of the series.

The section of connecting tubes, C, beneath the bar, D, can be compressed or decompressed by tightening or loosening the thumb screws marked S, S; the calibers of these tubes are thus reduced or increased, respectively. When their calibers are reduced and the quantity of fluid passing through them is, in consequence, diminished, but the amount of fluid entering the system from the pump remains

<sup>&</sup>lt;sup>2</sup> The model is not designed to illustrate the action of the heart, but only to show the physical principles governing the flow of liquid in narrow tubes. The ventricles, for example, do not "draw" blood from the auricles as the model indicates. On the contrary, the blood flows from the auricles into the ventricles under its own pressure.

approximately the same, then the pressure rises, as indicated by the heights of the fluid columns in the system on the left representing the arterial system. The pressure in the right-hand system, which represents the veins, falls, because less fluid now enters this system. Upon decompression of the rubber tubes, C—the quantity of fluid forced in by the pump again remaining unchanged—the pressure falls in the left-hand system and rises in the system on the right.

As the reader will learn later (pp 195 and 196), the arterioles can be narrowed or enlarged in caliber by means of vasoconstrictor and vasodilator nerves supplying their muscular coat. The bar, D, in the model, by the tightening or loosening of which the small rubber tubes can be compressed or decompressed, represents these nerves.

Other hydrodynamic principles applicable to the circulation can be illustrated by means of the model If, for example, the bulb, H, representing the left ventricle is compressed and decompressed more vigorously, more fluud will be pumped at each stroke into the system; the heights of the fluid in the side tubes, T, will show more pronounced fluctuations, and the mean pressure level will rise. More rapid strokes of the pump, with or without an increase of fluid forced into the system at each stroke, will also cause a rise in the mean level of the fluid in the side tubes. Changes of a similar nature in the action of the heart cause corresponding effects upon the arterial blood pressure.

If one of the small rubber tubes on the left of the bar, D, is loosened, punctured, or disconnected, so that a leak occurs in the system representing the arterial side of the circulation, a progressive



Fig 4 13 (Description in text.)

or a sharp fall in the levels of the fluid in the side tubes will occur; if the leak is considerable, the pressure cannot be restored to its original level even though rapid and vigorous strokes are made with the pump. In a similar way, loss of blood by hemorrhage causes a progressive or a sudden fall in blood pressure, the rate of the fall depending upon the severity of the bleeding. A leak in the right-hand

(venous) system would likely have a slower and a much less pronounced effect.

Let us now turn from the consideration of fluid pressure to a study of velocity of flow in a system of tubes. Physical principles governing the velocity of a fluid in a system of tubes comparable in size to the blood vessels are best illustrated by means of the model shown in Figure 4.13. The fluid in the reservoir, R, is kept at a steady level by means of the flow from a faucet. Whatever quantity of fluid enters the reservoir by the faucet must leave by the mouth, M, of the horizontal tube, H; otherwise the reservoir would overflow or be gradually emptied. It follows then that equal quantities of fluid must pass along equal lengths of the

tube, H, in any stated time. But section A is narrower (i.e., its capacity is less) than the sections of tube on either side. In order then to transmit the same volume of fluid, the velocity of flow in the narrow section must



Fig 4 14 (Description in text.)

increase. Section B is much broader; the flow in this section is, therefore, slower. It is immaterial to the general principle whether section B is a single channel or is divided into a number of smaller ones as in Figure 4.14, for, provided that the total cross area—that is, the sum of the cross areas of the small tubes in B—is greater than the cross area elsewhere along the horizontal tube, the flow through section B will be slower. Though each individual tube may be extremely narrow, the velocity of flow in each of the constituent tubes of B will be less than in any other part of the horizontal tube, H. Thus, with respect to velocity of flow, one of these fine tubes, which constitutes but a small part of the total cross area of the section which they occupy, differs from the single narrow tube, A, since the latter offers the sole channel at this point of tube H giving passage to the fluid In the circulation of the blood, the differences in velocity between, say, the aorta or large veins on the one hand and the capillary region on the other are due entirely to variations in the total cross areas of the vascular channels in the two situations (see also pp. 134-5).

Though, in general, the physical principles of the circulation can be demonstrated in a model, such a model differs from the vascular system in certain important particulars. In the first place, the tubes in the model, except those of the small rubber tubes in Figure 4.12, have rigid walls; they lack the elasticity of the living vessels, and this

property of the vascular structures is essential for maintaining an adequate supply of blood to the tissues in the intervals between the beats of the heart (p. 148). Again, certain physiological factors come into play in the living body under varying conditions, so that it is not always possible to predict what will happen under a given set of circumstances The complexity of these physiological factors, reactions and counterreactions, adjustments and compensations, preclude their being duplicated in any inanimate system.

The pressure of blood throughout the vascular system. In the model (Fig. 4.12), it will be observed that the oblique line R-R in the system on the left indicating the pressure levels at intervals along the




Fig. 4.15. Curve of blood pressure throughout the circulation—from left ventricle to right auricle

horizontal tube, if continued to the right, would not meet the corresponding line O-O in the right-hand system, but would fall above it This means that a fall in pressure more pronounced than elsewhere occurs in the region of the small rubber tubes representing the minute vessels of the vascular system This relatively sharp drop in pressure is due to a greater amount of energy being expended in over-coming the frictional resistance offered by the small-calibered tubes, C; for the total surface over which the fluid flows is greatly increased by these subdivisions of the horizontal tube. Thus, in the vascular system, a large part of the ventricular contraction is expended in overcoming the resistance offered by the small-calibered arterioles; an abrupt fall in pressure, therefore, occurs in this region of the circulation. A more gradual fall in pressure occurs from this point on through the capillaries and veins to the right auricle, which is filled at a very low pressure-namely, about 5 centimeters of water (see Fig. 4.15). It will be seen (Fig. 4.18, p. 134) that when the velocity of the blood flow throughout the vascular system is plotted, the curve obtained is quite different.

The work of the heart-the cardiac output. The energy whereby the pressure is maintained in the vascular system and the blood driven through the systemic circulation is derived from the contraction of the left ventricle. The right ventricle drives the blood through the lungs. The work performed by these two cardiac chambers will now be considered. The work performed by a machine is expressed in foot-pounds or in kilogram-meters; that is, the vertical distance, in feet or meters, through which a given weight, in pounds or kilograms, is lifted. Thus, if a weight of 2 kilograms is raised to a height of 4 meters the work done is 8 kilogram-meters. The efficiency of the machine is the proportion of the total energy expended which appears as work. Thus

work total energy expended = efficiency

The work performed by the heart when it contracts may be expressed in the same way, for it ejects a certain weight of blood at a mean pressure of about 1.25 meters of water (or blood), which is equivalent to saying that the heart raises the blood to a height of 1.3 meters. As we shall soon see, if a long glass tube be inserted into a large artery, and held vertically, the blood actually rises to a height of several feet.

Under ordinary conditions of bodily rest, by far the greater proportion (98 to 99 percent) of the work performed by the heart may be calculated from the following formula:

#### W = 7 OR

W = work in kilogram-meters Q = quantity of blood ejected, in kilograms R = mean blood pressure in the aorta, in meters of blood

To give an example: if the left ventricle discharges 60 grams of blood at each beat (contraction) against an aortic pressure of 125 mm Hg, that is equivalent to saying that the ventricular contraction raises 0.06 kilograms of blood to a height of 1.62 meters (mercury is approximately thirteen times heavier than blood). Then the work performed by the left ventricle at each beat is (1.62 × 0.06) 0.97 kilogram-meters, or (0.97 × 70) 68 kilogram-meters per minute, at a heart rate of 70 beats per minute.

The pressure in the pulmonary artery is only & of that in the aorta. That is, the pressure developed by the right ventricle is only a that developed by the left. The quantity of blood ejected by each ventricle is, however, the same The fraction  $\frac{7}{6}$ , therefore, is used to obtain the work of the whole heart—that is, of both ventricles

The cardiac output. The amount of blood ejected by one ventricle per minute is called the minute volume of the heart or the cardiac output. The minute volume of the human heart under ordinary conditions of mental and physical rest amounts to from 4 to 5.5 liters, depending on the size of the individual all must be remembered, how ever, that, per minute, an equal volume of blood is ejected by the right ventricle, and that an equal volume passes through the lungs and through the systemic circulation. The minute volume of the heart is therefore also referred to as the circulation rate. In other words it is the quantity of blood that passes per minute through the circulatory system. The quantity of blood ejected by one ventricle at each beat of the heart is called the stroke volume, it amounts to 60 or 70 cc under resting conditions. The cardiac output or minute volume is the product therefore of the stroke volume and the heart rate. Thus,

cardiac output (per minute) = stroke volume X heart rate.

The cardiac output is greatly increased during muscular exercise. In a robust healthy young man doing strenuous work it may amount to over 20 liters. The minute volume of the heart is raised by increasing the output per beat (stroke volume) and usually by increasing the number of beats per minute (heart rate) as well. In persons of poor muscular development, the heart rate accelerates in response to exercise to a greater extent than in those of athletic build. That is, the former in order to increase their cardiac output, depend upon increase in heart rate to a greater extent than do the latter. In some athletes for example, the minute volume may be increased several fold with little or no change in heart rate.

The minute volume of the heart is also increased during digestion at high environmental temperatures during emotional excitement and in the later months of pregnancy Among pathological conditions which increase the cardiac output are hiperthyroidum (p 411), fever

 The card ac output bears a closer relationship to the surface area of the body than to the body weight. The value is around 3 liters per square meter of body sur

face.

<sup>&</sup>lt;sup>4</sup> The cardiac output is affected by so many physiological conditions that in order for the values obtained in different persons to be strictly comparable and have a definite meaning with respect to cardiac function, the determination must be carried out under uniform or hoard cond to so. The subject must have been without food for 12 hours and be at mental and physical rest in a room kept at an ordinary comfortable temperature—about 25° C.

and severe anemia. It is reduced in hypothyroidism and in certain forms of cardiac disease.

The cardiac output in man can be measured only indirectly. The method of measurement is based upon what is generally known as the Fick principle. Fick pointed out that, if the quantity of carbon dioxide which was given off in the lungs (or of oxygen absorbed from the lungs) by each 100 cc of blood were known, and if the total quantity of CO<sub>2</sub> given off (or of oxygen absorbed) over a given period of time were also known, then the volume of blood which had passed through the lungs in that time could be calculated. For example, if the venous blood coming to the lungs contained 58 cc of carbon dioxide per 100 cc and the arterial blood leaving the lungs contained 52 volumes per 100 cc, then 6 cc of carbon dioxide per 100 cc must have been given off by the blood in its passage through the lungs. We must then first determine the gas content of the blood (either CO<sub>2</sub> or oxygen) in the arterial blood and in the venous blood. It is a comparatively simple matter to obtain a sample of arterial blood for analysis (by puncturing an artery with a hollow needle attached to a syringe). It is much more difficult to determine the gas content of the venous blood in the vessels of the lungs, or of the blood coming to the lungs (e.g., in the pulmonary artery). It can be obtained indirectly by a rather complicated procedure from the alveolar air, or by passing a slender rubber tube up the large vein of the arm into the right auricle of the heart and withdrawing a sample of blood. The oxygen or carbon dioxide content (that is, the load of either gas) of the blood in the auricle will, of course, be the same as that in the pulmonary arterv.

The total quantity of carbon dioxide eliminated (or of oxygen consumed) over a period of a few minutes is next determined (pp. 356-7). Let us say that this amounts to 240 cc per minute. We now have all the data necessary for the calculation of the cardiac output. Thus:

 $\frac{\text{(Total CO}_2 \text{ eliminated per minute)}}{240 \text{ cc}} \times 100 = 4000 \text{ cc cardiac output}} \times 100 = 6 \text{ cc} \times 100 \text{ cm minute volume}$ (Difference between volumes of CO, in venous and arterial bloods)

### The Velocity of the Blood

The velocity of the blood varies in different parts of the vascular system. It does not, like the blood pressure, show a continuously pro-

134

gressive diminution from the aorta to the right auricle. The blood moves swiftly through the large arteries. In the human aorta, for ex ample, its speed is around 0.4 meters per second when the body is at rest. The velocity decreases moderately throughout the arterial tree, but is greatly reduced in the capillaries, where it averages only 0.5 mm per second. The blood upon reaching the venous system speeds.




Fig 4.16 Diagram showing the pressure and velocity of the blood in different parts of the vascular system (Modified from Frederics) Note the relation between blood velocity and vascular area and the absence of rhythmical variations in pressure and velocity in the cap libries and veins

up again and, in the great veins feeding the heart (superior and inferior venae cavae), flows at a velocity not greatly less than that of the blood in the aorta

The hydrodynamic principle stated on page 129 and illustrated by the models in Figures 4 13 and 4 14 is of such importance that it will be well to apply it more directly to the circulation. It should be emphasized that except for very brief periods the quantity of blood returned to the right side of the heart in any given time must equal that ejected by the left ventricle. Variations in the velocity of the blood stream are due simply to differences in the dimensions of the

total cross area of the vessels through which the blood flows. The broader the passage for the blood, the slower will be its flow, and vice versa. It is quite immaterial to the principle whether the expansion of the bed of the stream is due to a single large channel, or to a division into a number of small, even minute, separate channels. Provided that the total cross area of a region of the vascular bed, which




Fig 4.17 Diagram illustrating the relative sectional areas of the aorta, capillary bed, and great veins. The area of the "capillary lake" is actually much greater relatively than is shown. (After Best and Taylor, The Physiological Basis of Medical Practice.)

constitutes the only passage for the blood from the left to the right side of the heart, is greater, the velocity of flow in the expanded part of the system will be reduced.

The human aorta is a little over an inch in diameter. All the blood ejected by the left ventricle must pass through this channel; the blood velocity must therefore be much greater here than in more peripheral parts of the vascular tree where, as a result of the numerous branchings and rebranchings, the total cross area of the vascular bed is increased. The flow in a carotid artery, for example, is slower than that in the aorta, for there are two carotids, as well as several other branches given off by the aorta, and the sum of their cross areas is considerably greater than that of the latter vessel. In the region of the capillaries where, as just mentioned, the flow is greatly slowed, the most pronounced widening of the bed is found. For this reason, the capillary area is frequently referred to as the "capillary lake." Though the cross area of each capillary is only a very small fraction of that of the aorta, the sum of all the cross areas of the capillaries is from 600 to 800 times greater (see Figs. 4 16 and 4.17).

5 The average diameter of a capillary is about 10 microns, the diameter of the human aorta is about 25 mm (approximately 1 inch). 136

In the venous system, the successive junctions of smaller veins to form larger trunks result in a progressive narrowing of the cross area of the vascular channels until, at the entrance to the right auricle, the two blood columns (in the superior and inferior venae cavae) have together a cross area only about double that of the aorta; and that is why the velocity of flow in these large veins is not so greatly less than the flow in the aorta. When the velocities throughout the vascular




Fig. 4.18 Curve of velocity throughout the circulation—from left ventricle to right auricle. Compare with Figure 4.15.

system are plotted, it will be found that the second half of the curve is radically different from the curve of pressure. See Figure 4.18 and compare with Figure 4.15, page 130.

The velocity of the blood in the arteries, but not in the veins, varies with the heart beat above and below a mean value. During the ejection of blood from the ventricle, the velocity of the arterial blood rises to a maximum, but falls again when the heart pauses. The minimum value is reached just before the next heart beat.

#### The Arterial Blood Pressure

That the blood in the arteries is under a relatively high pressure was not demonstrated until about a century after Harvey's discovery of the circulation. In 1733, an English clergyman, the Rev. Stephen Hales, inserted a small glass tube (called a cannula) into the artery of a horse's thigh and connected it by means of the windpipe (trachea) of a goose with a glass tube some 8 feet long. The long glass tube was placed vertically. The flexible trachea served the purpose of a rubber tube which would be used today, but had not then been invented. When the clamp, previously placed upon the artery, was released, the blood rushed into the tubing and rose nearly to its upper

end. The height of the blood column was observed to fluctuate several centimeters with each beat of the heart.

This method of measuring the blood pressure in animals has obvious disadvantages. In the first place, owing to the length of the tubing required, it is inconvenient. It is also inaccurate, since the blood which fills the long tube is out of circulation; the blood volume is correspondingly reduced, and the pressure as recorded is, in consequence, less than that which actually existed Finally, the experiment is soon interrupted by coagulation of the blood in the tubing. It was not until nearly another century had passed that an advance on Hales' method was made. In 1828, the French physicist Poiseuille used a glass U tube filled with mercury for registering the pressure and an anticoagulant fluid to fill the tubing which connected the U tube (now usually referred to as a manometer) with the cannula in the artery. Since mercury is over thirteen times heavier than blood, the mercury rises only a few inches instead of several feet, as when the pressure was measured by the height of a column of blood itself. Before the clamp is released from the artery, the anticoagulant fluid is introduced under a pressure equivalent to what the blood pressure is expected to be. If the pressure of the fluid is adjusted so that it exactly equals the blood pressure, blood does not leave the artery, nor does an appreciable amount of anticoagulant enter the circulation. It has been customary ever since Poiseuille introduced the mer-curial manometer to express the blood pressure in millimeters of mercurial manometer to express ne oncou pressure in mannecess of mer-cury (mm Hg). Some years later Ludwig, a German physiologist, improved the method by placing a float, fastened to a wire bearing a writing point, upon the mercury column. The height of the column of mercury and its fluctuations could in this way be recorded upon a writing surface—a revolving drum covered with smoked paper—and permanent tracings of the blood pressure secured. The instrument bearing the writing surface is run by clockwork, and is known as a kymograph. Figures 4.19 and 4.20 illustrate the manner in which the measurement and calculation of the blood pressure are made.

The phases of the arterial blood pressure. The maximum blood pressure of warm-blooded animals has a value of from 110 to 150 mm of mercury. Unlike certain other physiological functions (e.g., heart rate, output of the heart, basal metabolism, etc.), no relationship exists between the height of the blood pressure and an animal's size. The blood pressure of the rat, for example, is about the same as that of man. The arterial blood pressure, like the velocity of the arterial blood, varies rhythmically with the beating of the heart, rising to a




Fig 419 Showing method of recording the blood pressure in animals

maximum during the ventricular contraction (systole), when blood is pumped into the arteries, but falling again when the ventricle relaxes (diastole). The pressure reaches its lowest level just before the next beat. These variations are shown in Figure 4.21, which is a blood pressure tracing obtained from the carotid artery of an animal by the method described above.

The maximum or systolic pressure in a young man has an average value under ordinary resting conditions of around 120 mm Hg It may be a little above this value or 10 mm Hg or so below, and still be considered within the normal range. The minimum or diastolic pressure under similar conditions is around 80 mm Hg. The pulsation in the arteries is due to the difference between these two pressures, and is therefore called the pulse pressure, it amounts to




Fig. 4.20. U tube of mercury enlarged to show more clearly the manipulation of the manometer and the calculation of the blood pressure. The spring clamps 2 and 3 shown in Figure 4.19 are closed and clamp 1 has not been applied The mercury in both limbs of the U tube are therefore open to the atmosphere and at the same level indicating zero pressure (left hand sketch Fig. 420) A base line at this level is now inscribed on the rotating drum. Clamp 1 is ap plied (right hand sketch Fig. 4.19) clamp 2 is opened and the tubing up to the mercury column filed with an anticoagulant fluid from the bottle set at a suitable height (to give a pressure approximately equal to that of the blood) Clamp 1 is released momentarily to permit the escape of air. The mercury owing to the pressure of the anticoagulant fluid is depressed in the right hand limb and rises in the limb containing the float with the writing point When clamp 3 on the artery is now released the writing point may rise or fall a little depending upon how closely the animal's blood pressure has been predicted Fluctuations due to the heart beat now appear in the tracing. The height of the blood pressure is indicated by the distance in millimeters between A and Bthe levels of mercury in the two limbs of the U tube-or more conveniently by measuring with a millimeter rule and multiplying by two the d stance from the base line (zero pressure) on the drum and the blood pressure tracing

(120 - 80 -) about 49 mm Hg The value obtained by dividing the sum of the values for the systolic and diastolic pressures by two is called the mean pressure Under resting conditions, therefore, the mean pressure amounts to

$$\left(\frac{120 + 80}{2}\right) = 100 \text{ mm Hg}$$




Fig 421 Tracing of rabbits blood pressure Left hand sketch the larger fluctuations in blood pressure are due to the respirations the smaller ones to the heart beats The tracing near the bottom of the figure is made by a time marker. On the right a see ton of the tracing is ediagred S = systolic pressure. D = diagnostic actolic.

The fluctuations in the blood pressure due to the contractions and relaxations of the ventricle occur so rapidly that a mercurial manom eter, owing to the inertia of the mercury, fails to follow them closely, always showing a lag. The thrust given to the mercury by the systolic rise in blood pressure is of such short duration that the heavy liquid does not reach the height which it would do were the pressure rise developed more gradually or sustained for a longer time. The systolic pressure as recorded by a mercury manometer is, in consequence, lower than the true value. In Figure 4.21 for example, the fluctua tions due to the phases of the heart's action would be more pronounced if the recording apparatus were lighter in weight and, there




Fig 4.22 Illustrating a nonmercurial type of apparatus for recording blood pressure which obviates the inertia due to mercury

fore, showed less inertia. In animal experiments in which it is desired to show the true blood-pressure values or to reveal minor fluctuations on the main waves, apparatus of lighter weight is employed, the mercury being replaced by a lighter liquid; or tubing and a tambour as shown in Figure 4 22 is used. The method of recording the blood pressure in man (see below), in which the larger artery of the arm is compressed by air pressure which is measured by the height of a column of mercury, avoids the error due to the inertia of the mercury.

The clinical measurement of the arterial blood pressure. The blood pressure is measured indurectly in the human subject by means of an instrument called a sphygmomanometer (see Fig. 4.23). This comprises a flat rubber bag about 5 inches wide and 8 inches long, enclosed in an envelope of cotton cloth to prevent it from stretching under pressure. Two tubes, one connected to a manometer (mercurial or aneroid) and the other to a small hand pump provided with valves, communicate with the interior of the bag. The cloth-covered






Fig 4.23 Sketch illustrating the auscultatory method of determining the blood pressure in man. Below, drawing to illustrate the principle; cross section of the arm, with the armlet wrapped around it and connected with the manometer.

terior of the bag. The cloth-covered bag, which is usually referred to as the cuff or armlet, is wrapped snugly around the arm above the elbow, the tails of the cloth being wound bandage fashion to hold it in position. The pressure within the bag is raised by a few compressions of the hand bulb until the pulse at the wrist disappears—that is, until the air pressure overcomes the blood pressure and occludes the main artery (brachial) of the arm. The observer listens with a stethoscope placed over the artery in front of the elbow just below the armlet, while he gradually reduces the pressure by opening the valve close to the

bulb As the air pressure falls and allows the blood to escape beneath the cuff and fill the artery below, faint tapping sounds synchronous with the heart beats are heard. The reading of the manometer at which the sounds are first heard is taken as the systolic pressure. The sounds become progressively louder as the air pressure is reduced further, then undergo a series of changes in quality, and finally disappear. At a pressure level about 5 mm above that at which they disappear, the sounds acquire a soft muffled quality, the manometer reading at which this occurs indicates the diastolic pressure.

The method of measuring the blood pressure just outlined is called the auscultatory method (L. auscultare, to listen) It is the method most commonly practiced

Blood pressure variations in health. The arterial blood pressure is influenced by a number of physiological conditions-age, sex. muscu lar exercise, emotion, digestion, and posture. In the newborn baby, the systolic pressure averages 40 mm Hg, but reaches a value of around 80 mm Hg by the end of the first month It rises more gradually throughout childhood attaining a level of about 100 mm by the twelfth year and 120 or so by the seventeenth It remains around the latter level for a varying number of years. There is a tendency for the blood pressure to rise gradually after the twenty fifth year, the increase in systolic pressure averaging about 0.5 mm Hg per year. For example a systolic blood pressure of 135 mm Hg or so, though normal for a man between 50 and 60 years of age, would be too high for one between 20 and 30 But no fixed rule can be stated with respect to the increase in blood pressure with advancing years. It is questionable whether the tendency can be regarded as a physiological process, as is generally supposed, it is by no means of general occurrence. As compared with men of the same age, women usually have a somewhat lower blood pressure

Muscular exercise causes a pronounced temporary rise in the blood pressure, the systolic pressure rising during strenuous exertion to 180 mm Hg or more, the diastolic pressure shows a smaller proportionate rise. The pulse pressure is therefore, increased by exercise A change from a lying down to a standing position causes a rise in the diastolic, but little change as a rule in the systolic pressure, the pulse pressure is therefore reduced

Emotion (e.g., fear worry, excitement, etc.) may cause a rise of several millimeters of mercury in the systolic pressure, the diastolic pressure is raised to a smaller extent During digestion a moderate rise in the systolic pressure with little change in the diastolic occurs

A reduction of from 20 to 30 mm Hg in the systolic pressure occurs during restful sleep, but if the sleep is disturbed by exciting or terrifying dreams, the systolic pressure may be elevated to 180 or 200 mm He.

Arterial hypertension. A persistent elevation of the arterial blood pressure above the normal range is called arterial hypertension. The abnormally high pressure is due to spasm of the arterioles (i.e., to an increase in the peripheral resistance, see p. 146). The systolic pressure, in the most severe cases of the disease, may amount to 300 mm Hg, and the diastolic pressure to 150 mm Hg. The condition leads to serious consequences; it throws extra work upon the heart and subjects the arterial system to undue strain. Enlargement of the heart, followed eventually by cardiac failure, and deterioration of the arterial tissue (arteriosclerosis, p. 144) result. Rupture of a diseased cerebral vessel may cause death; or death may result from the kidneys failing to perform their functions as a result of pathological changes in the renal vessels. There are two types of arterial hypertension. In one type, called primary or essential hypertension, the condition occurs unpreceded by kidney or any other disease. Essential hypertension, therefore, may cause, but is not itself the result of, kidney disease. In the other form, the hypertension is a consequence of chronic kidney disease (chronic Bright's disease, chronic nephritis).

A few years ago, Dr. Goldblatt and his associates in Cleveland carried out an interesting and highly important series of experiments upon dogs, which has shed a light on the origin of the type of arterial hypertension developing in the course of chronic renal disease. By means of a specially devised clamp, the renal artery to one kidney or to both was constricted and the renal blood flow thus reduced. When the renal artery of one side alone was narrowed in this manner, the blood pressure rose moderately but returned to normal again within a few weeks. When, however, the blood flow to both kidneys was reduced, or to one kidney only and the other kidney then removed, the blood pressure rose to a higher level and remained permanently elevated.

It has been shown that a reduction in the renal blood flow causes the production by the kidney of a chemical substance which, upon its discharge into the circulation, results in the constriction of the arterioles throughout the body. Such vascular constriction (see peripheral resistance, p. 146) is the direct and immediate cause of the hypertension in the experimental animals. The substance formed by the kidney is known as renin; it has been demonstrated in the blood of animals

144

suffering from experimental hypertension, as well as in the blood of patients with the disease

The constriction of the peripheral vessels in hypertension, result ing from clamping of the renal artery, is not caused directly by renin but by the production of another substance called angiotonin or




Fig 4 24 Showing mechanism in experimental hypertension. The long arrow on right above represents the action of angiotomin.

hypertensin (Fig 4.24) Renin is an enzyme which acts upon a globu lin of the plasma to produce the latter material—which is the active vasoconstrictor agent

It is strongly suspected that the hypertension occurring in renal disease is also due to the renin angiotonin mechanism as a result of some interference with the renal blood flow

The possibility that essential hypertension is also due to some ab normality in the renal circulation has been investigated by a number of research workers, but definite evidence for such a connection has not been secured

Arterial disease Almost from birth the fabric of the arteries starts to deteriorate, imperceptibly in the earlier years, but often quite rapidly from middle age on Reduction of the elasticity of the arterial walls is the first physical change to be detected. The processes of degeneration vary, however, very greatly in different persons "A man is as old as his arteries" a saying attributed to Osler, and which means that generally speaking the usual signs of old age and the de terioration of the arteries go hand in hand

There are two principal types of arterial disease. In one form, the vessels become hard stiff and brittle (arteriosclerosis), owing to atrophy of the muscular and elastic tissues and their replacement by

salts of lime. This change in the vascular wall permits the superficial vessels to be felt beneath the skin and to be rolled under an examiner's finger. Such "pipe-stem" arteries as they are called are met with most frequently in old age. In another much more serious type of arterial disease, the endothelium (tunica intima) is primarily attacked. Patches of the endothelial lining break down and are replaced by a soft fatty material in which lime salts are later deposited. The patches tend to spread and strip up the inner coat, and even to invade the nuscular coat. This type of arterial degeneration is called atheroma, atheromatosis, or atherosclerosis (Gk. atherē, porridge + scleros, a hardening). It leads to weakening of the arterial wall, and is the cause of such vascular emergencies as cerebral hemorrhage (apoplexy) and coronary thrombosis. In the latter, the roughened inner surface by replacing the smooth endothelial lining has encouraged clotting within the vessel. This disease appears to be due to a disturbance in lipid metabolism. The atheromatous patches are composed of fats, cholesterol, phospholipids, and protein-lipid compounds (lipoproteins). Persons recovering from coronary thrombosis have an abnormally high concentration of plasma lipoproteins with a low density, and these, it is believed, are related to the atheromatous changes in the arterial walls These lipids are also in high concentration in the blood of persons showing other clear evidence of atheromatous disease, and their concentration in the plasma tends to rise with age.

The cholesterol and other lipids of the blood are derived largely from the diet, and it would appear that in atheromatosis certain of these substances when in excess in the blood are deposited in the arterial wall. It is advised, therefore, that obese persons and those with a high concentration of lipids in the blood, or who have a family history of vascular disease should restrict their consumption of fatty materials—eggs, cream, butter and other animal fats.

The factors responsible for the maintenance of the normal arterial blood pressure. The arterial blood pressure is the result of the discharge by the heart into the arterial system of a volume of blood which cannot all escape through the peripheral arterial vessels (arterioles) into the venous system (through the capillaries and veins) before the next heart beat occurs. This means that the arterial system during life is at all times overfilled. The elastic arterial walls are therefore always more or less stretched. The height of the blood pressure is thus dependent upon two main factors—the output of the heart, and the caliber of the peripheral vessels. These and other factors concerned are listed below and will be considered in some detail;

- 1. The cardiac output-that is, the volume of blood discharged by the heart per minute,
  - 2 The peripheral resistance,
  - 3. The total blood volume
  - 4 The viscosity of the blood,
  - 5 The elasticity of the arterial walls

The cardiac output. It has been mentioned that the cardiac output is, as a rule, augmented by increasing both heart rate and the stroke volume, but that the extent to which each of these factors participates varies. The larger amount of blood discharged per minute into the arterial system can be accommodated within the arterial system only



Fig 4 25 Showing the effects upon systolic, diastolic, and pulse pressures of an increase in stroke volume, heart rate, and of peripheral resistance (After Wiggers )

by further stretching of the arterial walls. Both systolic and diastolic pressures rise, but both pressures are not necessarily raised to the same degree When the increase in cardiac output is the result mainly of an increase in the stroke volume, the systolic pressure rises to a greater extent than the diastolic, with a consequent increase in pulse pressure (Fig 425) If, on the other hand, the heart rate is markedly increased, the diastolic pressure tends to rise to a greater extent than the systolic because, since the time between beats is shortened, the decline in pressure during the diastole of the heart is arrested at a higher level by the arrival of the next beat. The pulse pressure is there fore reduced

The peripheral resistance This refers to the small vessels at the periphery-the arterioles and capillaries-but especially to the former, which, as already mentioned, offer resistance to the passage of blood from the arterial system into the veins. The arterioles and capil laries are furnished with nerves through which their calibers may be altered from time to time

Constriction of the peripheral vessels reduces, temporarily, the outflow from the arterial system and, in accordance with the principle stated on page 127, a rise in arterial pressure occurs, provided the minute volume of the heart is not at the same time reduced. The arterial system must, therefore, contain more blood than formerly, and the arterial walls are more forcibly stretched. The pressure rises to the point at which the quantity of blood leaving the arterial system again equals that which is pumped into it by the heart, after that, no further rise in pressure occurs. Opposite effects are produced when the peripheral vessels dilate. The pressure falls until the outflow from the arterial system again just balances the inflow. The peripheral resistance may be compared to a dam in a river. If the dam is raised or lowered, the water continues to rise or fall, respectively (that is, its pressure increases or decreases), until it reaches the new level. From then on, the quantity of water which overflows in a given time is the same as it was at the original level.

Variations in the caliber of the arterioles of the abdominal (splanchnic) region are more effective than those of other vascular areas in causing changes in the blood pressure. The splanchnic vessels, when fully dilated, have a relatively immense capacity—sufficient to accommodate almost the entire blood volume. Some sudden and strong emotion may cause their dilatation, when a fall in arterial pressure and loss of consciousness (fainting or syncope) may result. On the other hand, stimulating the great splanchnic nerve (the large nerve of the abdomen through which nervous impulses reach the vessels, p. 195) in an anesthetized animal, as by the application of an electric shock, causes constriction of the abdominal vessels and, in consequence, a pronounced rise in blood pressure.

The blood volume. A sufficient volume of blood must be available to overfill the arterial system. That is to say, the arterial walls even at the lowest normal pressure (diastolic) are stretched. Any considerable reduction in blood volume, as by hemorrhage, causes the blood pressure to fall. Restoration of the blood volume by transfusion of blood or of a suitable blood substitute (p. 77) is followed by a rise in pressure.

The viscosity of the blood. Viscosity may be defined simply as the "thickness" of a liquid or the resistance which a liquid offers to its movement through a narrow tube. The viscosity of a particular liquid is determined by measuring the time which it takes, when acted upon by a constant force, to pass through a given length of a capillary tube. The viscosity of water is taken as unity, and that of various other

liquids is expressed relatively to that of water. Thus, alcohol and ether are less viscous than water, while glycerine, oil, and syrup are much more viscous Blood itself is some five times more viscous than water We know that should two piston syringes be filled, one with a highly viscous liquid, such as syrup, the other with water, a greater effort would be required to drive out the syrup than to drive out the water, and a higher pressure would be created in the first syringe than in the second The relatively high viscosity of blood increases the resist ance in the vascular system, it tends to impede the flow through the arterioles, and so to enhance the effect which these narrow vessels themselves have in maintaining the arterial pressure Variations in viscosity, therefore, have an effect upon the blood pressure similar to that caused by alterations in the caliber of the peripheral vessels The relatively high viscosity of blood is due to the plasma proteins, and, to an even greater extent, to the corpuscles The viscosity is lowered in conditions associated with low plasma protein concentration and in some anemias. It is increased in conditions associated with a high corpuscular concentration or when as a result of the loss of blood water (anhydremia), the protein concentration of the blood is raised

The elasticity of the arterial walls. If the arteries possessed rigid and not elastic walls the systolic pressure would be higher, other things being equal but there would be no diastolic pressure. That is to say, the arterial blood pressure would fall to zero between the beats of the heart. The absence of a rhythmical variation in the pressure—that is of a pulse—in the capillaries and veins is due, in turn, to the existence of a pressure in the arteries during diastole. These facts are best illustrated by means of a model.

In Figure 4 26 is represented a bulb syringe, S, valved at A, and having a short tube, B which dips into a basin of water Leading from the opposite pole of the bulb is a longer tube, C. If the bulb is alternately compressed and released, water will be drawn from the basin, and discharged from the mouth of the tube (Fig. 4.26, 1). If the walls of the latter are composed of some rigid material (e.g., glass), it will be found that when the pump is worked the fluid issues from the tube in spurts or jets synchronous with each stroke, but no flow occurs between the strokes. An increase in the frequency or force of the strokes does not after the intermittent character of the flow, nor does lengthening the tubing. If the peripheral resistance of the vascular system is imitated by attaching a nipple of small bore to the mouth of the delivery tube, thus increasing the resistance to the outflow of fluid, the issuing stream is finer and its velocity is increased, but it

still remains intermittent (Fig. 4.26, 2). Now let the elasticity of the arterial wall be imitated by replacing the rigid tube, C, by one of rubber, but let the mouth of the tube be left free and not constricted in any way (Fig. 4.26, 3). The intermittent character of the flow from the tubing is unaffected. However, if the small-bored nipple representing the peripheral resistance be fixed into the mouth of the elastic tubing, the issuing stream will be found to have lost its pulsatile character and to have become continuous (Fig. 4.26, 4). Two factors are therefore necessary to produce a continuous outflow: (a) elastic tubing




Fig 4 26 Diagram illustrating the dependence of diastolic pressure upon the arterial elasticity and peripheral resistance. (Description in text.)

and (b) resistance to the outflow. The reasons for this are clear. If the tubing is made of elastic material, its wall is distended by the force of the pump and energy is thus stored up which is expended between beats as the elastic wall recoils. (See Fig. 4.27.) Thus, the fluid is driven along the tube during the pauses of the pump. Nevertheless, if the fluid can escape freely from the tube, the pressure does not rise to a sufficient height to distend the rubber wall—that is, elasticity is not called into play, overfilling of the tube does not occur, and, in consequence, the latter acts simply as though it were composed of rigid material.

The foregoing facts apply directly to the arterial system. The elasticity of the vascular walls and the peripheral resistance are both essential for the development of the diastolic pressure. As the contents of the ventricle are forced into the already overfilled arterial system during systole, the added pressure which is then exerted upon the vascular walls causes their further distension. After the completion of systole, the elastic walls rebound (recoil) and, pressing upon the blood within their embrace, force it onward through the peripheral vessels. In other

words, the arterial lumen returns to its previous diameter and the energy that had been stored up during the stretching of the elastic tissue is in this way gradually expended during diastole (Fig. 4.27).

It may then be said that the elastic recoil of the arterial wall acts, in a sense, as a subsidiary pump to drive the blood onward in a con-




Fig 4.27 Diagram to illustrate the storing of energy of the heart's contraction in the elastic arterial walls (at root of aorta) during systole and its release during diastole. Small arrows indicate the direction of movement of the arterial walls and of the force exerted during the two phases. Large arrows indicate the direction of the blood flow, S. S. semilianar (aortic) value.

tinuous stream between the heart beats. Otherwise the pressure would fall to zero after each systole.

Interaction between the several factors maintaining the blood pressure. In health, the several factors mentioned in the foregoing paragraphs interact with one another to maintain the blood pressure at a fairly constant level. When a change occurs in one factor which, acting alone, would raise or lower the pressure, a compensating reaction of a reflex nature occurs which tends to offset its effects: For example,

a moderate reduction in the cardiac output or in blood volume (as through hemorrhage) is not necessarily followed by a fall in blood pressure, since reflex narrowing of the calibers of the small vessels occurs: that is, the peripheral resistance is increased. Again, dilatation of the arterioles in one region, which of itself would lower the blood pressure, is balanced by construction of the minute vessels in another (Lovén reflex). Also, increased blood volume or a rise in blood viscosity, which alone would cause an elevation of the blood pressure, is counteracted by vasodilatation. (See also pp. 196 and 197.)

#### Capillary and Venous Pressures

It has been mentioned that, except for the sharp drop which occurs in the region of the arterioles, the blood pressure falls in a gradual slope from the left to the right side of the heart (see Fig. 4.15, p 130). The pressure is about 30 mm Hg at the arterial end of the capillary and about 12 mm at the venous end. The pressure of blood in the great veins is low (although the velocity is relatively high) and at the entrance of these vessels into the right auricle is only some 5 mm H<sub>2</sub>O. In other words, the energy of the cardiac contraction has been almost entirely expended in overcoming the resistance offered to the flow of blood through the vascular system

The venous pressure may be measured in man by an indirect method based upon the same principle as that employed for the measurement of the arterial blood pressure, but water or saline is used in the manometer for, owing to the low pressure to be measured, mercury, which is some thirteen times heavier than these liquids, would show a scarcely measurable change in level A vein is selected upon the back of the hand and a small chamber with a transparent top is hermetically sealed by means of a special cement to the overlying skin (Fig. 4.28). The chamber is provided with two tubes, one of which is connected to a water manometer, the other to a hand bulb. The pressure within the chamber is raised by a few compressions of the bulb until the vein is seen to collapse. The pressure, which at this point just balances the pressure in the vein, is read from the manometer. While the measurement is being made, the vein under observation must be brought to the level of the right auricle; otherwise, the hydrostatic pressure of the venous blood—that is, the weight of the blood column extending from the vein level to the auricle-would be added to the pressure due to contraction of the left ventricle.




Fig. 428 Measurement of the venous blood pressure The small chamber with transparent top (A) is cemented to the skin over a superficial ven The pressure within the chamber is raised by means of the pressure-bulb arrangement B. C., D., until the ven collapses E, manometer (After Hooker and Eyster.)

The pressure of blood in the peripheral veins, such as those of the limbs, amounts to from 60 to 100 mm H.O, it falls progressively through the larger veins to the right auricle (See Fig. 4.15, p. 130.)

## The Effect of Gravity upon the Circulation

When the human subject assumes the erect posture, the blood pressure in the arteries of the head and arms is not reduced; as stated on page 142, the diastohe pressure actually rises This is so, even though the subject makes no effort, but is tilted on a moveable table from the horizontal to the upright position. If a fall in blood pressure of sufficient magnitude to deprive the brain of an adequate supply of blood occurred when the body changed from the recumbent to the standing position, fainting would result. Now, in order to maintain

the blood pressure, the heart pump must receive an adequate supply of blood. The question then arises, "How, when the body is standing, is the blood conveyed against gravity up the great veins to the heart?" The factors concerned are four in number:

- 1. The contraction of the left ventricle. This vis a tergo (force from behind), as it is termed, drives the blood through the arterioles and capillaries and along the veins to the right side of the heart.
- 2. The respiratory movements. The pressure within the thorax (see intrathoracic pressure, p. 232) is always a little below that of the atmosphere. This "negative" pressure, which is exerted upon the blood through the thin-walled auricle and venae cavae, fluctuates with the respiratory movements, being increased (i.e., becoming more nega-tive) during inspiration, when it amounts to about -6 mm Hg, and decreased during expiration to about -2.5 mm Hg. The intrathoracic pressure thus exerts a suction effect or pull (vis a fronte) upon the column of blood in the large veins of the abdomen.

The blood is also forced upward during inspiration, for when the diaphragm descends it increases the intra-abdominal pressure. The increased pressure is transmitted to the blood in the veins; since the blood is prevented by the presence of valves in the large veins of the blower limbs from passing downward, it is propelled upward. The respiratory movements thus act both as a force pump and as a suction pump to move the venous blood toward the heart.

The effect of the respiratory movements upon the flow of blood to the heart is especially important during muscular exercise when the

excursions of the chest are deep and rapid.

3. The abdominal and limb muscles. When one is standing, the abdominal muscles contract and offer a firm support for the large veins of the abdomen The venous walls are prevented from becoming over-distended under the weight of the blood which they contain. If the walls of the veins stretched and so increased their capacity, a larger mass of blood would be accommodated in the large veins of the abdo-

mass of blood would be accommodated in the large veins of the abdomen, and the flow to the heart correspondingly reduced.

The muscles of the limbs perform a similar duty for the limb veins. Furthermore, they exert, during muscular movements, a varying pressure or massaging action upon the veins. By virtue of the valves in the veins (Fig. 4.11, p. 125), any pressure upon the column of venous blood is converted into an upward propulsive movement.

4. Constriction of the vessels of the splanchnic area. Through the

action of the vascular nerves (p. 195), the small vessels—arterioles. capillaries, and small veins-in the abdomen undergo a reduction in

caliber, and so in capacity Thus, blood which otherwise would collect in these areas when the erect posture is assumed, is directed up ward into the large venous trunks

In quadrupeds the mechanisms for raising the blood against gravity are much less efficient than they are in man or in the ages. The dog however, is able to compensate fairly well for the gravity effect, but the domestic rabbit may become unconscious or even die, if held up by the ears for any considerable length of time. This result is due to the heart being inadequately supplied with blood which owing to the flabbiness of the abdominal muscles of the tame rabbit collects in the large veins. Sheep also not infrequently become unconscious if held in a vertical position during shearing Fainting (syncope) in the human subject is in most instances due, not to any affection of the heart itself but to reduction of the venous flow to its chambers for if the left ventricle does not receive an adequate supply of blood the blood pressure in the vessels of the brain is not sustained, and the con scious centers suffer from lack of oxygen Fatigue or some strong emotion such as fear or shock may, by inducing splanchnic vasodila tation and imperfect cardiac filling cause loss of consciousness But a factor probably of much greater importance than splanchnic vasodila tation is the widespread dilatation of the blood vessels in the muscles that occurs in most cases of fainting

A person who stands quietly for some time—a soldier standing at attention on parade for example—may fall in a faint because the venous flow is unaided by muscular movement. Such a result is the more likely to occur if he is fatigued or if the day is hot and the vessels of the skin are therefore dilated. The treatment of a subject in a faint is simple he is laid in the horizontal position the effect of gravity upon the circulation being thus abolished. He should not be supported in the sitting position. As a matter of fact, the subject of a faint usually does the right thing himself—he falls to the ground Pressure may be applied to the abdomen in order to force blood along the veins to the heart but such a procedure is usually unnecessary

## The Effects of Acceleration upon the Circulation

A constant speed however great has in itself no effect upon the circulation but acceleration—that is, a change in velocity either in the

line of motion (linear acceleration) or in a change in direction, as when a body traveling along a straight course turns into a curvilinear one (entripetal acceleration)—may cause profound circulatory effects Such effects not uncommonly result from the sudden and pronounced changes in direction entailed in military flying For example, a pilot pulling out of a power dive—1 e, changing direction at high velocity from a downward to a horizontal and upward movement—has his head directed inward toward the center of a circular course —has his head directed inward toward the center of a circular course and the lower part of his body directed outward. His blood, as a re sult of centrifugal force, is therefore "forced' away from the head into the lower part of the body. The high pressure of blood in the capillaries in the skin of the lower limbs may result in small cutaneous hemorrhages. The return of venous blood to the heart is reduced, with a consequent fall in the blood pressure in the cerebral and retinal vessels. Vision is temporarily lost and the pilot may become unconscious. Such effects are commonly referred to in the flier's parlance as blackout "The abdominal organs may also be forced downward and drawing upon the diaphragm embarrass respiration. Any maneuver in which the head is directed outward—that is, toward the circum ference of the circular movement-causes the reverse effect upon the circulation The blood is now "forced toward the head The vessels of the head and neck become engorged and severe throbbing pain in the head is experienced. The eyes feel as if they were being forced from their sockets. There may be mental confusion. Rupture of a cerebral or retinal vessel may result

The force exerted upon the body is expressed in units of gravita tional force and designated +G or -G, according to whether the acceleration is positive or negative, that is, according to whether the blood is driven away from or toward the head, respectively A force twice that of gravity is referred to as 2 Gs, one three times gravity, as 3 Gs, and so on Persons vary with respect to the centrifugal force which they can withstand before "blackout" occurs The limit for most pilots is between 5 and 6 Gs (+) During World War II, double-walled suits were fashioned which were worn by fighting airmen in order to prevent 'blackout" effects Air under pressure, or water be tween the two layers of the suit, by affording support to the tissues of the lower part of the body, prevented the distension of the vessels and the pooling of blood in this region Thus the vital centers of the brain were assured of an adequate supply of blood

#### The Arterial Pulse

The pulse in the arteries is due, of course, to the beat of the heart. Though all arteries, large and small, pulsate, the pulse is felt most conveniently by placing the finger upon the radial artery, which lies near the outer border of the palmar aspect of the wrist. In health, the rate of the pulse, when the body is at rest or engaged in some light occupation, is around 70 per minute. The pulse is a pressure change transmitted as a wave through the arterial wall and blood column to the periphery. Though caused by the discharge of blood into the arterial system, it is in the nature of a transmitted impact, and is not due to the passage of the ejected blood itself. For example, when one




Fig. 4.29 Showing successive stages in the transmission of a pulse in a ligated artery.

places a finger upon the artery at the wrist, the pulse beat which is felt is not due to the arrival of blood which has been discharged an instant before from the left ventricle into the arterial system, but is due rather to the shock or impact given to the resilient arterial walls by the sudden entrance of that blood into the aorta. The wave thus set up in the wall of the aorta and in the column of blood therein travels from ten to fifteen times more rapidly than does the blood itself, and quite independently of it. The blood, as we have seen, has a velocity of around half a meter per second in the aorta, and less than that in the smaller arteries. The speed of the pulse wave is from 5 to 8 meters per second, and is practically the same throughout the arterial tree. That the pulse travels quite independently of the blood flow is evident from the observation that in an artery which has been tied across so that no blood flows through it, the pulse is not abolished, but travels in the usual manner up to the point of occlusion (Fig. 4.29).

The velocity of transmission of the pulse wave is dependent upon


the character of the arterial wall. Were arteries made of unyielding material (e.g., glass), they and the incompressible blood within them would constitute a rigid system and, therefore, transmit an impact almost instantaneously from end to end. The arteries are, however, elastic and, in consequence, transmit a shock applied to their walls much more slowly.

The arteries become less resilient with age and the speed of the pulse wave therefore increases. At 5 years of age, for example, the velocity of the pulse wave is around 5 meters per second, whereas at the age of 80 it is in the neighborhood of 8 meters per second. In arteriosclerosis, in which condition the arteries are abnormally rigid, the pulse wave has a still higher velocity. The speed of the pulse wave can be measured by placing two levers, one upon an artery near the heart, the other near the periphery, and timing the arrival of the pulse at the two points. This distance between the two points is then measured. The distance (in millimeters) divided by the time (in seconds) which the pulse wave has taken to pass between the two points will give the velocity (in millimeters per second).

Thus,  $\frac{D}{T} = V$ , where D = distance in millimeters, T = time in seconds and V = velocity in millimeters per second.

The instrument employed for obtaining a graphic record (tracing) of the pulse wave is called a *sphygmograph* (see Figs. 4.30 and 4.31). The upstroke of the curve drawn by this instrument during the passage

Fig. 430 Upper drawing, Dudgeon's sphygmograph in position. Lower, diagram of mechanism. Ci., clockwork driving the smoked paper, Tr., under the writing point of the level, L. Sp. is a steel spring with a button, B, which is applied over the radial artery. With each expansion of the artery, the button is moved upwards and causes a movement I of the system of levers as indicated by the arrows.



of the pulse wave is abrupt and is called the anacrotic limb (Gk ana. up + crotos pulse) The downstroke has a more gradual inclination and is called the catacrotic limb (Gk cata down). The catacrotic limb shows a short distance from the peak of the primary wave, a sec ondary elevation known as the dicrotic wave, which is immediately preceded by a slight negative fluctuation called the dicrotic notch The latter is due to the slight backward movement (i.e., toward the heart) of the blood column in the artery, when the ventricle com mences to relax. The dicrotic wave is caused by the checking of this backward movement of the blood column and its rebound from the aortic valves which coming into accurate apposition at the end of



Fig 431 Tracing of radial pulse 1 anacrotic limb 2, dicrotic wave 3 dicrotic notch

nortic valves are removed, or hooked back with wires to pre vent their closing the dicrotic wave is not seen In normal cir cumstances dicrotic fluctuations can be revealed only by means of

ventricular systole (see p 170), prevent the leakage of blood into the relaxing ventricle When the

some graphic method of recording they cannot be felt by the finger placed upon the pulse In certain conditions however, in which the blood pressure is abnormally low and the arterial wall, in consequence, less tense, the dicrotic wave may be felt as a distinct tap following the main wave. A pulse of this character-the so called dicrotic pulse -is not uncommonly present in typhoid fever

It should be mentioned that the height of the pulse tracing gives no indication of the height of the systolic blood pressure. Nor does the pulse wave as traced by the sphygmograph give any indication of the length of the pulse wave itself. The pulse wave in the arterial system is actually a long swell measuring from 5 to 6 meters (16 to 20 feet), that is, it is several times longer than the arterial system itself

### The Structure and Physiological Properties of Cardiac Muscle

The fibers of heart muscle possess cross striations, but these are less well marked than those of skeletal muscle. Unlike the cells of skeletal muscle the cardiac fibers have no surcolemma separating or insulat ing one fiber from another. On the contrary, they give rise to branched processes-slips or bridges-which join neighboring fibers and thus



Fig. 4.32 Cardiac muscle fibers.

establish continuity between them. The heart muscle is therefore in reality a syncytium; that is, a large multinucleated mass of protoplasm rather than, like skeletal muscle, a number of separate units bound together (see Fig. 4.32).

The physiology of cardiac muscle will be taken up under the following heads: (1) excitability and contractility, (2) rhythmicity, (3) conductivity.

1. Excitability and contractility. Heart muscle responds by a contraction of its fibers to the various types of stimulus—thermal, chemical, mechanical, or electrical—which excite skeletal muscle. Its response follows the "all-or-none" law or principle. This law, which also applies to the conduction of the nerve impulse (p. 471) and to the contraction of an isolated fiber of

skeletal muscle (p. 488), states that a stimulus, if it produces a response at all, will produce the maximum response of which the tissue is capable under the conditions existing at the moment. That is to say, increasing the strength of a stimulus above that which is just necessary to excite the heart muscle (threshold strength) does not increase the force of its contraction.



Fig 4.33 Record of a series of contractions of frog's heart caused by electric stimulation of the ventricle. The ventricle was rendered quiescent by drawing a tight ligature (Stamuus' ligature) between the auricles and the ventricles. Tracings taken on a rapidly moving surface. First contraction (1) the lowest, last contraction (10) the highest. (After Hallibutron.)

160

Treppe When the heart muscle is stimulated at regular intervals, the first few contractions increase progressively in amplitude From the appearance given to the tracing by the initial contractions (Fig 4 33), this behavior is referred to as treppe or the staircase phenomenon. It is probably due to the beneficial effect exerted upon one contraction by the rise in temperature and the acid metabolites (carbon dioxide and lactic acid) produced during the contraction or contractions which have preceded. The phenomenon of treppe does not, as may at first appear, contradict the "all or none" law, for the conditions in the muscle—that is, a rise in the temperature and acidity—have been altered by the initial contractions.

Refractory periods Heart muscle will not respond to a stimulus while it is contracting, in this it differs from skeletal muscle (p 488). This time, during which the cardiac muscle is unresponsive to stimulation is called the absolute refractory period. The absolute refractory period of contraction. The absolute refractory period is followed by one in which the muscle gradually recovers its excitability. At the beginning of the latter—that is at the commencement of cardiac relaxation—a stronger stimulus is required for excitation, and the force of the contraction which results is less than the normal Excitability increases progressively and reaches its full value only when relaxation is complete. This phase of depressed but gradually increasing excitability is called the relatively refractory period.

2. Rhythmicity. This is an outstanding characteristic of the heart. The heart of the frog will contract spontaneously and rhythmically for a considerable length of time after its removal from the body, if simply kept moist with physiological saline (06 percent solution of sodium chloride). The rhythmicity is also an inherent property of the mammalian heart, it occurs quite independently of the central nervous system (see pertusion of the heart p 162).


When the heart of an animal is stimulated by an electric shock or a

When the heart of an animal is stimulated by an electric shock of a series of such shocks, the rhythm may be disturbed (made irregular or more rapid) but it will not, as will skeletal muscle (p 489), give a sustained (tetanic) contraction. The cadiac contractions are always intermittent. Summation of contractions and tetanius cannot take place in heart muscle. This fact is dependent upon the absolute refractory period and ensures a pause between contractions, an essen tital requirement in order that the heart can perform its pumplike function. The heart muscle must first cease to contract and recover at least

part of its excitability before another contraction can be evoked

Extrasystoles or premature beats of the ventricle. If, when the heart is beating naturally, the ventricle is stimulated during its relaxation phase, a contraction occurs which varies directly in strength with the excitability of the muscle; this, as just mentioned, rises progressively throughout the relative refractory period. The maximal response is therefore obtained at the end of the latter. The response caused by the artificial stimulus is called a premature beat or extrasystole. Premature beats occur in the human heart as a result of some abnormal stimulation (p. 184), such as by some chemical agent carried in the

Fig. 4.34 Extrasystoles Record of contractions of frog's ventricles Upper row of arrows and dotted lines indicate the arrival of the auricular impulse in the ventricular muscle. Slanting arrows in this and succeeding tracings indicate the application of the artificial stimulus. In tracings 1, 2, and 3, the artificial stimulus is applied during the contraction of the muscle, and is therefore ineffective (absolute refractory period). In 4, 5, and 6, the stimulus falls in the relative refractory period and gives rise to an extrasystole, its amplitude being greater the later in the relaxation phase that the stimulus is applied. A long pause follows the extrasystole because the auricular impulse reaches the ventricular muscle during the refractory phase, and is therefore ineffective. E = extrasystole.



blood stream. The excessive use of tobacco is not infrequently a cause. An unusually long resting interval known as the compensatory pause follows the extrasystole (see Fig. 4.34). The pause is caused in the following way. The impulses causing the normal heart beat originate at regular intervals in the upper part of the right auricle and, spreading downward, excite the rest of the heart. When an extrasystole of the ventricle occurs, the next regular impulse from the auricle usually reaches the ventricle while it is absolutely refractory—that is, while it is still in the contracted state as a result of the abnormal stimulus. The muscle therefore fails to respond. Not until the next normal impulse arrives is the heart capable of contracting.

3. Conductivity. We have seen that, as a result of the fusion of branches or slips of one fiber with those of its neighbors, the heart muscle is a continuous protoplasmic mass. When, therefore, one part of the muscle is stimulated, the excitatory process spreads without interruption to all its parts. In the mammalian heart, the property of

rapid conduction has been developed to a very high degree in a system of specialized tissue in the interior of the heart. This system will be dealt with shortly (p. 165)

#### Perfusion of the Excised Heart

Under appropriate conditions the excised mammalian heart will continue to beat for several hours. Even the human heart has been revived after death, and made to beat rhythmically for a time after its removal from the body.

In the perfusion of the excised heart, a fluid resembling the plasma, in so far as its inorganic constituents are concerned, must be employed. The perfusion fluid should be delivered under pressure to the blood vessels of the heart (coronary system, p. 204) in order that it shall reach all parts of the cardiac muscle, the temperature of the fluid should be that of a warm blooded animal, namely, about 98 6° F. The perfusion is performed in the following way A cannula is used into the aorta and connected by rubber tubing to a reservoir filled with the perfusion fluid and placed at a height of about 5 feet above the level of the heart, the fluid is thus driven into the coronary arteries which arise from the root of the aorta just beyond the semilunar valves. The aortic valves are closed by the pressure of the fluid so that none enters the ventricle directly. After passing through the coronary system (arteries, arterioles, capillaines, and veins), the fluid enters the chambers of the heart from which it is allowed to escape through knife cuts made in the walls of the ventricles (see Fig. 4.35)

The heart muscle must be supplied with oxygen. This is accomplished by bubbling oxygen or simply air through the fluid in the reservoir.

It is of the utmost importance that the perfusion fluid contains the three inorganic cations Na<sup>+</sup>, Ca<sup>++</sup>, and K<sup>+</sup>, in properly balanced proportions. This fundamental fact was first demonstrated many years ago by Sidney Ringer an English physiologist (see Fig. 4.36). Experimenting with the excised frog's heart, he found that the heart beat could be maintained for a short time only, when physiological saline (0.6 percent sodium chloride solution) was used. The addition of a very small amount of criteium restored the beat, but the pauses between beats (diastolic periods) became shorter and shorter, and the heart came to a standstill in a firmly contracted state, this condition is called calcium rigor. When potassium instead of calcium was added to the physiological saline, the diastolic periods became progressively

longer, and the heart ceased to beat in a fully relaxed state (diastole). This effect is referred to as potassium inhibition. A fluid prepared to contain sodium, calcium, and potassium chloride in suitable proportions (sodium chloride 0.6 percent for the frog heart or around 0.9 percent for the mammalian heart, with the other salts in the proportions shown in Table 4-1) will sustain the beat for long periods. This is known as Ringer's solution. If the perfusion fluid contains calcium in excess, the effect is the same as if it contained calcium in normal concentration but lacked potassium; calcium rigor ensues. On the other hand, when potassium is in excess the effect is the same as when calcium is lacking and potassium present in the correct concentration; potassium inhibition is induced. The effects of these two cations upon the action of the heart muscle are therefore antagonistic, one favoring contraction, the other relaxation. Only when they are present in balanced proportions. as in blood plasma, can the normal rhythm of the heart be sustained.

The reaction of the perfusion fluid is also of prime importance. The optimum reaction is around a pH of 7.4. If the fluid is too alkaline, an effect is produced upon the heart resembling that of an excess of calcium; if the hydrogen ion concentration is too high, it induces an effect similar to that caused by an excess of potassium.

Locke's solution and Tyrode's solution are modifications of Ringer's original fluid. They contain glucose, added for the purpose of furnishing fuel for the heart muscle, and sodium bicarbonate (0.02 percent) which maintains the fluid




Fig. 4.35 Showing apparatus for the perfusion of the isolated mammalian heart A, tube conducting Ringer's solution to the heart The lower end of this tube is tied into the aorta so that fluid is distributed through the coronary circulation to all parts of the cardiac muscle. B. waterjacket through which warm water (37.5° C) circulates, C, thermometer.

at optimum hydrogen ion concentration. Tyrode's solution contains,




Fig 4.36 Tracings showing the effects of sodium, potassium, and calcium upon the action of the heart. Not until all three cations are in the proportions present in Ringer's solution (lowest tracing) is cardiac behavior normal in character (kindness of Dr. K. G. Watkin).

as well as bicarbonate, the phosphate buffers—disodium and monosodium phosphates (Na<sub>2</sub>HPO<sub>4</sub> and NaH<sub>2</sub>PO<sub>4</sub>). See Table 4-1.

Until quite recently it had been thought that glucose constituted the chief fuel of cardiac muscle. It now appears that the heart uses glucose to a minor extent, its main fuel is lactic acid Lactic acid is formed during muscular contraction from the breakdown of glycogen (p. 368). It is also formed from glucose in the blood, and in the lungs at all times. The lactic acid so produced is utilized by the heart muscle to furnish energy for its contraction.

The hearts of lower orders, such as frog, are not provided with a special circulating system (coronary system) as is the heart of man or other mammals The frog's heart will therefore continue to beat for

Table 4-1

Percentage composition of Ringer's, Locke's, and Tyrode's solutions, used in perfusion of the mammalian heart

	Ringer's solution	Locke's solution	Tyrode's solution
Sodium chloride	0.900	0 900	0.80
Potassium chloride	0 040	0.042	0.02
Calcium chloride	0.025	0 024	0.02
Magnesium chloride	_	_	0.01
Sodium bicarbonate	0.020	0.020	0 10
Monosodium phosphate	0 001		0.005
Glucose	-	0.10	0.10

some time if simply immersed in Ringer's or other physiological solution.

# The Origin and Conduction of the Heartbeat

In the slowly beating heart of the frog, each beat may be seen to arise at the junction of the great veins with the sinus venosus-a chamber situated between these veins and the right auricle (see Fig. 4.37). The sinus venosus is present in the mammalian embryo but disappears during development. In the heart of the frog or the turtle, the beat spreads from the sinus venosus through the muscle of the auricles and ventricles When a ligature is drawn tightly around the frog's heart at the junction of the sinus with the auricle (called the 1st Stannius liga-ture), or at any lower level, the impulse through the muscle is blocked and the heart below the block ceases for a time to beat. In the adult mammalian heart, a ring of fibrous tissue is interposed between the auricles and the ventricles: the beat therefore cannot be transmitted from the upper to the lower cardiac chambers by the ordinary muscular tissue of the heart. But a system of specialized tissue—the junc-tional or nodal tissues—has been developed in the mammalian heart, which possesses the property of thythmical impulse formation and conductivity to a higher degree than does the muscle composing the aurícular and ventricular walls. The beat is generated in a small mass of tissue belonging to this system, situated in the wall of the right auricle near its junction with the superior vena cava. This circumscribed




Fig 4 37 The frogs heart, upper drawing, from in front, lower drawing from behind

Inferior vena cava

knot of tissue differs in structure from the surrounding auricular muscle, being composed of muscle cells of a primitive type (Purkinje fibers), together with nerve fibers and a few nerve cells. It is called the sinoauricular or sinoatrial (S-A) node (Fig. 4.38), and is believed to be derived from the wall of the sinus venosus of the embryonic heart where this chamber joined the auricle. Heating or cooling the S-A




Fig 4.38 The conducting (junctional) tissues of the heart. Wall of the right ventricle cut away exposing the septum

node accelerates or slows, respectively, the rate of the heart From such and other facts, it seems quite certain that the impulse which causes the beat of the heart is generated within this node of special tissue. The sinoauricular node is therefore commonly referred to as the pacemaker of the heart. The impulse spreads as a wave—the excitation wave—in all directions through the muscle of the auricles. The nature of the excitation wave cannot be defined precisely, but at any rate it is known to be accompanied by an electrical change, and is followed immediately (within 0.02 second) by contraction of the cardiac muscle.

The remainder of the specialized conducting system of the mammalian heart consists of the auriculo- (atrio-) ventricular (A-V)

node, the auriculo- (atrio ) ventricular bundle and the Purkinje network

The auriculoventricular node lies in the lower part of the interauricular system. It is composed of tissue similar to that of the sinoduricular node. The auriculoventricular bundle arises from the auriculoventricular node and, after descending for a short distance, divides into two branches—one to each ventricle. Each branch descends beneath the endocardium covering the interventricular septum and, after giving off twigs to the papillary muscles, breaks up into a rich network of fine filaments—the Purkinje network—which is continued beneath the endocardium of the ventricular wall to the base of the ventricules.

The muscle fibers composing the strands of the Purkinje network are of essentially the same character as those forming the rest of the junctional ussues like the latter, they contain a relatively large quantity of sarcoplasm but few myofibrils. They are, however, of much larger diameter than the fibers of the S-A and A-V nodes and the A V bundle. They merge with the muscle fibers forming the ven tricular walls and through them the excitation wave or impulse to contract is carried to the latter.

The impulses are discharged rhythmically from the S A node, and conducted at a uniform rate through the fibers of the auricular muscle. The impulse is then picked up by the A-V node, and conducted to the ventricular muscle by the bundle and its ramifications. Each part of the muscle of the ventricel is excited when the impulse arrives at the part of the Purkinje system immediately underlying it. The impulse, normally is not conducted from point to point in the ventricle through muscular tissue, but follows this specialized pathway. Every muscle fiber of the ventricle is believed to receive a filament of the Purkinje network. The rate of conduction through ventricular muscle (which may in certain abnormal conditions conduct the impulses) is around 500 mm per second, whereas the rate through the specialized tissue is about 5000 mm per second.

The junctional or nodal tissues of the heart may be compared to a radio system, the S A node to the broadcasting station, and the A-V node to a receiving center which, after picking up the message, transmits it by a telephonic system. In the heart, the bundle and its branches and the terminal network of Purkinje correspond to the telephone wires.

It should be noted that the SA and A-V nodes are not connected by a tract of specialized tissue, the impulse in this part of its course is conducted solely through the ordinary auricular muscle. Also, the A-V bundle is the only pathway along which the impulse can reach the ventricles, for all other tissue in the mammalian heart joining the auricles and ventricles is of a fibrous nature and incapable of conducting an impulse. This slender strand of tissue is therefore a very important and vulnerable point in the conducting system of the heart. It is a "bottleneck" which the impulses must traverse. If interrupted, either experimentally in animals or by disease in man, the path of the impulse from auricle to ventricle is completely and permanently blocked. The ventricle does not, however, cease to beat; for the specialized tissue below the block also possesses, though to a less degree than does the S-A node or the A-V node, the power of generating rhythmical impulses. The ventricular muscle excited by impulses arising below the block then beats at the relatively slow rate of about 35 per minute. The auricle being still under the control of the S-A node beats at the usual rate of 70 per minute. This dissociation of the rhythms of the auricles and ventricles, which follows interruption of the A-V bundle above its point of division into the two main branches, is called complete heart block or auriculoventricular block (see p. 183).

In other instances one or other branch of the A-V bundle may be interrupted by disease. The defect in conduction caused in this way is called bundle branch block or intraventricular block.

### The Action of the Heart, the Cardiac Cycle

The action or beat of the heart consists of a succession of events which follow one another with great rapidity. It is therefore impossible, except by the use of special apparatus and improved methods, to analyze the beat into its different components. Starting with any given event in cardiac action (e.g., contraction of the auricle or of the ventricle), the series of changes which takes place in the heart until the first-noted event is repeated, is called a cardiac cycle. The cardiac cycle in the human subject, when the heart is beating at the usual rate of about 70 per minute, is \$\frac{9}{2}\$ (0.85) second. It is shorter than this, of course, when the heart rate is more rapid. Contraction of the auricular or of

<sup>&</sup>lt;sup>8</sup> The most common condition affecting the A-V bundle is gradual hardening and narrowing of the coronary arteries which, by interfering with the blood supply to this vital ussue, convert it to a fibrous (nonconducting) structure.




Fig 439 Diagram to show the different phases of the cardiac cycle A auri cles filling valves between auricles and ventricles closed ventricle relaxed valves at arterial openings closed (period of isometric relaxation) B the valves between the auricles and ventricles have opened and blood is pouring into the ventricles which are still relaxed (period of rapid inflow) C, the auricles and ventricles are full there is little or no movement of the blood (period of di astasis) D the auricles contract (auricular systole) and empty into the ven tricles which are over filled with consequent further stretching of the ventricu lar muscle fibers E the ventricles immediately contract (period of isometric contraction) and close the openings into the auricles (auriculoventricular orifices guarded by the corresponding valves) The pressure then rises rapidly in the ventricles and when sufficiently high to overcome the pressure in the pulmo nary artery (and in the aorta) the valves guarding the onfices of the arteries open In this sketch the valves are still closed the ventricle is therefore a com pletely closed cavity F the arterial valves have been forced open and the blood is now being driven into the pulmonary system (or aorta) (period of ejection) The left side of the heart is drawn as though empty in order to show more clearly the positions of the valves but it must be remembered that whatever is occurring on the right side of the heart is occurring at the same instant on the left side The figures refer to the duration of the periods at a heart rate of 70 beats per

the ventricular muscle is called systole so we speak of auricular systole or ventricular systole, respectively Auricular diastole and ventricular diastole are corresponding terms which refer to the relaxation phase of the respective chambers When the word systole or diastole is used alone, it is the systole or diastole of the ventricle that is meant

The phases of the cardiac cycle are shown diagrammatically in Figures 4 39, 4 40, and 4 41 In animals the cardiac cycle is most readily analyzed into its several phases by means of records, taken




Fig. 4.40 Intraventicular, intra-aortic and intra-auricular pressure curves from heart of dog together with an electrocardiogram. The vertical lines pass through synchronous points in the four curves. Detween the closure of the auriculoventricular valves and the opening of the aortic valves is the isometric period of contraction. From the opening of the aortic valves to the closure of these valves js the period of ejection, and from the latter event to the opening of the A-V valves is the period of isometric relaxation. The divisions in the line at the bottom of the diagram indicate  $\frac{1}{2}$  second.

simultaneously, of the pressure changes occurring, respectively, in an auricle, a ventricle, and the aorta. Instruments specially designed to record rapid variations in pressure are placed in these situations. The records are taken simultaneously so that synchronous events in the curve can be compared.

We shall now describe in their proper sequence the several events in the cardiac cycle. The part of the cycle at which the description starts does not matter. The period immediately following the contraction (systole) of the ventricle will be chosen. (See Fig. 4.39.) At this time the ventricles are closed cavities. The valves guarding the orifices of the aorta and the pulmonary artery are closed, as are also those guarding the auriculoventricular orifices. The ventricular muscle is




Fig. 4.41. Diagram of the card ac cycle. The black part of the outer circle represents ventricular systole, the white part ventricular diastole, division marks and cate  $\frac{1}{10}$  second.

now relaxing but since no blood can enter the ventricles from the auricles to stretch the muscle fibers these are of the same length as during the preceding systole. This phase of the cardiac cycle at the commencement of ventricular diastole is therefore called the period of isometric (Gk isos same + metron measure) relaxation (Figs 440 and 441) it has a duration (at a heart rate of 70 per minute) of 06 to 08 second. At its termination the auriculoventricular valves open and the blood flows rapidly into the ventricles from the auricles. This phase which lasts for about 0.1 second is called the period of rapid filling of the ventricles.

As the ventrules fill the flow of venous blood becomes slower and then virtually ceases This interval following the period of rapid filling is termed the period of diastasis. It varies considerably in length, being longer when the heart is beating slowly. During rapid action of the heart it may disappear entirely. Auricular systole—that is, auricular contraction—follows the period of diastasis; it gives an impetus to the blood flowing from the auricles into the ventricles. Auricular systole has a duration of about 0.1 second; at its termination the ventricles contract.

It should be emphasized that throughout all the phases of the cardiac cycle just described (vis., isometric relaxation, rapid filling, diastasis, and auricular systole) the ventricles are relaxing or are completely relaxed—that is, these events occur during the resting phase or diastole of the ventricle.

Ventricular systole which, as just mentioned, follows auricular systole is divided into two phases: the period of isometric contraction and the period of ejection. The rise in pressure within the ventricles when they contract firmly closes the auriculoventricular valves. The aortic and pulmonary valves, which were brought into apposition at the end of the preceding systole, have not yet opened; they are held closed by the higher pressure of blood in the aorta and pulmonary arteries. The ventricle being thus again a closed cavity but filled with blood which, of course, is incompressible, the muscle fibers obviously cannot shorten during this phase of systole. They contract isometrically. This phase, therefore, is appropriately called the period of isometric contraction. The pressure within the ventricles continues to rise rapidly, however, during this time. When it reaches a height sufficient to overcome the diastolic pressure in the aorta (about 80 mm Hg) and pulmonary arteries (about 15 mm Hg), the valves guarding these vessels are thrown wide open, and the ventricular contents ejected. The ejection period lasts for about 0.3 second (with heart rate of 70). It terminates with relaxation of the ventricular muscle. The pressure within the ventricles now falls below that in the large arteries, and the aortic and pulmonary valves close again as a result. The next phase of the heart's action is the period of isometric relaxation—the phase with which this description started. The cardiac cycle has been completed.

The intra-auricular pressure curve, it will be seen, consists of three positive waves, a, c, and v, and three depressions, x, x, x and y. The wave a is due to auricular systole, the depression x is due to auricular relaxation (auricular diastole). The wave c is caused by ventricular contraction which bulges the thin auriculoventricular valves into the auricle. The depression x is dependent also on the ventricular systole.

factors During auricular systole, the A-V valves and, during the ventricular systole, the pulmonary and aortic valves are held open by the blood flowing through the respective openings. When auricular systole comes to an end, eddy currents arising in the ventricle bring the leaflets of the A-V valves gently into position. The pressure created immediately thereafter by ventricular systole closes them firmly. The closure of the aortic and pulmonary valves is caused in a similar manner, namely, by the higher pressure in the arteries (aorta and pulmonary artery) than in the ventricle when the ventricular muscle commences to relax.

The heart sounds. If one places his ear, or the receiver of an instrument called a stethoscope upon a man's chest a little below the left nipple, the beat of the heart can be clearly heard. Two sounds, separated by a short silent interval and followed by a longer pause, can be distinguished. They are called, respectively, the first and second heart sounds, and have been compared to the syllables 'lub' 'dup' The first sound ('lub') is louder, of lower pitch, and longer than the second ("dup"). The first sound is caused by two factors (a) the contraction of the ventricular muscle and (b) the vibrations set up by the tight closure of the aunculoventricular (tricuspid and mitral) valves, and of the chordae tendineae as they are put upon the stretch by the rising pressure in the ventricles.

The second sound is caused by the closure of the aortic and pulmonary valves

Valvular disease. The function of one or other valve of the heart may be interfered with as a result of some inflammatory disease (e.g., acute rheumatic fever or septicemia) The affected valve becomes de formed or partially destroyed so that its cusps do not come accurately into apposition to prevent leakage. The valve is then called incompetent and the leakage of blood is referred to as regurgatation. For example, when the aortic valves fail to close properly at the end of ven tricular systole, blood flows from the aorta into the left ventricle during disatole, since the pressure in the aorta is higher than that in the relaxing ventricle, when the mitral valve is incompetent, blood is driven into the left auricle during ventricular systole. Now, these reverse movements of the blood cause abnormal sounds or murmurs to the heard. In aortic incompetence, the clear second sound of the normal heart is replaced by a murmur, and this is often continued through

Any skeletal muscle causes a faint sound when it contracts as a result of the vibration of its fibers. This can be verified by listening over the contracting biceps of the arm with a stethoscope.

a part of, and sometimes throughout, the diastolic period; that is, a blowing or rushing sound is heard during the silent period which normally follows the second sound and separates it from the first sound of the next heart cycle. In mitral or tricuspid incompetence, the abnormal sound is heard during ventricular systole, and alters or replaces the normal first sound.

In other instances the diseased valve loses its resilient character, becoming stiff and its surface roughened. The cusps are not brushed aside by the current of blood but, projecting into the stream, offer an impediment. The orifice is therefore narrowed. This valvular condition is called stenosis. Thus, according to which valve is affected, mitral, aortic, pulmonary, or tricuspid stenosis is spoken of. Owing to the narrowing of the orifice, the blood passes through it at a higher velocity than usual The blood flowing at high velocity over the roughened and irregular cusps causes a murmur. When an auriculoventricular valve is affected, the murmur is heard during those times when the blood normally enters the ventricle at highest velocity, namely, during the period of rapid filling—i.e., early in diastole, or during auricular systole which is just before ventricular systole (presystolic murmur). In aortic stenosis the murmur is heard during the ejection of blood from the ventricle.

It must be evident that the rigidity and deformation of a stenosed valve is likely to render it incompetent as well. Two sounds will then be present, one being due to the stenosis, the other to regurgitation.

A diseased valve increases the work of the heart. The heart is at a mechanical disadvantage. In aortic regurgitation, for example, the blood must eject not only the blood that it receives from the auricles but also that which leaks back from the aorta Moreover, the disease which attacked the valve may have also injured the heart muscle. In order to carry on its work efficiently, the cardiac muscle fibers must become larger—longer and thicker—the chambers of the heart thus become enlarged and the ventricular walls much thicker. Dilation of the heart and increased bulk of the heart muscle (hypertrophy) is, therefore, a consequence of valvular disease. Such adjustments, or compensation, as they are called, may enable a person with valvular disease of the heart to enjoy good health for years and suffer little inconvenience from the mechanical defect.

Electrical changes in the heart. Contracting muscle is relatively negative (electrically) to resting muscle. The electrical changes in skeletal muscles are described on page 494. The same principles apply, in general, to heart muscle. When the excitation wave spreads

through the heart, changes in electrical potential are created which may be recorded in animals by placing electrodes directly on the sur face of the heart and connecting them through wires to a galvanom eter It is, of course, impossible to take records (electrograms, as they are called) in this way of the changes in electrical potential occurring





Fig 4.43 Diagram illustrating the electric field surrounding the heart and the relation of the three stand ard electrocardiographic leads (After Pardee)

in the human heart Neverthe less the blood and tissues of the body have a relatively high electrical conductivity (Fig 4 43) the beating heart may, therefore, be looked upon as an electric generator immersed in a con ducting medium, and the elec tric currents (action currents) produced during the cardiac cycle may be "tapped,' as it were, by connecting two parts of the body with a sensitive galva nometer Einthoven, a Dutch physiologist, was the first to de vise a galvanometer for this pur pose The moving part or indi cator of the galvanometer con sists of a very fine fiber of quartz glass through which the cardiac action currents are transmitted In order to render it conductive the fiber is coated with a silver

deposit (as in the backing of a mirror), and suspended between the poles of an electromagnet. The fiber moves outward (away from the arch of the magnet) or inward according to whether the cardiac current at the moment is being transmitted in an upward or a downward direction. By means of a projection system and a camera of special design, the movements are photographed. The record thus obtained is called an electrocardiogram (abbreviated ECG), and the recording instrument—that is, the galvanometer together with the projecting and photographing apparatus—is known as the electrocardiograph (see Fig. 4.44). The parts of the body most usually connected to the instrument are the two hands (or arms) or the right hand (or arm) and left foot (or leg). In order to obtain a record of the electrical changes occurring in the heart, these members are joined through the



photographic paper moving behind the lens, Cl, which narrows the shadow, ST.S.H., of the string ST. to a thin vertical line and focuses it upon the sensitive paper. By means of ruled etchings upon the lens, horizontal lines I and \(\frac{1}{2}\) mm apart are thrown upon the record, the heights of the waves in the electrocardiogram can thus be measured and the electromotive force in each wave calculated. T is a rotating toothed disc which breaks the light beam at regular intervals of \(\frac{1}{2}\) or \(\frac{1}{2}\) or \(\frac{1}{2}\) seconds and by throwing vertical shadow lines upon the record serves as a time marker. O and C indicate the positions of the lenses in the projection system; S and N are the poles of the electromagnet, L is the source of light. The current from the subject passes through the control box below. Switches on the dial enable a record from any lead to be taken RH, LH, and LL represent the three standard leads—right hand, left hand, and left leg (Kindness of Dr. L. N. Katz.)

instrument in any one of the following combinations (see Figs. 4.43 and 4.45). Each combination is referred to as a *lead*. Thus:

Right hand (or arm) and left hand (or arm) is called lead I. Right hand (or arm) and left foot (or leg) is called lead II. Left hand (or arm) and left foot (or leg) is called lead III.

A switch on the instrument board enables records to be taken in succession from each of the three leads.

A fourth lead (lead IV or chest lead), in which one electrode is placed on the chest over the heart and the other on the left foot or ankle, has been introduced more recently, and is usually taken in series with the other (so-called standard) leads.




Fig 4.45 Showing the three standard leads of the electrocardiograph and a chest lead. Contacts are made with the skin by means of suction cups—small rubber builb and metal bell—the skin being first smeared with a paste containing electrolyte in order to ensure good conduction between the skin and the metal.

The normal electrocardiogram shows five waves or deflections, known simply by the letters P, Q, R, S, and T (Figs 4 46 and 4 47). The P, R, and T waves are positive—that is, above the base line of the record, the Q and S waves are negative P and T are blunt waves of relatively low amplitude. The R wave is a tall spike, it is the most conspicuous wave in the tracing and represents the highest voltage developed by the contracting ventricular muscle. The P wave is caused by the spread of the excitation wave over the auricle—that is, it precedes auricular systole by a fraction of a second. The Q, R, and S.




Fig 446 Normal electrocardiogram taken in lead I (Courtesy of Dr John Hepburn )

deflections are produced during the transmission of the excitation wave or impulse through the junctional tissues—auriculoventricular bundle and its branches, and the Purkinje network. The actual contraction of the ventricular muscle commences a fraction of a second after the beginning of the R wave. The T wave is recorded while the ventricle is relaxing (see also Fig. 4.40, p. 171)

The electrocardiograph is one of the most valuable instruments that the physician possesses for the investigation of the various abnormal heart conditions It is now possible with its help to study diseases of the heart with a precision which hitherto was impossible An electrocardiogram is now taken as a matter of routine, being almost as commonplace a procedure as the determination of the blood pressure Important information is given by the electrocardiogram concerning the state of the conducting system of the heart and of the coronary system Since the P wave is caused by the excitation wave




Fig. 4.47 The waves of the electrocardiogram enlarged and represented diagrammatically

passing over the auricle and the R wave by the excitation of the ventricle, the time interval between the commencement of P and the commencement of R (P-R interval) gives the conduction time from auricle to ventricle—i.e., over the A-V bundle. The normal time does not exceed \( \frac{1}{2} \) (0.2) second. It is usually around 0.12 second. Lengthening of the interval indicates depression of conductivity through the auriculoventricular connections—i.e., incipient or partial heart block. Disease of one or other branch of the bundle (intraventricular block) is also readily detected by means of the electrocardiograph. Marked distortion of the record occurs in this condition.

The venous pulse. The arterial pulse is not transmitted through the capillaries to the venous system (p 148). The peripheral veins, therefore, do not pulsate, but the *µggular vein* is in direct communication with the right auricle, and any change in intra-auricular pressure causes a reflux of blood and volume changes in the blood within this vein which can be recorded clinically by means of suitable apparatus the receiver of which is placed over the vein at the root of the neck just above the clavicle (collar bone).

The jugular tracing shows a series of waves resembling, and having the same significance as, those already described for the intra-auricular pressure curve (p 173) Since the a wave is due to auricular systole and the c wave to the commencement of ventricular systole (which causes the bulging into the auricle of the closed auriculoventricular valve), the length of the a-c interval gives the time of conduction of the impulse from auricle to ventricle. This interval, therefore, corresponds to the P-R interval of the electrocardiogram and, like the latter, does not normally exceed  $\frac{1}{2}$  (0.2) second




Fig 448 A normal venous pulse and a radial pulse tracing. The dotted lines join synchronous points in the two tracings. (After Lewis.)

The waves of the jugular record are not always easy to identify unless an arterial tracing (carotid or radial) is taken simultaneously with it. The instrument employed for taking synchronous records of jugular and arterial pulses is called a polygraph. The upstroke of the carotid tracing occurs simultaneously with the jugular c wave, which

can thus readily be identified. The positive wave immediately preceding the c wave must therefore be the a wave, and the positive wave immediately following c is the v or stasis wave caused by the filling of the auricle (see p. 173 and Fig. 4.48).

The apex beat. The contraction of the heart causes a slight impact

The apex beat. The contraction of the heart causes a slight impact upon the chest wall which can be felt by placing the fingers on the skin between the fifth and sixth ribs, and in a line with the center of the left collar bone—that is, about an inch below the left nipple (in the male). This impact is called the apex beat. The slight outward movement of this area of the chest wall can also be seen if the bare chest is looked at from one side, in a good light. The movement is caused by the pressure exerted by the left ventricle during its contraction. All dimensions of the ventricles are reduced when they contract, but no change in the level of the apex of the heart occurs. The base of the heart, therefore, must move toward the apex. The auricles and great vessels are thus elongated; they cannot move downward as a whole, for they are fixed above. At the same time, the heart rotates to the right, bringing the wall of the left ventricle and particularly that part forming the apex, against the chest wall. During ventricular diastole the base of the heart moves upward again. The up and down movement of the auriculoventricular groove (the junction between the auricles and ventricles) during the cardiac cycle can be seen quite plainly in the exposed heart of an animal.

Disorders of the heart beat. Certain abnormalities of cardiac rhythm in the human subject (e.g., heart block and extrasystoles have already been mentioned. These, and some of the other cardiac irregularities, will now be briefly described.

Hear block or auriculoventricular block. This results from disease, usually a degenerative nature, affecting the auriculoventricular bundle. Interference with the blood supply to the heart, including the specialized tissue, as a result of 'hardening' (sclerosis) with narrowing of the coronary vessels is a common cause of defective conduction through the bundle. The disease may completely interrupt the passage of impulses from auricles to ventricles; the auricles and ventricles then beat quite independently. In such an event, the radial pulse is around 35 per minute while the a waves (due to auricular contractions) in the venous pulse occur at the rate of 70 per minute. There is no regular time relationship between the auricular and ventricular waves of the venous pulse; corresponding abnormalities are seen in the electrocardiogram, there being about two P (auricular) deflections to each R (ventricular) deflection.

In less severe grades of the disease, the conduction time is lengthened, but no actual block exists. The P-R intervals of the electrocardiogram and the a-c intervals of the jugular tracing are prolonged. In other instances, block occurs intermittently. That is, after every two, three, or four auricular beats the impulse fails to get through to the ventricle which, in consequence, misses a beat. Three stages in disease




Fig 4.49 1, heart block. Note absence of the normal relationship between the auricular (P) and the ventricular waves (Q, R, and S). 2, a ventricular extrasystole. Note the long pause following it, 3, ventricular paroxysmal tachycardia, actually a series of extrasystoles. 4, auricular fibrillation. Waves marked f replace the normal P waves (1, 2, and 3 by the couriesy of Dr. John Hepburn, 4 from Lewis)

of the A-V bundle are therefore recognized, namely: (a) delayed conduction, (b) missed beats or incomplete heart block, and (c) complete heart block (Fig. 4.49, 1).

Extrasystoles or premature beats (see also p. 161). The abnormal impulse which excites the heart may arise in the auricle, the A-V node, or in the special conducting tissue of the ventricle. Thus, three types of extrasystole occur: auricular, nodal, and ventricular. In the auricular type, the impulses arise in some part of the auricle other than the S-A node; each of these impulses spreads to the ventricle, excites it in the usual way, and causes a premature beat of the whole heart. There is, as a rule, no compensatory pause (p. 161). In the nodal type, the impulse arising in the A-V node passes upward

through the auricular muscle and downward to the ventricle. Both chambers, therefore, beat almost or quite simultaneously. In the ventricular type, which has already been described on page 161, the premature beat is followed by a compensatory pause. In this variety, the abnormal impulse, since it excites one ventricle before the other, disturbs the electrical balance of the heart. The QRS group of waves of the electrocardiogram is therefore distorted (Fig. 4.49, 2).

Paroxysmal tachycardia. This name is given to a cardiac affection in which the heart suddenly commences to beat at an extraordinarily rapid rate and continues to do so for a period which varies in length in different instances from a few seconds to several hours or even days. The heart rate is from 150 to 250 per minute. The rapid beating is caused by impulses arising in some part of the heart other than the S-A node. They may be generated in the auricle, the A-V node, or in the ventricular part of the special conducting system. The condition therefore consists, actually, of a series of rapidly recurring extrasystoles (Fig. 4.49, 3).

Auricular flutter. In this heart condition the auricle beats more rapidly than in paroxysmal tachycardia—namely, from 250 to 300 per minute. Attacks of flutter are also usually of longer duration than those of paroxysmal tachycardia. Furthermore, the A-V bundle, even though healthy itself, is unable to transmit impulses at as high a frequency as they are produced by the fluttering auricle. Only a proportion of the impulses gets through; the ventricle, therefore, beats at a slower rate than the auricle. In other words, there is a relative or functional heart block. The ratio of auricular to ventricular beats is usually 3:1 or 2:1.

Auricular fibrillation. The auricle in this disorder does not contract as a whole, nor in an effective manner. Its walls twitch rapidly and incessantly at a rate of from 400 to 600 per minute. The auricle is never emptied by any effective contraction of its walls; the blood flows into the ventricle unaided by auricular systole. The a waves are therefore absent from the venous pulse, and the P waves from the electrocardiogram; they are replaced in either record by numerous small fluctuations called f waves (Fig. 4.49, 4). The auriculoventricular connections (A-V bundle) are capable of transmitting only a fraction of the impulses arising in the auricular muscle. Those that do reach the ventricle are irregular in time, and induce contractions of the ventricle of variable strength. The pulse is, therefore, very irregular and usually rapid. Auricular fibrillation is seen very frequently when the heart, as a result of disease of the heart muscle, is

184

weakening and performing its work with difficulty. The serious nature of the condition is not due to any direct effect upon the dynamics of the circulation resulting from the absence of auricular systole, but rather to the irregular and rapid excitation of the ventricular muscle. The drug most commonly used in the treatment of auricular fibrillation is digitalis, which acts chiefly by depressing the conductivity of the A-V bundle. The ventricle is therefore shielded from many of the impulses arising in the fibrillating auricles; the pulse usually becomes slower and more regular in consequence; the general circulation is greatly improved.

## The Regulation of the Heart's Action

The heart rate. The heart rate is more rapid in small animals than in large. The heart of the canary, for example, beats 1000 times per minute, whereas the average heart rate of the elephant is 25 per minute. In adult man, at rest, the usual rate is about 70 per minute, but a rate of 75 or of 60 per minute is not unusual in perfectly healthy persons. Athletes have, as a rule, a slower resting heart rate than have persons leading a sedentary life; the pulse rate of some famous runners is from 50 to 60 per minute, and it may show little increase in frequency during physical exercise. The heart rate diminishes progressively from birth, when it is around 130 per minute, to adolescence. The heart is accelerated in muscular exercise, by emotional excitement, at high environmental temperatures, and during digestion. Among abnormal conditions which cause an increase in cardiac rate are hemorrhage, surgical shock, fever, hyperthyroidism, and certain cardiac disorders (p. 185).


Tachycardia is a general term meaning increased heart rate. Bradycardia means an unusually slow rate of the heart

The cardiac nerves. Although the heart will continue to beat thythmically after all nervous connections have been cut, in the ordinary conditions of life, it is under the constant influence of nervous impulses, discharged from centers in the brain. These centers are influenced in turn by impulses conveyed to them from various parts of the body along afterent nerves (p. 460), as well as by impulses received from the higher centers (psychic, emotional) of the brain. It is common knowledge that emotional disturbances—anger, fear, excitement, etc.—exert a profound influence upon the pulse rate. The efterent cardiac nerves—ie, the nerves which transmit impulses from the nervous centers to the heart—are the vagus and

the accelerator (or augmentor) nerves (Pl. 5A). Thus, through its nervous connections, the heart may undergo slowing or acceleration as a result of the stimulation of nerve endings situated in regions remote from the heart itself. These reflex mechanisms are dealt with in greater detail on page 190.

The vagus nerves arise, one on each side, from the medulla oblongata (p. 546). The great majority of the fibers of which they are composed belong to the parasympathetic division of the involuntary or autonomic nervous system. The vagus branches are very widely distributed, passing to the bronchi, lungs, stomach, and small intestine, as well as to the heart. The fibers going to the heart separate from the main trunk of the nerve in the upper and lower parts of the neck. They terminate around nerve cells (ganglion cells) situations.

Fig. 4.50 showing the effect of an electric stimulus applied to the right vagus nerve Heavy white line indicates the commencement and duration of the stimulus. Note the escape of the heart from the vagal stimulation.



ated in the auricular muscle. The impulses are relayed by the axons of the ganglion cells to the sinoauricular and auriculoventricular nodes. The fibers in the first link of the nerve path—that b, from the center in the medulla (generally known as the cardoinhibitory center) to the ganglion cells—are called preganglionic fibers. The axons of the ganglion cells, which constitute the second link, are called postganglionic fibers. The right vagus is distributed mainly to the S-A node; it affects the rate of the heart to a somewhat greater extent than does the left nerve. The left vagus is distributed chiefly to the A-V node; its effect upon auriculoventricular conduction is greater than that exerted by the nerve of the right side. (See below.) The distributions of the two nerves overlap to a considerable extent, however, since both the S-A node and the A-V node receive fibers from both.

Action of the vagus nerves. The vagus is the inhibitory nerve of the heart. When it is stimulated, slowing or stoppage of the heart results. Diminished force of the beat usually accompanies the reduction in rate These effects result in a fall in the arterial blood pressure (Fig. 4 50). The vagus exerts no direct effect upon the ventricular muscle. Its action is upon the auricle, and the auriculoventricular bundle; stimulation of the vagus after section of the A-V bundle does not affect the ventricular rate During vagal stimulation excitability of the auricular muscle is reduced, and its refractory period shortened, the rate of conduction from auricle to ventricles—that is, through the auriculoventricular bundle—is depressed. The slowing of the heart following vagal stimulation is due mainly to lengthening of the diastolic period and, when the heart stops, it does so in the diastolic phase. For this reason, the vagus is sometimes called the diastolic nerve of the heart. The action of the vagus upon the heart closely resembles that caused by an excess of potassium (p. 162).

When the heart is arrested by vagal stimulation, it very often com-

When the heart is arrested by vagal stimulation, it very often commences to beat, though the stimulus is still being applied. That is to say, the heart breaks away from the vagal restraint This is spoken of as the "escape of the heart" from vagal inhibition. The first few beats of the heart after it has escaped from vagal control are usually of exceptional force (Fig. 4.50). This is due most probably to the fact that the muscle fibers of the ventricle which, since the latter is fully distended by the venous blood during the period of inhibition, are very strongly stretched, and, therefore, cause an unusually powerful contraction (see Starling's law, p. 193)

The tone of the vagus. When the physiologist says a nerve or muscle possesses tone, he means that it exerts a continuous action. In this sense, the cardiac vagus (or rather its center in the medulla) has tone, it exerts a ceaseless restraint upon the action of the heart, sending a continuous stream of impulses along the vagus nerves which holds the heart in check. When this restraint is removed, as by cutting both vagus nerves, the rate of the heart increases very greatly. This method of demonstrating vagal tone cannot, of course, be used in man Atropine, a drug which temporarily paralyzes the vagus, can, however, be given without danger. When a full dose of this drug is given, the heart rate is more than doubled (150 to 180 beats per minute).

The accelerator or augmentor nerves. These are nerves belonging to the sympathetic division of the involuntary or autonomic nervous system (p 561) Like other pathways of the autonomic nervous system, the pathway for the accelerator impulses to the heart consists of preganglionic and postganglionic fibers. The preganglionic fibers are the axons of cells situated in the lateral horns of spinal gray

matter in the upper five thoracic segments. The postganglionic fibers arise from the stellate ganglion, and the inferior, middle, and superior cervical ganglia of the sympathetic cord; they pass to the heart by the inferior, middle, and superior cardiac nerves (Pl. 5Å). Fibers also pass to the heart from the upper two thoracic ganglia. Stimulation of the accelerator nerves increases the rate of both

Stimulation of the accelerator nerves increases the rate of both auricles and ventricles and increases the force of their contractions. The nerves exert a direct action upon the ventricular muscle. They also enhance A-V conduction. Like the vagus, the accelerators exert a continuous effect upon the cardiac rate—that is, they exercise a tonic action. Consequently, when their influence is removed by excision of the sympathetic ganglia from which they arise, the heart rate slows.

Cardiac reflexes. The stimulation of almost any sensory (afferent) nerve in the body may cause a change in cardac rate. In the frog, merely tapping the abdomen results in slowing of the heart, and it has been shown that in man, the stimulation of abdominal organs by manipulations during surgical operations causes changes in the heart rate. Depending upon the nature of the stimulus and the nerves stimulated, the heart may increase or reduce its rate during an operation. Again, irregularities in the action of the heart may result due to the production, reflexly, of extrasystoles (p. 184). Afferent impulses arising in a diseased organ may also influence the rate of the heart. Cardiac inhibition may be caused by stimulation of the endings of the fifth nerve in the nose, as by some pungent odor, or of afferent fibers in the lungs by irritant vapors. A similar effect may follow pressure upon the outer angle of the eye (oculocardiac reflex). In animals, stimulation of a spinal sensory nerve, such as the sciatic, is not infrequently followed by cardiac acceleration.

Among the afferent fibers which are most effective in bringing about reflex changes in heart rate are those contained within the vagus nerves themselves, for these nerves are made up of fibers which convey impulses to as well as away from the cardioinhibitory center. The afferent vagal fibers end in the heart and in the arch of the aortia (see cardiac depressor nerve, p. 199). The endings in the aortic arch and in the left side of the heart are stimulated by a rise in arterial blood pressure; the impulses increase the tone of the cardioinhibitory center; and cardiac slowing results. Those on the right side of the heart are stimulated by a rise in the pressure of venous blood in the great veins and auricle; they cause acceleration of the heart by depressing the tone of the cardioinhibitory center.

The sinus nerve. Another afferent nerve which, when stimulated causes reflex slowing of the heart is the sinus nerve. Its endings are situated in the walls of the carotid sinus (p 199). The normal stimulus to these nerve endings is a rise in arterial blood pressure.

lus to these nerve endings is a rise in arterial blood pressure. Two well known circulatory phenomena will now be described. The first is known as Marey's law, after a French physiologist, the other is called Bambridge's reflex after an English physiologist. Marey's law. Marey is law states that the heart rate is inversely related to the arterial blood pressure—that is, a rise or a fall in the arterial blood pressure causes, respectively, slowing or acceleration of the heart. These effects are brought about through (1) the after ent fibers of the vagus ending in the aortia and the left side of the heart, and (2) the sinus nerves. Since reflex changes in the caliber of the peripheral vessels are also brought about through these nerves any further description of their effects will be postponed until we come to deal with the control of the blood vessels (p. 199).


Banhridge's reflex. This is the name eigent to the increase in heart.

Banbridge's reflex This is the name given to the increase in heart rate which, as stated above results from a rise in venous pressure

rate which, as stated above results from a rise in venous pressure. The rise in venous pressure that occurs when the heart becomes filled acts as a stimulus to the afferent vagal endings in the right auricle Impulses are set up that depress the tone of the cardioinhibitory center. The heart rate is, in this way automatically adjusted to the quantity of venous blood flowing into its chambers. This reflex comes into effect to produce increased heart rate during muscular exercise, cardiac action being thus adjusted to pump the increased quantity of blood brought to it from the contracting muscular exercise, cardiac action being thus adjusted to pump the increased quantity of blood brought to it from the contracting muscular exercise, cardiac action being thus adjusted to pump the increased quantity of blood brought to it from the contracting muscular exercise, cardiac action being thus adjusted to the neart. The heart, like other internal organs, is insensitive to the ordinary types of stimulus. Those stimuli for example, which arouse the sensation of touch heat, cold, or pain when applied to the skin cause no sensation when applied to the heart. The heart may be touched, pinched, or cut without any sensation being aroused. In disease however (e.g., angina pectoris) severe pain may be experienced. The pain is felt over the front of the chest and down the left arm. The pain impulses travel by afferent sympathetic fibers in the middle and inferior cardiac nerves. None apparently are conveyed by the vagus. They enter the cord via the apparently are conveyed by the vagus. They enter the cord via the posterior roots of the upper four or five thoracic segments. Removal of the upper sympathetic ganglia, or the injection of alcohol around these ganglia, interrupts the pain fibers or abolishes their conductivity, and is sometimes resorted to for the relief of the severe, intrac table pain of angina pectoris

The action of certain drugs upon the heart. Atropine, as mentioned on page 188, causes an increase in the heart rate by abolishing vagal tone It acts by antagonizing the action of acetylcholine—the vagus substance (see below) Muscarine, a principle in poisonous mushrooms (amanita) slows or arrests the heart, thus, its action is similar to that following stimulation of the vagus. The effects of atropine and muscarine are, therefore, antagonistic Pilocarpine, physo-

Fig 4 51 The perfusion apparatus A is furnished with an overflow tube B the height of which can be varied to allow alterations in the perfusion pressure Fluid is delivered through the tube C The fluid from A passes through the tube D to the Inflow limb E of the double cannula which supplies the donor heart. Fafter irrigating the inside of the donor heart, the fluid passes by the outflow limb G to the glass cannulated tube H to which is attached the recipient heart 1 This cannulated tube is provided with a lateral overflow so that the hydrostatic pressure of the fluid supplied to the recipient heart remains constant J and k are the levers to which the hearts F and I are respectively attached (After Bain)



stigmine (also called eserine), choline, and its ester, acetylcholine, have the same effect upon the heart as that of muscarine Necotine acts upon the ganglion (vagal) cells in the heart first stimulating and later paralyzing them, slowing of the heart followed by acceleration results Digitalis depresses conduction in the auriculoventricular bundle, thus causing ventricular slowing. It also exerts a beneficial effect upon the heart muscle, and is probably the most valuable drug which the physician possesses for the treatment of a failing heart Excessively large doses of digitalis may cause complete functional dissociation of the auricles and ventricles—heart block

The chemical (humoral) transmission of vagus and accelerator effects. About 35 years ago, the astonishing and important discovery was made by Loewi, a German pharmacologist, that the vagus and

accelerator nerves exert their actions upon the heart, not through the direct effect of nerve impulses, but through the intermediary of chemical substances. At the terminals of the vagus in the heart, acetylcholine is produced, and the cardiac slowing or arrest, which results from vagal stimulation, is the result of the direct action of this substance upon the heart muscle The production of this vagal substance or Vagusstoff, as Loewi called it, was demonstrated by the following experiment Two frog hearts were set up as shown in Figure 4 51. The perfusion fluid, after passing through the heart on the right (donor heart), then passed through the heart on the left (recipient heart) Now, when the vagus of the donor heart was stimulated for a time, inhibition of the recipient heart followed The latter effect must have been due to a chemical substance passing into the fluid, perfusing the donor heart during the period of vagal stimulation, and then being conveyed to the recipient heart. Subsequent experiments have left little doubt that the chemical mediator of the vagal effects is an ester of choline, namely, acetylcholme. Experiments of a similar nature have proved that adrenaline (Chap. 10) is produced during stimulation of the accelerator nerves.

The vagal substance is quickly destroyed after its production. In order that a chemical shall serve as a mediator of vagal effects, it is absolutely necessary that it be removed rapidly after it is produced. Otherwise it would continue to exert its effect and so cause prolonged slowing or even arrest of the heart. The destruction of acetylcholine is brought about by the action of an enzyme present in the heart muscle as well as in the blood and other tissues. The enzyme is called cholmesterase The action of physostigmine (eserine) upon the heart s., as mentioned above, similar to that of the vagus, or of acetylcholine itself. It has been shown that eserine acts simply by antagonizing the action of the cholinesterase. In other words, it prevents the destruction of acetylcholine which is being liberated constantly from the vagal endings and, as a consequence, enhances the tonic action of the vagus.

More recent work has shown that the conception of a chemical or humoral mechanism in the transmission of nervous effects has a much wider application Not only the effects of the vague and sympathetic upon the heart, but also those of other parasympathetic and sympathetic nerves (e.g., of the stomach, intestine, blood vessels, etc.) are mediated by chemical materials (see p. 566).

The control of the action of the heart by chemical substances carried in the blood stream. The importance of the minerals sodium, potassium, and calcium upon the action of the heart has been described (see p. 162). Adrenaline, the hormone of the medulla of the adrenal glands, acts upon the heart in a manner similar to that of sympathetic stimulation (see p. 420). The carbon dioxide and oxygen pressures in the blood exert a powerful influence upon cardiac action. Excess of carbon dioxide (e.g., in asphyxia) increases the tone of the cardio-inhibitory center, reduces the rate of impulse formation in the S-A node, and depresses conduction in the auriculoventricular bundle; the heart is therefore slowed. When the pressure of carbon dioxide is such that a marked change in blood reaction toward the acid side occurs (e.g., to pH 7.0), heart block resurts. Reduction in the carbon dioxide pressure in the blood (as may be produced by prolonged forced breathing, p. 255) causes the reverse effects—namely, decrease in the tone of the cardioinhibitory center, increased rate of impulse formation in the S-A node, and enhanced conduction in the A-V bundle.

A low oxygen pressure in the blood causes an increase in heart rate. If the oxygen lack is severe or prolonged, slowing of the beat occurs, irregularities appear, and the heart fails. The heart muscle is unable to contract for long unless the oxygen supply is adequate; it is much more susceptible to oxygen lack (anoxia) than is skeletal muscle.

The length of the muscle fiber in relation to the force of the cardiac contraction. It has been demonstrated experimentally that the well-nourished heart possesses great reserves of energy. For example, when, as in muscular exercise, the quantity of blood entering the ventricles during their period of filling (diastole) is increased several fold, the heart ejects without difficulty the greater quantity of blood during the succeeding systole. Again, when the arterial blood pressure is raised to double the normal value, the left ventricle readily overcomes the great resistance and discharges its contents as easily, apparently, as it did at the lower pressure. Starling found that the dog's heart was capable of discharging, per minute, a weight of blood nearly sixty times that of the heart itself. We have already seen that the minute volume of a large robust man, whose heart weighs about 300 grams, may amount to over 35 liters (approximately 35 kilograms).

How does the heart gain the required energy for the performance of the extra work? Now, it is well known that a skeletal muscle contracts more forcibly if it is stretched by a weight, than if it is permitted to contract unweighted. The degree of stretch to which the muscle is subjected—that is, the length of the muscle fibers before excitation determines the force of the ensuing contraction. The cardiac muscle fiber behaves in a similar manner. The greater the weight, within physiological limits which is applied to the muscle during diastole and, consequently, the longer the fiber, the more powerful will be the following systole In the case of the heart, the weight which stretches the muscle during diastole is, of course, the mass of blood entering the ventricles. Thus, automatically, the heart finds the necessary energy to perform the extra work entailed by an increased venous return In order to gain power to eject blood against a raised arterial pressure, the heart does not completely empty itself for the first few beats after the pressure has been raised. The residual blood thus in creases the diastolic volume of the heart-that is, increases the length of the muscle fibers. From then on the heart empties its chambers completely during systole

The ability of the heart to do work above the ordinary requirements of the resting state of the body is called its reserve power One might therefore in a sense, compare the cardiac muscle fiber to an elastic band which as we know, rebounds most powerfully when it is fully stretched. The behavior of the heart in this regard has been expressed by Starling in the following words "The energy liberated by a cardiac contraction is a simple function of the length of the muscle fibers of the ventricle at the end of diastole 'This is known as Starling's law of the heart

When, as a result of disease, the contractile power of the heart is reduced, it must dilate. In other words, it must increase the length of its fibers to a greater degree than does a healthy heart in order to per form the same amount of work Thickening-hypertrophy-of the muscle fiber also occurs Enlargement of the heart is therefore a sign of heart disease Nevertheless, the enlargement in itself is beneficial, and may be looked upon as a compensatory process which enables a heart which is at a mechanical disadvantage (e.g., one with a valvular lesion) to discharge its functions. Indeed the hypertrophied heart may have a reserve power almost as great as that of the normal heart When the heart muscle itself is diseased, it may be so greatly weakened that it cannot even at maximal dilatation, contract forcibly enough to maintain an efficient circulation. The signs of heart failure then ensue—breathlessness (dyspnea, p. 260), edema (p. 101), and the appearance of a bluish tint of the skin, especially of the lips, ears, and fingertips (cyanosis p 263)

It is now generally accepted that a healthy heart cannot be "strained" by overwork Athletics, even though of a strenuous nature, do not injure a normal heart. In other words, the yolume of blood brought to the heart from the contracting muscles is not great enough to tax the normal healthy heart beyond its reserve powers; the muscles will fatigue or come to the limit of their capability before the heart is injured. On the other hand, if a person with valvular, coronary, or other organic cardiac disease indulges in strenuous muscular exercise, serious injury may be inflicted upon the heart, for the work demanded of the heart may then exceed its reserve power, which has been reduced as a result of deterioration of the heart muscle caused by the existing disease.

## Nerve Control of the Blood Vessels

As already mentioned on page 124, the walls of the arterioles are composed largely of smooth muscle fibers arranged in such a fashion that when they contract the vascular lumen is narrowed. This muscle, in so far as its nerve supply is concerned, may be compared to the muscle of the heart. One set of nerve fibers—pasoconstrictors—exite

the muscle—that is, cause constriction of the arterioles. The other set—vasodilators—inhibit the muscle, causing it to relax and so cause vascular dilatation.

The vasoconstrictor nerves. The vasoconstrictor nerves belong to the sympathetic division of the involuntary nervous system. They arise, in common with other sympathetic fibers (p. 563), from cells in the lateral horns of the gray matter of the spinal cord lying between the levels of the first thoracic and the third lumbar segments. The vasoconstrictor fibers to the vessels of the limbs, for the most part, are carried in the mixed spinal nerves (e.g., ulnar, sciatic, etc.). Those to the head and neck pass from the cervical ganglia of the sympathetic to the carotid arteries which they surround in a netlike (plexiform) manner. These nerve nets or plexuses invert the vascele to this could be the sympathetic to the carotid articles the care of the sympathetic to the carotid arteries which they surround in a netlike (plexiform) manner. These nerve nets or plexuses



Fig. 4 52 Showing effect upon arterial blood pressure of stimulating the right splanchnic nerve. (After Macleod)

invest the vessels to their smallest branches, ultimately furnishing filaments to the arterioles. The vasoconstrictor fibers to the abdomen pass to the vessels in three well-defined strands—the greater, the lesser, and the least splanchnic nerves. The greater splanchnic nerve supplies the larger proportion of the arterioles of the abdomen; when stimu106

lated, the wide-spread vasoconstriction which results causes a very pronounced rise in blood pressure (Fig. 4.52).

The vasodilator nerves. We may group the vasodilator nerves into

The vasodilator nerves. We may group the vasodilator nerves into three categories: (a) those which belong to the sympathetic division of the sympathetic nervous system, (b) those belonging to the parasympathetic division, and (c) those that leave the central nervous system in the posterior spinal nerve roots. The sympathetic vasodilators arise from the thoracic and lumbar segments of the spinal cord and follow the same general course as the vasconstrictors. They appear to be distributed almost exclusively to the vessels of the muscles and viscera.

Of the parasympathetic vasodilators, some leave the brain in the facial, glossopharyngeal, and vagus nerves; others arise in the sacral part of the cord and are carried in the pelvic nerve (p. 563) to the vessels of the bladder, rectum, and external genital organs. The vasodilators which leave the brain in the facial nerve enter its chorda tympani branch, and are ultimately distributed to the vessels of the tongue and salivary glands.

Vasodilator impulses conveyed along the pelvic nerve to the vessels of the penis or of the corresponding organ of the female—the clitoris—are responsible for the phenomenon of erection (Ch. 14). The vasodilator fibers of the posterior spinal nerve roots appear to be distributed, mainly at any rate, to the skin of the limbs. There has

The vasodilator fibers of the posterior spinal nerve roots appear to be distributed, mainly at any rate, to the skin of the limbs. There has been much discussion as to the nature of these nerves. When the posterior nerve roots supplying fibers to a limb are stimulated, the vessels may dilate. Many believe that the ordinary sensory fibers, of which the posterior roots are mainly if not entirely composed, must transmit the vasodilator impulses. That is to say, impulses pass to the vessels, apparently, over fibers which have been thought to be capable only of conveying impulses to the spinal cord—ie, sensory impulses. According to this conception, the vasodilator impulses travel over the nerve in a direction opposite to that of the ordinary sensory impulses. For this reason, they have been termed antidromic (Gk. anti, against + dromos, a running) vasodilator impulses. Other physiologists offer another explanation of the vasodilatorin which follows stimulation of the posterior roots, claiming that efferent fibers are contained in the posterior roots, and that these alone are the transmitters of vasodilator impulses. The question remains unsettled.

The vasoconstrictor and vasodilator centers. The vasoconstrictor and vasodilator centers are situated in the medulla oblongata, in the floor of the fourth ventricle (see p. 559). The vasoconstrictor center is

connected by tracts of fibers with the nerve cells in the lateral horns of the spinal cord which, as mentioned above, give rise to the vaso-constrictor fibers. These cells may be loked upon as constituting a spinal vasoconstrictor center. Centers controlling the vessels also probably exist at higher levels than the medulla—for example, in the hypothalamus and even in the cerebral cortex.

The vasoconstrictor center in the medulla and, though to a much less degree, the vasodilator center as well, possess tone. That is to say,

Fig. 4 53 Showing the vessels of rabbit's ear. The sympathetic (vasoconstrictor) nerves supplying the right ear have been cut. The nerves to the left ear are intact. Note the dilatation of the vessels of the right ear as compared with those of the left. The difference in caliber of the vessels on the two sides would be still more pronounced if the sympathetic nerves of the left ear had been stimulated.



impulses are constantly passing from the centers along the vasoconstrictor and vasodilator nerves. Under ordinary circumstances, vasoconstrictor tone predominates—the vessels being always slightly constricted. This tonic vasoconstrictor action is at once abolished if the spinal cord is cut across in the lower cervical region—the stream of impulses from the medullary center to the minute blood vessels being thus interrupted. The blood vessels then dilate, the peripheral resistance, which is such an important factor in maintaining the arterial blood pressure, is greatly reduced, and a profound fall in pressure results. The low blood pressure persists for a time, which varies with the animal species, being longer in mammals than in lower forms

and then gradually feturns to normal. The restoration of the blood pressure to the normal level is due to the centers in the spinal cord assuming the functions of the isolated centers in the brain Sectioning the splanchnic nerves by depriving the abdominal vessels of tonic vasoconstrictor impulses, acts in a similar way to cause a fall in blood pressure.

That dilatation of vessels occurs when the vasoconstrictor fibers to a part are cut can be very simply demonstrated upon a rabbit The vessels of the rabbit s ear are plantly visible when the ear is held up in front of a bright light. If the sympathetic nerve in the neck (cervical sympathetic) is cut while the ear is being observed in this way, the ear becomes flushed and warm and the individual vessels are seen to dilate see Figure 4.53. The effect is still more striking if the opposite ear (with nerve uncut) is compared with the ear on the operated side. This experiment first performed by Claude Bernard the great French physiologist of the nineteenth century, is taken to mean that a constant stream of (tonic) vasoconstrictor impulses is carried along the intact nerve which thus maintains the vessels in a state of partial constriction. These impulses are interrupted when the nerve is cut.

painful sensation is aroused, reflex vasoconstriction is more likely to result.

The aortic and sinus nerves. There are two sets of afferent fibers which exert a very special controlling influence upon the circulation. When stimulated, they cause reflex slowing of the heart together with vasodilatation, especially of the abdominal (splanchnie) region of the vascular system. As a consequence of these effects, the blood pressure falls. One set of such afferent fibers is contained in the vagus nerve. They terminate peripherally in the wall of the aorta and base of the heart. An increase in the blood pressure, by stretching the aorta, stimulates the nerve terminals embedded in the arterial walls; impulses

ascend to the cardiac and vasomotor centers in the medulla; impulses are discharged, in turn, down the efferent fibers of the vagus and the vasodilator nerves. The tonic vasoconstrictor impulses are, at the same time, reduced in frequency. The vasodilation is therefore due to reduced vasoconstrictor tone as well as to increased tone of the vasodilators. Although, in man, these afferent (depressor) fibers are bound up in the vagus trunk with the efferent fibers, in certain animals (e.g., the rabbit) they are contained in a separate branch of the vagus, known as the cardiac depressor or aortic nerve, shown in Figure 4.54.

The other set of afferent fibers are contained in the sinus nerve, a small branch of the glossopharyngeal nerve. The filaments of this nerve terminate in receptors situated in the wall of the carotid sinus, the name given to a slight enlargement of the common carotid artery where it divides into the internal and external carotids (Fig. 4.55). These receptors, since




Fig 4.54 Showing the carotid sinus and arch of aorta with the sinus and aortic nerves. I, arch of aorta; 2, common carotid; 3, carotid sinus, 4, internal carotid, 5, external carotid; 6, vagus nerve; 7, glossopharyngeal nerve; 8, sinus nerve, branch of the glossopharyngeal nerve.

they respond to a pressure or stretch stimulus, are called pressoreceptors. When stimulated in this way, as by a rise in blood pressure, nerve impulses are transmitted along the sinus nerve to the cardiac and vasomotor centers in the medulla oblongata. Reflex slowing of the heart, vasodilatation, and a fall in blood pressure follow. Mere compression of the sinus between the finger and thumb in an animal, or even pressure through the skin of the neck in a proportion of normal human subjects, will stimulate the nerve endings in the sinus wall and evoke the reflex. In certain rare instances, the human carotid sinus is abnormally sensitive. In such persons the lightest pressure in the neighbor-




Fig 4.55 Showing the carotid sinus region in man 1, common carotid artery; 2, carotid sinus, 3, internal carotid, 4, external carotid, 5, sinus nerve; 6, glossopharyngeal nerve

hood of the sinus (as during shaving, or buttoning a collar) may cause pronounced slowing of the heart, and a sharp fall in blood pressure. The fall in blood pressure may, by depriving the brain of an adequate supply of blood, cause syncope (fainting) or convulsions.

The maintenance of the arterial blood pressure within the normal limits is dependent to a large extent upon the activity of the sinus and aortic nerves. These nerves transmit at all times a stream of impulses to the cardiac and vasomotor centers, and thus exert continuously a depressing effect upon the blood pressure. Any marked rise in blood pressure increases the frequency of the afferent


impulses and magnifies the constant depressor effect upon the medulary centers; the blood pressure is thus automatically reduced again. In a corresponding manner, any tendency for the blood pressure to fall is countered by an opposite effect, namely, reduction of the frequency of the impulses going to the center and, in consequence, reflex cardiac acceleration and vasoconstriction (Fig. 4.56). The sinus and aortic nerves thus exert a governing or buffering effect upon the blood pressure, preventing it from fluctuating greatly in either direction. For this reason they are sometimes referred to as the "buffer nerves" of the circulation.

In hemorrhage, for example, the fall in blood pressure occasioned by the reduction in blood volume, diminishes the intensity of the pressure stimulus normally applied to the sinus and aortic nerves, thus lowering the frequency of the afferent impulses impinging upon the centers. Reflex vasoconstriction and cardiac acceleration follow, which tend to restore the blood pressure to its normal level. The vasoconstriction seen in surgical shock, and the ability of the circulation to

compensate for the effect of gravity and maintain the blood pressure when the erect posture is assumed, are mainly dependent upon the activity of the sinus and acrtic nerves.

Filaments of the sinus nerve also end in the carotid body, a collection of specialized cells in the neighborhood of the carotid sinus; but since this body is concerned with the control of respiration rather than with the circulation, its structure and function will be deferred.

Fig 4.56 Record A, upper tracing, blood pressure. Lower, record of impulses from a single isolated fiber of the sinus nerve Record B, upper tracing, blood pressure at a higher level than in A. note that the impulses recorded in lower tracing from the sinus nerve are at a higher frequency and persist even throughout the diastolic phase



A similar structure, known as the aortic body, is to be found on a small artery arising from the arch of the aorta; its receptors are innervated by terminals of the aortic nerve, and stimulated by a low oxygen tension of the arterial blood. When so stimulated, impulses travel along the aortic nerve, and reflex vasoconstrictor results. Thus, while the carotid body is concerned, chiefly at any rate, with the regulation of respiration, the aortic body is part of the mechanism controlling the circulation.

The reflex control of the heart and blood vessels is shown diagrammatically in Plate 5B.

## Special Features of the Circulation in Certain Parts

The capillary circulation. The capillaries are the smallest vessels of the circulatory system, measuring about 1 mm in length, and having an average diameter of 8 to 15 microns. Their walls are composed of a single layer of endothelial cells cemented together at their edges, and are of such delicate construction that they are seen with the

greatest difficulty, even under careful microscopic examination. The blood within the capillaries (the corpuscies being clearly seen through their walls) enables one to locate the position of these vessels. In amphibians, here and there, peculiar spiderlike cells are to be seen in close relation to the capillary. The bodies of these cells possess several long slender processes which embrace the endothelial wall. They are called Rouget cells (see Fig. 4.10, p 125).

Though, of course, the existence of small vessels between the arternal and venous systems was implicit in Harvey's theory of the circulation of the blood, it was not until some years after Harvey pro-




Fig 4.57 Vessels in a capillary area. (After Chambers and Zweifach, modified.)

until some years after Harvey propounded his theory that the capillaries were actually demonstrated by the Italian anatomist, Malpighi. It has been mentioned elsewhere (p. 125) that the whole purpose of the elaborate circulatory system is to drive blood through the capillaries, for it is through the walls of these vessels alone that oxygen and nutritive materials pass from the blood to the cells of the tissues, and carbon dioxide and waste materials from the tissue to the blood.

Each arteriole is continued into a vessel of about the size of a capillary but differing from the latter in that its walls contain a small quantity of smooth muscle; it is called a metarteriole (Fig. 4 57). Such vessels run centrally through a capillary area. This central or thoroughlare channel gives off a number of true capillaries—that is, simple endothelial tubes which, after forming a network of loops, return the blood to the central

channel nearer its venous end. The metarteriole, by virtue of its smooth muscle and of its possessing a vasoconstrictor and vasodilator nerve supply, is capable of altering its caliber and thus of decreasing or increasing the quantity of blood flowing through the capillary area which it serves The capillaries themselves, in opposition to older accounts given of these vessels, are not able actively to alter their calibers. They

are merely dilated passively by an increased volume of blood entering them when the metarterioles dilate. Changes in the blood flow within the capillaries occur frequently in accordance with the state of activity of the surrounding tissues (Fig. 4 58). In a resting muscle, for example, a large proportion of the capillaries are closed, the channels being completely obliterated. When the muscle contracts, large num-



capillaries of guinea pig injected during life with India ink, showing different degrees of constriction. Blank oval and circular areas indicate red corpuscles. (After Krogh)




Fig. 4 59 Showing capillary loops in the human skin at the base of the fingernail, as seen under low power of the microscope. The divisions in the scale at the bottom of the illustration represent tenths of millimeters. (After Lewis )

bers are opened up, thus increasing the blood flow through the tissue several fold, and assuring an adequate oxygen supply for the contraction process.

A number of chemical materials exert an influence upon the diameter of the metarterioles and so, indirectly, upon the caliber of the capillaries. Carbon dioxide and lactic acid, produced during tissue activity, cause dilatation of the metarterioles. Iodine, urethane, silver nitrate, and histamine have a similar effect. Adrenaline and pituitrin cause constriction.

The capillaries of the human skin. The most superficially placed capillaries of the human skin are in the form of hairpinlike loops situated in the papillae of the dermis, and running more or less at right angles to the skin surface with their convexities directed outward. They lie just beneath the papillae of the skin. These loops may

be seer when the skin at the base of the fingernail is examined with a strong light beneath the microscope (PI 6A and Fig 459) One limb of a capillary loop receives blood from the arterial side, the blood in the other limb, having given up a part of its oxygen load, is more venous in character. The venous limb drains into a plexus of capil. lary vessels lying beneath the bases of the papillae of the skin, and named the subpapillary venous plexus. It is the blood in this plexus, and not that in the capillary loops, which gives color to the skin. When the flow is rapid and the vessels of the venous plexus dilate, the skin is bright red in color When the vessels are constricted, the skin is pale. If the blood flow through the plexus becomes slowed, more oxygen is given up by the blood within the vessels, a greater proportion of the hemoglobin is in the reduced state, and, in consequence, the skin acquires a bluish tint (see cyanosis, p 263) This is seen most commonly in subjects with heart disease, the slowing then being due to failure of the heart pump But slowing of the cutaneous blood flow, to a less degree, may occur in normal persons if exposed to cold When the body is chilled, the arterioles of the skin become constricted (an attempt on the part of the body to conserve heat, p 376), as a result of the diversion of blood to deeper parts, the blood flow through the capillaries of the subpapillary venous plexus may be slowed to such an extent that a definitely blue tint is given to the skin, especially of the finger tips, ears, and nose (See also pp 376-7)

The blood supply to the heart muscle, or coronary circulation. The heart of mammals possesses a well developed system of vessels—arteries, arterioles, capillaries, and veins The blood is delivered to the heart muscle by the right and left coronary arteries (Pl 7A) and, after passing through the small vessels, is drained into the right auricle by a large vein known as the coronary sinus (which receives blood from the great, middle, and small cardiac veins) and by the two anterior cardiac veins A number of smaller and more deeply situated veins known as the veins of Thebesius, after an eighteenth-century anatomist who first described them, empty directly into the auricular and ventricular cavities About 60 per cent of the blood escapes from the coronary system by the coronary sinus, the remainder by the anterior cardiac veins and the veins of Thebesius The coronary arteries arise from the aorta just beyond the attachments of the aortic valves

Unlike any other organ or tissue of the body, the greatest flow of blood through the heart muscle takes place during the diastolic phase of the heart During the ejection phase of ventricular systole, the

coronary blood flow is greatly reduced, for the cardiac muscle fibers when they contract compress the coronary vessels; the flow of blood is then reduced through some vessels and completely arrested through others.

The mean blood pressure during the diastolic phase of the cardiac cycle is, therefore, the important factor in filling the coronary vessels and determining the magnitude of the coronary blood flow.

The coronary vessels receive fibers from both the vagus and the sympathetic nerves. The vagus causes vasoconstriction; the sympathetic nerves cause vasodilatation Adrenaline also causes dilatation of the coronary vessels, in animal experiments, a threefold increase in coronary blood flow has been obtained by the use of this substance.

Angina pectoris and coronary occlusion. The term angina pectoris means breast pain, and is used to denote an attack of pain, usually severe, felt beneath the breast bone (sternum), and in many cases in the left shoulder and arm. It is due to the heart muscle being inadequately supplied with oxygen. The commonest cause is disease (arteriosclerosis) of the coronary vessels. The disease causes narrowing of the caliber of the arteries and, as a consequence, reduces the blood supply to the heart muscle. The attack most frequently follows some muscular effort or emotional excitement which increases the cardiac output—that is, the work of the heart. In other words, the heart is called upon to do work for the performance of which it is unable to obtain an adequate supply of oxygen. A drug such as amylinitrite, or nitroglycerine, both of which are dilators of the coronary arteries, is commonly given to arrest an attack.

The severe pain associated with angina pectoris is believed to be due to the action of a chemical upon the nerve endings. This substance, when the oxygen supply to the heart is adequate, as in health, is removed or destroyed about as quickly as it is formed, but when the supply of oxygen is deficient its accumulation within the heart

muscle gives rise to anginal pain.

Complete occlusion of a large branch of a coronary artery is not an uncommon event. The occlusion is usually the result of the clotting of blood within a vessel (thrombosis) that is the seat of arteriosclerosis. Coronary thrombosis, or occlusion of a large coronary branch from whatever cause, is accompanied by intense pain, nausea and vomiting, breathlessness, and collapse. Death may occur almost instantly. If recovery takes place, the area of heart muscle deprived of its blood supply dies (necrosis), and is replaced by scar tissue. Rupture of the diseased area may, however, occur before healing

has taken place. The electrocardiogram taken shortly after an attack shows certain special features which are of the greatest value in diag

The pulmonary circulation. The blood passes from the right to the left side of the heart through the lungs. The same quantity of blood as a discharged in a given time by the right side of the heart is received and discharged by the left. In other words, the quantity of blood passing through the lungs per minute is the same as that passing through the systemic circulation. The vessels of the pulmonary circuit, as of any other part, consist of arteries, arterioles, capillaries, and veins. The main arterial vessel is called the pulmonary artery it arises from the right ventricle and carries venous blood—that is, blood in which the hemoglobin is in the reduced state. The main veins are four in number (two from each lung), they contain arterial (oxygenated) blood, which they deliver into the left auricle.

The capillaries of the lung differ from those of any other tissue in that they are separated from air only by the very delicate and incomplete membrane composing the walls of the pulmonary alveoli. The pulmonary capillaries can increase their capacity to a relatively great extent Ordinarily the lungs contain from 6 to 10 percent of the total amount of blood in the body, but under certain conditions they may contain 20 percent of the total blood volume. If, for example, there is some resistance on the left side of the heart (e.g., mitral stenosis or weakness of the left ventricle) while the right ventricle continues to contract forcibly, the pressure in the pulmonary circuit rises, the capillary bed becomes distended, and a much greater volume of blood is contained in the lungs. This is likely to cause breathlessness—a common symptom of heart disease—because the dilated vessels encroach upon the pulmonary alveoli (p. 230). The greater amount of blood in the lungs are "stiffer" and do not distend and deflate during respiration with their normal residency. This tends to impair the exchange of the respiratory gases between the lungs and the atmosphere.

The pressure in the pulmonary circulation is about one sixth of that in the systemic circulation. In the dog, the mean pressure in the pulmonary artery is around 25 mm Hg as compared with 130 to 150 in the aorta.

Variations in intrathoracic pressure occur during ordinary quiet

<sup>†</sup> The blood carried to the lungs by the pulmonary artery does not serve to
supply the pulmonary tissue useful with oxygen and nutriment This function devolves
upon the systemic circulation. The bronchial tree and pulmonary tissue are supplied
by branches of the aorts.

respiration. The intrathoracic pressure amounts to about -6 mm Hg during inspiration, and about -2.5 mm Hg during expiration. More blood, therefore, is "sucked" into the great veins of the thorax and right auricle during inspiration than during expiration; more blood is discharged from the right heart into the pulmonary system during inspiration. The quantity of blood received and given out by the left side of the heart must, of course, vary with the quantity discharged by the right side. The variations in output of the left ventricle cause




Fig 4.60 Illustrating the effect of the respirations upon the blood pressure.

corresponding fluctuations in the systemic blood pressure. The increased discharge of the left ventricle and the rise in systemic blood pressure are not, however, synchronous with inspiration, but owing to the increased vascular capacity of the lungs themselves, caused by the inspiratory movement, are somewhat delayed. The rise in systemic blood pressure caused by inspiration, therefore, does not commence until near the end of this movement, and reaches its maximum toward the end of expiration (see Fig. 4.60).

The pulmonary vessels are supplied with fibers from the sympathetic and the vagus. The former are mainly vasoconstrictor in action, the latter mainly vasodilator.

The hepatic circulation. The blood from the gastrointestinal tract and spleen is collected by the portal vein and conveyed to the liver. The blood from the small intestine contains materials derived from the food—that is, glucose, amino acids, as well as a proportion of the fat. The portal vein breaks up within the liver into numerous branches; its smallest subdivisions run between the lobules of the liver,

208

and are called interlobular veins. These open into capillarylike (but relatively wide) vessels called sinusoids, lying between rows of liver cells (the liver cords). All the sinusoids of a liver lobule converge toward the center of the lobule where they drain into a central vein. The central veins drain in turn into a system of larger (sublobular) veins which ultimately join to form vessels called the hepatic veins;




Fig 4 61 The hepatic circulation, see also Figure 8 27, page 329.

these open into the inferior vena cava (see Figs. 4.61 and 8.28, p. 330). Thus the portal vein resembles an artery in that it breaks up into branches which lead into capillary vessels. It, therefore, lies between two capillary beds, one in the gastrointestinal tract and spleen which it drains, the other in the liver to which it is distributed. The blood of the portal vein has a pressure of only from 10 to 12 mm Hg; it has a relatively low content of oxygen and a relatively high content of carbon dioxide. But the liver cells must receive well-oxygenated blood under high pressure. This is delivered through the hepatic artery -a branch of the celiac artery. The latter vessel arises directly from the abdominal aorta. The blood of the hepatic artery is delivered into the sinusoids of the liver where it mixes with the blood of the portal vein (see Fig. 4.61). Thus, whereas the portal blood carries blood loaded with digestive products which are subjected to various changes by the liver cells, the latter cannot survive if they receive only the small quantity of oxygen which this blood can supply. The function of supplying the hepatic tissue with adequate quantities of oxygen devolves upon the blood of the hepatic artery.

The cerebral circulation. Blood is delivered to the brain through two pairs of arteries—the internal carotids and the vertebrals—and is drained away by the internal jugular veins. The vertebral artery is a branch of the subclavian artery. It must be remembered that the brain is enclosed in a rigid case—the skull. Since the contents of the skull—brain, blood, and cerebrospinal fluid—are incompressible, their volume can alter very little from time to time. More blood can be accommodated in the cranial cavity only through the displacement of an equal quantity of cerebrospinal fluid into the spinal canal, where room is made for it by the stretching of membranes situated between the joints of the vertebrae. Though the quantity of blood within the skull changes comparatively little from time to time, the velocity of the blood flow through the cerebral vessels varies considerably. The flow is determined largely by the blood pressure in the rest of the body, a rise in pressure in the aorta increasing the cerebral blood flow, and vice versa. For example, when a person assumes the standing position, the carotid sinus and aortic nerve mechanisms come into play and prevent a fall in the pressure of blood in the arteries going to the head, which otherwise would result from the effect of gravity upon the circulation. The brain is thus assured of an adequate blood supply.

The flow through the brain is dependent also upon the pressure gradient—that is, upon the difference between the pressures in the cerebral arteries and the cerebral veins. An increase in the pressure in the internal jugular veins (the pressure in the cerebral arteries being unchanged) will therefore tend to duminish the cerebral blood flow.

Variations in the blood flow through the cerebral vessels can occur, however, quite independently of the general blood pressure for, though denied for years it is now accepted that these vessels are supplied with nerves, which can bring about active changes in vascular calibers. Though the systemic blood pressure remains unchanged, an increased cerebral blood flood can therefore be effected through intracranial vasodilatation. Constriction of the cerebral vessels, with a constant systemic pressure, will cause the opposite effect—namely, a reduction in the blood supply to the brain. The vasodilator fibers to the cerebral vessels are derived from the facial nerve; the vasoconstrictors, from the sympathetic.

## The Defensive Mechanisms of the Body against Disease and Injury; Shock, Anaphylaxis, Allergy

Infecting agents Man animals, and plants are constantly liable to attack by many lower forms of life—namely, bacteria fungi or molds (which cause ringworm and other affections of the skin), and protozoars. These latter are the infecting organisms in some of the most virulent of human diseases such as malaria amebic dysentery, African sleeping sickness etc. Bacteria and fungi belong to lower orders of the vegetable kingdom. Protozoa are unicellular animal forms (Fig. 5.1)

The viruses, another large group of infecting agents, are the cause of a number of well known diseases such as measles mumps chick-enpox, smallpox influenza one type of pneumonia anterior polio myelitis (infantile paralysis), rabies, psitacosis (a disease of parrots and allied species transmissible to man), and the common cold They are also responsible for several diseases of plants and even attack bacteria. It is the general opinion that viruses are living organisms,

<sup>&</sup>lt;sup>1</sup> Viruses that infect bacteria are called bacteriophages. There are several varieties each of which attacks a specific type of bacterium—e.g. staphylococci colon bacilli etc.

or relatively large molecules of nucleoprotein (p 366) on the border line between the living and the nonliving. They are of minute size, as compared with bacteria, most are invisible under the ordinary microscope and pass readily through a fine filter (filtrable viruses). Unlike bacteria, they cannot be grown apart from living cells. The intensely infective nature of these agents is evident from the fact that, in the case of one at least, the particular disease of which it is the cause can be transmitted by as small an amount as 1 cc of a solution containing it in a dilution of only 1 part in 10 billion.

It had been suggested by various medical scientists from time to time throughout the centuries even in Roman days, that disease was caused by minute invisible particles carried in the air But it remained for the French chemist, Louis Pasteur (1822-95) to demon strate that many diseases were caused by living organisms which we now know as bacteria Joseph Lister (later Lord Lister, 1827-1912), an English surgeon, brought Pasteur's discoveries to bear upon the practice of surgery. He showed that if strict antiseptic methods were applied, wounds would remain surgically clean—that is, free from infection. The work of these pioneers has made possible the great ad vances in surgery that have marked the last 90 years or so

But even before this, Ignaz Semmelweis (1818-65) of Vienna and Oliver Wendell Holmes in America showed clearly to unprejudiced minds, (of which there were few at that time) that puerperal fever (blood poisoning following childbirth) was caused by an invisible something carried to the patient on the hands and instruments of the attending physician, and that death from this disease, which in hospitals sometimes reached almost the proportions of a plague, could be greatly reduced by the use of antiseptics. See Figures 5.2, 5.3, 5.4, and 5.5.

Bacterial forms. The types of bacteria are innumerable, but fortunately only a small proportion cause disease. Also, only a relatively few kinds of fungi and Protozoa threaten man or animals. Several bacterial forms are actually beneficial to man and some are even essential for human welfare. Those, for example, which inhabit the intestinal tract effect the final breakdown of food material with the formation of gas which, together with the bulk by which the bacteria

This calls to mind Dean Swift's amusing little rhyme which was written at least 100 years before bacteria were discovered

"So naturalists observe a flea Hath smaller fleas that on him prey" And these have smaller still to bite 'em, And so proceed ad Infinitum."




Fig 5 1 Some pathogenic microorganisms

themselves add to the feces, stimulate the bowel movements Othe bacterial types present in the intestine are capable of synthesizing several factors of the vitamin B complex, as well as vitamins C and K  $^{\rm 2}$ 

<sup>2</sup> It sometimes happens that a pat ent receiving an antibacterial agent ("sulfa" drug or antibiotic p 221) especially when given by mouth develops a deficiency in certain vitamin or vitamins as a result of the destruction of intestinal bacteria

From others again, and from molds, anti-infective substances of var-'ious kinds have been isolated (see p. 221).

The commoner types of disease-producing (pathogenic) bacteria can be classified, on basis of their microscopical appearances, into four main groups, cocci (sing. coccus), bacilli (sing. bacillus), spirilla (sing. spirillum), and spirochetes (see Fig. 5.1).




Fig 5 2 Louis Pasteur.

Fig. 53 Joseph Lister.

The cocci are dotlike bodies (Gk. kokkos, a berry) which appear singly, in parts (Diplococci, a species of which causes lobar pneumonia), in clumps like grapes (Staphylococci), or in chains (Streptococci). These last two types are the cause of acute pyogenic inflammatory conditions. One strain of streptococcus is the cause of scarlet fever.

The bacilli are rod-shaped organisms; examples of which are the typhoid, tetanus, and diphtheria bacilli and the tubercle bacillus.

Spirilla are slender organisms possessing a shape like the shaft of a corkscrew.

The spirochetes, one genus of which is the cause of syphilis, have a slender spiral structure.

The reactions of the tissues to infecting agents. A pathogenic organism or virus shows, as a rule, a highly selective action with respect

to the tissue which it or its town injures, and the symptoms characteristic of the disease which it causes are the manifestations or the results of such injury, or represent the reaction of the tissue cells to invasion by the particular type of microorganism or virus. The town of the tetanus bacillus, for example, singles out nervous structures lowering the "resistance" at synaptic junctions, so that a relatively







Fig 5 5 Oliver Wendell Holmes

mild stimulus—a mere touch a faint sound or a light—may cause widespread muscular responses. The viruses of rabies and anterior poliomyelitis also direct their attack to the central nervous system the former injuring nerve centers in the brain, the latter destroying chiefly the motor cells of the spinal cord. The typhoid bacillus cause ulceration of the mucous membrane of the small intestine, the tubercle bacillus has a predilection chiefly for pulmonary, bone, and joint tissues (though almost any tissue may be attacked). The bac terra of general septic infections the staphylococci and streptococci, show much less selectivity of action, for they may cause an inflamma tory condition in any tissue (see also p. 92.). The spirochete of syph ilis is also devoid of selective action, skin, bone, nervous or any other tissue may be attacked. The reactions of the tissues that it infects tristue may be attacked. The reactions of the tissues that it infects are, however as are also those of the tubercle bacillus quite characteristic. The infected tissues react to both of these types by attracting small round cells, mostly lymphocytes, which form masses around

the invading organism. Thus a tubercle is formed in the case of tuber culosis, or a tumorlike structure called a gumma if spirochetes of syph ilis are the invading organisms

The processes of immunity The body is capable of putting up a defense which is, in part or entirely, successful against the inroads of most of the bacteria and viruses. The protection when it develops fully is called immunity. One who has suffered from an attack of small-

tully is called immunity. One who has suffered from an attack of small-pox for example, cannot (except, perhaps, in the rarest instance) contract the disease again. With respect to that particular disease he is immune. Vaccination also confers immunity against smallpox. The defenses of the body against the various disease producing (pathogenic) organisms depend upon the activity of living cells. Cells, such as the neutrophil polymorphonuclear leucocytes and certain wan dering cells of the reticule endothelial system exert their protective action in an obvious, overt or, one might say, in a mechanical way, they ingest the bacteria. The phagocytic property of the neutrophil leucocytes has been already described in Chapter 3. The immunity which the leucocytes confer is not specific, for they attack various types of bacteria indiscriminately though they are particularly active against the pus producing (pyogenic) forms. Other cells of the body, especially the lymphocytes in the lymphoid tissue generally, throughout the body, produce and liberate chemical substances closely associated with the globulin of the plasma, which have a specific neutralizing or destructive action upon the agent causing the disease. These izing or destructive action upon the agent causing the disease. These substances which go under the generic term of antibodies are as it substances which go under the generic term of antibodies are as it were, made to order, each for protection against only one pathogenic agent. Thus the antibody which is responsible for immunity against smallpox is mactive against measles or any other infectious disease. There are several types of antibody—namely, antitoxins—which combine with and neutralize the bacterial poison, a substance generally called a toxin Others called cytolysins or bacteriolysins destroy the foreign cell itself, to this group belong the specific hemolysins (p. 84) which cause the destruction of the red cells of one species when injected into the blood of another Agglutinis with which we are familiar (they were referred to in connection with the blood groups, p. 78) render bacteria innocuous by clumping them together in a mass. Agglutinis are produced in the blood of a patient suffering from typhoid fever. A test for the diagnosis of typhoid fever is based upon this fact, the serum of a patient with typhoid fever alone is capable of agglutinating a suspension of typhoid bacilli. Other antibodies,

known as precipitins, form an insoluble compound with the bacterial toxin and with foreign proteins \*

The foreign agent—whether a red ceil in incompatible blood, a tissue ceil of another species, a bacterial ceil, or a virus, which stimu lates the tissue cells of the host (i e, of the infected person or animal) to produce the antibody—is called an antigen (anti [body] + Gk root gen, to form or produce) The response of the tissues of the host to the antigen and the production of the antibody, together with the reaction between the two, is called the antibody antigen reaction. Paul



Fig 56 Diagram illustrating the production of antitoxin according to Ehrich's side chain theory Q = toxin black toxphore group blank circle haptophore group bly = receptors or hapture of the toxin and through on the cell Under the stimulus of bon the cell Under the stimulus of

which the latter exerts its harmful effect upon the cell. Under the stimulus of the toxin (antigen) receptors are produced in excess and escaping into the general blood stream constitute the antitoxin


Ehrlich of Berlin (1854-1915) many years ago sought to explain this reaction and the immunity which it conferred by what has come to be known as Ehrlich's side chain theory of immunity (see Figs 56 and 5 7) Thus, in the case of immunity due to an antitoxin, he conceived that the protoplasm of the tissue cells, stimulated by the antigen, formed and put forth active chemical groups, or side-chains, called receptors or haptins these combined with the bacterial toxin, and, being thus linked to the tissue cells, brought about the injury characteristic of the disease. But the tissue cells produced an excess of such receptors which, becoming detached passed into the general circulation where they combined with toxin before it could reach and damage the cells. These free receptors constituted the antitoxin. The same conception could be made to embrace other types of antibody Ehrlich suggested that the toxin molecule possessed two active chemical groups, one, known as the toxophore group, was responsible for its injurious action. The other, called the haptophore group, united

A delicate and specific test for human blood in medico-legal (murder) cases is based on the precipitin reaction A rabbit is given a series of injections of a solution of human blood or serior When the serior of the rabbit which as a result of the injections has developed a precipitin active against human serior is added to a specine of the supported blood (e.g. from stans on clothing, a presumed weapon etc.) dissolved in a test tube a slight cloudy precipitate appears if the material in question is human blood. If it is the blood of an animal the solution remainst clear.

with the cell receptor or haptin, and thus enabled the toxin to bring its action to bear upon the tissue cell.

Three types of immunity are recognized: natural, active, and passive. Natural immunity, also called inborn or innate, is that which has not been acquired by a previous attack of the disease, or in any other way. Thus, we are naturally immune to some diseases which afflict animals, and they possess a natural immunity to a great many human

Fig. 57 Diagram illustrating the production and action of specific cytolsyns, hemolysins, bacteriolysins, etc. According to Ehrlich's theory a substance destroyed by heat and called the complement (\psi\) is present in nearly all normal sera. It be-



comes joined to the receptor ( ) of the foreign cell by means of another substance called the amboceptor ( ), intermediary, or immune body When this umon takes place the foreign cell is destroyed. The amboceptor, which is not destroyed by heat, is produced by the body cells of the animal receiving the injection of antigen—e.g., foreign cells bacteria, blood cells, or tissue cells of another species, it is specific, having a destructive action only upon the kind of cells which were injected. The complement is nonspecific, its nonspecificity can be demonstrated in the following way. When the immune serum is heated, the complement but not the amboceptor is destroyed and the serum loses its cytolytic action; this is immediately restored, however, by adding any normal serum.

diseases—for example, measles and typhoid fever, to mention only two Many persons appear to be naturally immune to anterior poliomyelitis, and to some other human diseases

Active immunity is that which has been developed as a result of the activity of the body's own cells, as when a person has recovered from an attack of an infectious disease, or has had the antigen administered by injection as explained below.

Immunity produced by the injection of an antibody—e.g., diphtheria antitoxin—formed in the body of an animal or of another person is called passive. Diphtheria antitoxin, for example, is produced by the horse into which the toxin (antigen) of diphtheria has been injected. Serum from such animals is employed in treating diphtheria in patients suffering from this disease. In a similar way, serum from patients who have recently recovered from a disease—such as mensles, scarlet fever, or anterior poliomyelitis—is used to immunize passively another person suffering from the same disease

It is not always easy to say whether a person who has been exposed to some infection or other, but does not "catch" it, is naturally immune to the disease or has been actively immunized against it by having at some time previously contracted it in such a mild form that it was not recognized An apparently natural immunity probably occurs in this way to the virus of anterior poliomyelitis and to the typhoid bacillus

Active immunity may be conferred artificially in order to prevent persons from contracting a disease should they be exposed to it They are injected (inoculated) with an emulsion of bacteria which have been killed by heat or chemical means. The toxin is not destroyed, and it is the essential antigenic substance Such preparations of dead bacteria are called vaccines. This method of immunization is employed against typhoid fever and infections with pyogenic organisms Other methods of vaccination are practiced for protection against the viruses of smallpox, anterior poliomyelitis and rabies Immunization against smallpox is carried out by infecting the subject with a mild form of the disease known as cowpox ' The lymph of calves which have been inoculated with the disease is pricked or lightly scratched into the subject's skin. In the case of rabies (hydrophobia), the victim of the disease is inoculated with a solution of the virus which has been weakened by drying and aging Finally, the toxin of the causative organism-e g, that of diphtheria or of tetanus (lockjaw)-may be rendered nontoxic by chemical treatment without destroying its antigenic properties-that is its ability to stimulate anti

\*Vaccination against smallpox has a long history it was practiced in Eastern employee before Jenner was a rather dangerous procedure. But considering the method employee before Jenner was a rather dangerous procedure. But considering the robte ravages of smallpox at was justified. In this old method the person one if was desired to protect against smallpox was incoulated with a little material (usually was desired to protect against smallpox was incoulated with a little material (usually little and the little and the little material (usually little and little material (usually little and little material) and little material (usually little and little material (usually little and little material) and little material seenging to a desired and in the material seenging to a desired and Jenner an English physician originated the modern method of vaccination against smallpox in 1796 The ledes came to him from the remark of a milkmad "I cannot take the smallpox for I have already had the cowpor. The truth of this statement was known generally among the country people of the district Jenner accordingly inoculated a boy of 8 years named James Phipps with pus from the hand of a dalipmand infected with cowpor. Cowpor though alided to smallpox is a quite mild diverse Two months later Jenner inoculated the boy with virulent small pox virus This the crucial test proved the efficacy of vaccination Phipps 4d not contract smallpox An outstanding example of the value of vaccination against small pox is afforded by the statut set of the Franco-Prussan Max (1870) in the Prussan Army but not in the French vaccination was practiced. Only a few Prissan solders

body production. In other words, according to Ehrlich's theory, the toxophore groups but not the haptaphore groups of the toxin are destroyed. Immunizing agents prepared in this way are called toxoids. By the administration of diphtheria toxoid of young children, deaths from diphtheria, which used to run into high figures annually, have been reduced almost to the vanishing point.

Another immunizing procedure, especially against measles and infectious hepatitis, is the administration of gamma globulin (pp. 58 and 101).

Pain and rest. The pain and temporary loss of function of an inflamed or injured part are definitely protective provisions. Though
we may think that pain has been imposed upon us by a malicious
nature and could be dispensed with, we would soon be destroyed
were it not for the warning which it gives of injury or of a threat of
injury. In a certain nervous disease affecting the sensory fibers in the
spinal cord which carry impulses giving rise to the sensation of pain
and temperature from the hands, the patient is continually damaging
his fingers; sores develop from such injuries, which are neglected because they cause no discomfort. It is also true that the most potent
stimulus for evol.ing a protective reflex, such as jerking the hand away
from a hot object, is a painful one. In the case of the eye, the winking
reflex which shields the eye is extremely sensitive. Even the sight of
an approaching object which might injure the eye will elicit the reflex;
a stimulus applied to the cornea, the anterior surface of the eye which,
if directed to any other part of the body surface would result in nothing more than a mild sensation of touch, when applied to the eye
instantly causes a reflex closure of the eyelid. The cornea is richly supplied with pain fibers which have a very low threshold; almost any
stimulus, except the very blandest such as a gentle jet of warm water,
if it gives rise to any sensation at all, causes pain.

Of an importance as a protective device, nearly as great as that of pain, is the suspension of function in an injured structure. Rest, whether of a broken bone or a lung, enables the healing power of nature (vis medicatrix naturae) to exert most effectively its beneficent action. In some instances, such as in pulmonary tuberculosis where one lung is solely or predominantly affected, man, by artifice, does what nature cannot or fails to do. The severely damaged lung is immobilized—that is, put at rest by collapsing it (as by the injection of air into the pleural cavity of that side) or by cutting one phrenic nerve, and thus paralyzing the corresponding half of the diaphragm. So, also, a broken bone is immobilized by putting it in splints, and in-

flamed eyes are protected by a bandage or dark glasses (though some natural protection is afforded by contraction of the pupil and closure of the cyclids). Pain, of course, not only serves to warn and protect, but forces a part to be put at rest.

A recapitulation of protective mechanisms described elsewhere. There are a number of other defensive measures which the body can take against the variety of adverse conditions and dangers occurring in an ever-changing environment Among such provisions of nature, most of which have been or will be dealt with elsewhere in this book, are: the coagulation of the blood and the other reactions which tend to ensure against fatal hemorrhage (pp 102 and 76); the protection against a general blood infection afforded by the lymph nodes; the cilia and mucus of the upper respiratory passages as well as the cough reflex which serves to prevent the entrance of foreign material into the finer airways of the lungs (p. 258); and the reactions of the sympathoadrenal system (pp 424 and 564).

Finally, it should be mentioned that in order for the various safe-

Finally, it should be mentioned that in order for the various safeguards to serve their purposes most effectively, since they are manifestations of living cells, all physiological processes must be manitained in states of prime efficiency. Unless this is so, the normal biochemical balance of the internal environment (Homeostasis, p. 53) is not secured. When the general health is underpined, a way is opened for invasion by pathogene microorganisms. Diabetic patients, for example, are especially susceptible to infection; victims of this malady, and of nephritis and other general diseases, are often carried off, not directly by the disease itself, but by some "intercurrent infection" such as pneumonia or a septic condition.

Chemotherapy. This is the treatment of disease by the administration of a chemical compound, in many instances synthetic, by means of which the causative microorganisms are destroyed within the body of the patient, or their multiplication arrested. The science of chemotherapy has made enormous strides within the last 15 or 20 years. The aim of the investigator in this field is to prepare a chemical compound that acts selectively against one or more types of pathogenic organism, but, in the dosage used, exerts little or no toxic action upon the patient.

Paul Ehrlich, the founder of the science of chemotherapy, introduced some forty years ago an arsenical compound having a destructive action upon the spirochetes of syphilis. It was named Salvarsan or arsphenamine It was also commonly known as "606" (six-o-six) because it was the last of a long series of compounds (presumably 606) which had been experimented with in the search of a cure for syphilis Several compounds of arsenic and antimony have since been synthesized which are active against the protozoan organisms of malaria, African sleeping sickness, and other tropical diseases

The "sulfa" drugs In 1928, it was discovered by Dr Domagk, a German scientist, that prontosil, a red dye, when administered to mice enabled them to withstand an injection of patheogenic bacteria that was ordinarily lethal. It was found later that a chemical group containing sulfur was the active antibacterial agent. This compound was prepared separately, it is colorless and known as sulfanilamide. The use of this drug and its derivatives—the so called "sulfa" or sulfonamide drugs—has opened up a fresh vista in chemotherapy. Several bacterial diseases from which the death rate was extremely high have been treated with these compounds with spectacular success. A large number of derivatives of sulfanilamide have been synthesized, such as sulfapyridine, sulfanilaziole, sulfacetamide, and sulfadiazine, to mention those in most common use. The sulfonamides are not equally active against different bacterial forms. One drug may be active against one group of bacteria but may fail against another group

Antibacterial agents such as the "sulfa" drugs do not kill the bacteria, like antiseptics or other chemicals such as the arsenical compounds They prevent the bacteria from multiplying They are bacteriostatic 5 (L statos, standing) rather than bacteriocidal (L caedo, I kill)

Antibiotics An entirely new class of highly effective antibacterial agents has been discovered within recent years. These substances, known as antibiotics, are obtained from other living organisms, especially fungi, molds, and bacteria. It was through a happy accident and alert observation that the first of these principles was discovered. In 1929, Dr. (later Sir) Alexander Fleming, an English bacteriologist, noticed that the growth of a culture of staphylococci, which had been exposed to the air of the room, was arrested in the neighborhood of the growth of a mold that had fallen upon the same culture plate. He suggested that the mold had produced a substance mimical to the bacteria Dr. (later Sir.) H. Florey and his associates at Oxford instituted a series of experiments during World War II which re-

a The mode of action of sulfanilamide is believed to be as follows: A factor of the vitamin B complex para aminobenzone acid is an essential requirement for some enzyme system in the bacterial cell. Owing to the similarity of their chemical structures the "sulfa" drug and the vitamin compete for inclusion in the enzyme system. When the drug is taken up by the latter, the entrance of the vitamin into the system is "blocked". The enzyme is thus rendered inactive, and the bacteria are tin able to multiply.

sulted in the isolation of the antibacterial principle from the mold The mold had previously been identified as *Penicillium notatum*, the active principle was, therefore, called *penicilliu* Both Fleming and Florey received the Nobel Prize in 1945

A number of antibiotic principles have since then been isolated from various molds and bacteria. The most valuable of these agents are streptomycin, terramycin, aureomycin, and tyrothricin. Streptomycin is obtained from the funguslike bacteria, Streptomyces it is active against the tubercle bacillus and a number of other pathogenic bacteria. Aureomycin is furnished by another species of Streptomyces, it is active against some virus infections and several types of bacteria. Tyrothricin is derived from a bacterial form found in milk and soil, it is especially valuable for local application in staphylo coccic and streptococcic infections of the skin. Most of the anti-biotics are bacteriostatic in their action, but some are also bacteriocidal. Others are active against pathogenic fungi.

Pathogenic organisms show a remarkable adaptability to the action of various agents employed against them. They tend to become gradually resistant and can then tolerate larger doses of the anti-bacterial substance. When this occurs, the physician's strategy when feasible, is to attack with some other antibacterial substance. Another person may become infected with the resistant bacteria. When, for example, infection is derived from a patient with tuberculosis treated with streptomyen the effectiveness of this valuable drug, in so far as the second infected person is concerned, is greatly reduced.

#### Shock

Shock is a term which has been given to a number of different conditions associated with a state of prostration or collapse. A variety of qualifying words is used to distinguish the several, often unrelated, states to which the term shock has been applied Primary of nerious shock, secondary or traumatic shock, anaphylactic shock, hemorrhagic shock, shell shock, spinal shock, and insulin shock are a partial list of such states. Two severe types of shock—secondary or traumatic shock and anaphylactic shock—will be described.

<sup>•</sup> Terramytin and aureomytin are also known as oxysteticycline and chloriters recline respectively. Since the volation and great success of periodlin the most in tensive search for other antibiotics has been instituted besides the four mentioned above a host of new ones have been volated. Some of these have proved of great value—e.g. erythromycin chloromycenn bacinacin carbomycin etc.—while others are not very active or are too toxic for use.

Secondary or traumatic shock. Immediately following a severe wound, the subject is weak and feels faint and may pass into a state of collapse as a result of pain, fright, apprehension, or other emotion. This state immediately following an injury is known as primary or Inis state immediately following an injury is known as primary or nervous shock. It is of comparatively small physiological interest and does not end fatally in a previously healthy person. If the wound is of moderate severity, the subject's condition improves after a pe-riod of rest and appropriate treatment. If, on the other hand, the injury is very severe—especially if much blood has been lost—the wounded person gradually sinks into a condition of great weakness with falling blood pressure. This state of extreme prostration which becomes most profound in from 3 to 5 hours following the injury is known as secondary or traumatic shock. It is also known as deferred. wound, or surgical shock. The chief features of traumatic shock are associated with failure of the peripheral circulation—very low blood pressure, constriction of the arterioles with dilatation of the capillaries. There is usually also a loss of plasma from the circulation into the tissues, with a resulting concentration of the blood in red cells, and a reduction of the total volume of blood in the circulation. The patient usually remains conscious but is extremely weak and listless, with a cold clammy skin and a thin faint pulse. Shock has been the subject of intensive research, especially during the two world wars, but the underlying cause leading to its development has not been entirely revealed. Many believe that it is due simply to a reduction in blood volume. either as a result of the loss of plasma into the tissues at the site of the injury or to hemorrhage. Others maintain that some other factor, as well-probably a blood-pressure lowering principle produced by the liver and discharged into the circulation—is responsible for the profound fall in blood pressure.

The only treatment which is of any avail in shock is the transfusion of plasma or of blood, but all too frequently even this fails to save the patient,

Anaphylaxis, anaphylactic shock. If an animal is given an injection of a foreign protein—such as serum albumin of another species—and after a period of 10 days or more is given a second injection of the same protein, a sudden and great fall in the injected animal's blood pressure results. Death usually occurs within a few minutes. The first injection of protein is said to "sensitize" the animal to the second injection which may be only a very minute quantity. The foreign protein acts as an antigen (sometimes called specifically the anaphylactogen) causing the body cells, in the interval between the

two injections to produce an antibody (p. 215), which reacts in some way with the second dose of protein. The precise nature of this antigen antibody reaction and its relation to anaphylactic shock are not altogether clear, but it is generally believed that it is associated with the liberation of histamine (an amine derived by the loss of carbon dioxide from the amino acid histidine) and that this amine or a closely similar substance is responsible for the effects. The anaphy lactic reaction shows pronounced species differences with respect to the tissues upon which the effects are manifested. In the dog for example the chief anaphylactic effects apart from the fall in blood pressure are upon the liver and alimentary tract there are vomiting diarrhea and engorgement of the vessels of the liver stomach and bronchioles occurs and the animal dies in asphyxia The subcutane ous injection of antigen into a rabbit causes a severe tissue reaction at the site of the injection where an abscess forms and death of tis sue ultimately occurs. If this animal receives the second dose of pro-tein intravenously circulatory failure results and the animal passes into a state resembling traumatic shock. Most of the effects enumerated as well as the species differences can be induced by injec-tions of histamine which strengthens the belief though does not absolutely prove that histamine liberation is the cause of the ana-phylactic manifestations. Histamine can be extracted from normal trissues but it exists apparently in some innocuous form Large amounts are found in the blood of the dog when anaphylactic shock supervenes and in the blood and lungs of the guinea pig. The mech anism which causes the liberation or accumulation of histantine in these tissues in anaphylactic shock is unknown

In man death from anaphylactic shock is unknown. In man death from anaphylactic shock has occurred on several occasions as a result of the injection of foreign serum (horse or bovine) for therapeutic purposes as in the treatment of diphtheria or tetanus the patient having been sensitized by the injection of serum at some time previously. Milder reactions of an anaphylactoid nature, such as skin rashes vomiting or circulatory disturbances, are not very uncommon following the injection of foreign serum and are spoken of as serum sickness."

Allergic conditions—such as hay fever (from the inhalation of pollen dust), gastrointestinal upsets rashes etc caused by eating certain foods by susceptible persons—are anaphylactoid in nature Essentially similar mechanisms are involved

# Respiration

All Irving things, with a few exceptions absorb oxygen. The oxygen combines with the carbon and hydrogen furnished by food material Carbon dioxide (CO<sub>2</sub>) and water (H<sub>2</sub>O) are produced. These oxida tions take place in the cells of the tissues, they generate heat and furnish the organism with energy for the performance of work (see Chap 9). The carbon dioxide, so formed is eliminated, the oxidation of the hydrogen constitutes an important source of body water (the water of metabolism). The exchange of gases (O<sub>2</sub> and CO<sub>2</sub>) be tween an organism and its environment is termed respiration.

In unicellular organisms and many multicellular forms of lower orders the gaseous ex change takes place directly be tween the cells and their sur roundings (see Fig 6 1). Higher forms of animal life are composed of cells of which the great mass are removed from direct contact with the external environment. But the cells of such higher forms like those of the most primitive are bathed in fluid. This tissue fluid we may speak of as the *internal e* 





Fig 6.1 Respiration of a unicellular organism (e.g. an amoeba)

may speak of as the internal environment from it the cells absorb oxygen (as well as nutriment) and into it the cells discharge their waste materials including carbon dioxide (Fig 6 2) Through the medium of the blood vascular system oxygen is transported from the external to the internal environment, thus a steady supply of oxygen to




resp ration Capillary shown filled

the cells is secured By the same means, carbon dioxide is carried from the tissue fluids. In air breathing animals, the exchange of gases between the blood and the external environment takes place in the lungs This is called external respiration. In fish it is a function of the gills The exchange of gases between the tissue cells and their fluid environ ment is termed internal respiration

## An Outline of the Structure of the Respiratory Organs

The air passages These are the nasal cavities the pharynx larynx trachea bronchi and bronchioles (see Figs 63 and 64) The in terior of the nose is divided into two lateral halves by the nasal sep tum From the lateral wall of each nasal cavity spring three spurs of bone called respectively the inferior middle, and superior turbinates (or conchae) These incompletely divide each side of the nose into three passages (the meatuses of the nose) Situated between the su perior turbinate bone and the floor of the skull is a small recess con taining the olfactory epithelium (Chap 13) The main air currents do not enter this recess the meatuses of the nose alone serving as airways The meatuses are lined with ciliated epithelium the move ments of the cilia are toward the exterior, thus mucus and dust or any other foreign particles that may enter the nasal passages in the in spired air, are carried to the nostrils

Fig 6.3 The inner surface of the outer wall of the nose. 1, inferior concha (turbinate); 2, middle concha; 3, superior concha; 4, olfactory nerves; 5, olfactory bulb; 6, opening of auditory (Eustachian) tube The arrows indicate the course taken by the afr in breathing



The nasal cavities open posteriorly into the pharynx which, therefore, serves as a common passage for the transmission of air to the larynx, and of food from the mouth to the esophagus. During swallowing, the food has the right of way, the respirations being then for a moment arrested.

The larynx. The larynx is situated in the neck in front of the lower part of the pharynx and at the commencement of the trachea or windpipe. It contains the vocal cords, and will be described more fully later.

The trachea and bronchi. The human trachea is a tube about \(\frac{2}{3}\) inch in diameter and 4 inches long, extending from the larynx to a little




Fig. 64 Showing trachea and bronchi. Portion of the lung tissue removed to show branchings of the bronchial tree.

below the upper boundary of the chest, here it divides into two branches—the right and left bronchi. Each bronchus enters the corresponding lung and divides, like the limbs of a tree, into a number of smaller branches. The larger of these are also called bronch, but the finer terminal twigs are referred to as bronchioles. It is, of course, imperative that this important airway always be kept open. The patency of the trachea is ensured by the presence of a series of C shaped rings of cartilage which support its wall in front and at the sides. The gaps in the cartilages posteriorly, as well as the intervals




Fig 65 Showing the Jungs in relation to the heart and chest wall R.L. right lung L.L. left lung H, heart N, mpple D, dome of the dia phragm

between them, are bridged by a strong fibroelastic membrane. The trachea is lined by ciliated columnar epithelial cells among which are many gobiet cells. The walls of the two main bronch are constructed upon a plan almost identical with that of the trachea, but the cartilage in the walls of the smaller bronch is in the form of thin isolated and irregular plates which become smaller and fewer in number with each successive branching of the bronchial tree.

The bronchioles are about 1 mm or less in diameter. Their walls are composed of smooth muscle lined by mucous membrane of the same character as that lining the trachea and bronchi except that goblet cells are absent. Cartilage is entirely lacking. The muscle fibers of the bronchioles are arranged circularly. Their contraction, there fore, causes narrowing of the bronchiolar lumen. In astimatic attacks, marked bronchiolar constriction occurs and is responsible for the respiratory distress characteristic of this condition. The muscle of the bronchioles receives its nerve supply from both the vagus and the sympathetic, the former is constructor, the latter dilator in action. The bronchiolar muscle is also relaxed by adrenaline and ephedrine.

The lungs. The left lung is divided by a fissure into two lobes, an upper and a lower (see Fig. 6.5). Two fissures divide the right lung into three lobes. The bronchi and all but the very finest bronchioles possess no truly respiratory function; no interchange of gases can




Fig. 6.6 Diagram of a lung unit. AT, atrium; AS, air sac.

occur across their relatively thick walls. They serve merely as airways.

The lung tissue proper consists of an immense number of air spaces arranged in clusters. Each cluster is called a lung unit. Each of the terminal twigs of the nonrespiratory part of the bronchial tree (terminal bronchioles) opens into a lung unit. One of the latter with its several named parts—respiratory bronchiole, alveolar ducts, air

230



Fig 67 Microscop cal section of pulmonary tissue

sacs and pulmonary alveoli—is illustrated diagrammatically in Fig ure 6.6 The pulmonary alveoli are seen as from four to six outpouch nigs of the cavity of each alveolar sac A single layer of flat cells and a network of fine elastic fibers compose their walls, a rich network of blood capillaries surrounds them Moreover, according to most observers the walls of the alveoli are incomplete and show numerous gaps. In many places therefore the capillary wall alone intervenes




Fig. 68 Model of a lung unit

between the blood and the alveolar air. Thus the blood as it flows through the lungs is separated from the air in the alveoli by, at the most, two membranes—the alveolar and capillary walls—so extremely thin that little hindrance is offered to the free exchange of the respiratory gases. The number of pulmonary alveoli in both lungs has been estimated to be about 750,000,000.

A microscopical section of pulmonary tissue is shown in Figure 6.7, and a model of three lung units in Figure 6.8.

#### The Mechanics of Respiration

The chest or thorax is a closed cavity, bounded laterally and behind by the ribs and the vertebral column; in front by the ribs and sternum (breast bone); below by a dome-shaped sheet of muscle (the diaphragm), which separates it from the abdominal cavity; and above by the upper ribs and the tissues of the neck. The lungs, heart, and great blood vessels almost completely fill the thorax.

The lungs are covered by a thin membrane called the pleura, which passes from each lung at its root (i.e., at the point where the main bronchus and pulmonary vessels enter it) on to the interior of the chest wall and upper surface of the diaphragm. Thus, two membranous sacs, called the pleural cavities, are produced, one on each side of the chest, between the lungs and the thoracic boundaries. Since the pleura covering the lung and that lining the chest are in contact under all conditions of health, no actual space exists; these so-called cavities are potential ones only. In disease, however, air, fluid, or blood may collect and separate the pleural layers to produce an actual space.

The part of the pleural membrane lining the thoracic wall is called the parietal pleura (L. paries, gen. parietis, a wall), while that portion covering and firmly adherent to the surface of the lungs themselves is called the pulmonary or visceral pleura. Inflammation of the pleura is called pleurisy. The inflammatory process may cause the production of little or no fluid—dry pleurisy—or may be accompanied by the exudation of a considerable amount of clear fluid—pleurisy with effusion. If the fluid becomes purulent, the condition is called empyema.

The central part of the thoracic cavity lying between the lungs, and bounded in front by the sternum and behind by the vertebral column, is known as the *mediastinum*. It is divided for the purpose of description into an upper and a lower part, called the superior and the in232

ferior mediastinum, respectively. The latter consists of three parts—anterior, muddle, and posterior mediastina. The middle mediastinum contains the heart, enclosed in the pericardium, and the commencement of the aorta The division of the trachea into the right and left bronchi is contained in the posterior mediastinum. The anterior mediastinum, in front of the heart, is almost obliterated by the approximation of the two pleural sacs. It contains only a little areolar tissue and two or three lymph nodes.

The intrapleural pressure. The pressure between the layers of the pleura (intrapleural pressure), and indeed throughout the thorax




Fig. 69 Tracing from x-ray photograph demonstrating the elasticity of the bronchial tree. A, full expiration, B, full inspiration. (After Macklin.)

generally (intrathoracic pressure), is below that of the atmosphere. This subatmospheric or so-called negative pressure is due to the elasticity of the lungs. Before birth the lungs are airless, the air sacs are of small size and contain a little fluid. The thoracic cavity is filled by the almost solid lungs, the heart, and great vessels. The pressures within the pleural cavities and in the interior of the lungs are equal. With the first breath after birth, the thorax is expanded in all its diameters. This reduces the pressure within the thorax—that is, on the pleural aspects of the lungs—and tends to reduce the pressure within the air sacs (intrapulmonary pressure). The air sacs, however, are in communication with the outside air through the air passages. Atmospheric air, therefore, enters the lungs to equalize the pressure, and, were the lung tissue inelastic, equalization of pressure between the atmosphere and the pleural cavities would also result. But the bronchial tree and the lung substance itself are richly supplied with elastic tissue. When the lungs expand and fill the enlarged thoracic cavity, the elastic tissue is stretched. The lungs in postnatal life are, therefore, constantly under tension with a tendency to recoil to their original dimensions. This pull of the elastic lungs away from the thoracic

walls, though it cannot separate the layers of the pleura, creates a "negative" pressure.

Throughout life the intrapleural pressure undergoes rhythmical variations with the respiratory movements as a result of changes in the thoracic dimensions and the consequent variations in the degree of stretching to which the lungs are subjected (see Fig. 6.9). The intrapleural (or intrathoracic) pressure is around -6 mm Hg during inspiration and -2.5 mm during expiration. The importance of the




Fig. 6 10 (Description in text.)

subatmospheric pressure within the thorax in aiding the return of venous blood to the heart has been pointed out (p. 153).

The respiratory movements—the intake and expulsion of air. The drawing of air into the lungs is called *inspiration*, and its expulsion expiration. The capacity of the thorax is enlarged during inspiration and reduced during expiration. These two movements alternate rhythmically at a frequency, in the adult, of from 18 to 20 per minute. The rate is more rapid in infants and young children.

At the end of expiration, the pressure of air within the lungs (intrapulmonary pressure) equals or nearly equals the atmospheric pressure. The inspiratory movement, as mentioned above, reduces the
pressure upon the pleural surfaces of the lung. The air sacs are dilated thereby and the air within them is rarefied—i.e., reduced below
that of the atmosphere. Air, therefore, flows into the lungs until the
intrapulmonary pressure again almost equals that of the atmosphere.
During expiration the intrapleural "negative" pressure is reduced, the
air sacs return to their previous dimensions, and the intrapulmonary

pressure rises above that of the atmosphere; air is therefore expelled from the lungs.

It should be emphasized that the lungs play a purely passive role in the respiratory movements. The changes in volume which they undergo, and to which the intake and expulsion of air are directly due, are brought about solely through changes in the capacity of the thoracic cavity. The inspiratory and expiratory movements of the thorax may be compared to the opening and closing of a fire bellows. A pair of bellows, which may be taken to represent the thorax, is shown in Figure 6 10. Two clastic bags, contained within it and communicating with the outside air through the nozzle, correspond to the lungs and air passages. The space surrounding the bags is hermetically sealed and corresponds to the pleural cavities. Opening the bellows ("expiration") causes the bags to expand and fill with air; closure of the bellows ("expiration") to the stop S causes their partial deflation. Since the bags, like the lungs, are made of elastic material, they are always tending to recoil against the distending force. This "pull," as between the layers of the pleura (i.e., in the pleural cavity), produces a "negative" pressure between the bags and the walls of the bellows

Having outlined the general principles underlying the ventilation of the lungs, the muscular mechanisms whereby the changes in thoracic capacity are brought about will now be described.

The enlargement of the thorax during inspiration. All diameters—vertical, anteroposterior, and transverse—are increased during the





Fig. 6.11 Showing position of the diaphragm at the end of a full inspiration and a full expiration, respectively.

inspiratory phase. Increase in the vertical diameter of the thorax is brought about by the descent of the diaphragm. This is an arched tendomuscular sheet which separates the thorax from the abdominal cavity. The muscle fibers arise from the spinal column and lower ribs and, arching upward and inward, are inserted into a central leaf-shaped tendon. This tendon forms the highest part or dome of the diaphragm and is called the central tendon of the diaphragm (Fig.

6.11). The central tendon of the diaphragm blends with the under surface of the pericardium; on either side of the pericardium the diaphragm is in contact with the bases of the lungs. When the muscular fibers of the diaphragm contract, the structure shows little change in shape, but is seen under the x-rays to move downward as a whole very much like the stroke of a piston (Fig. 6.11). The diaphragm is the most important muscle of respiration, about 60 percent of the volume of the air respired being due to its action.

During inspiration the thorax is enlarged in its anteroposterior diameter by the movement, into a more horizontal position, of the third, fourth, fifth, and sixth pairs of ribs, which at the end of ex-





piration slope obliquely downward and forward (Fig. 6.12 A and B.) This movement is brought about by the contraction of the external intercostal muscles. The fibers of the latter take a slanting direction across the intercostal space from the lower border of one rib downward and forward to the upper border of the next rib below. When the external intercostal muscles contract, the anterior ends of the four pairs of ribs are elevated and the sternum thrust forward and upward. Any downward pull which the contracting external intercostal muscles exert upon the upper of two adjacent ribs is antagonized by the upward pull of certain muscles of the neck (scalenus muscles) which are inserted into the first and second ribs. Thus the upper of any two adjacent ribs serves as a fixed point for

235

the intercostal fibers, so that, when they contract, the lower rib of each pair is raised anteriorly and assumes a more horizontal position. The part of the thorax bounded by the seventh, eighth, ninth, and tenth ribs is increased in its transverse diameter by rotation of these

ribs around an axis directed from the sternum obliquely backward



Fig 6 13 Illustratbucket-handle movement of lower

and a little outward (Fig. 6.12 C). At the end of expiration these ribs are bowed outward and downward, and their movement during inspiration has been compared to raising a bucket handle from its position of rest toward the horizontal plane (Fig. 6.13).

Expiration. This phase is effected mainly in a passive manner in quiet normal breathing. At the end of the inspiratory phase, the external intercostal muscles and the diaphragm relax. The walls of the thorax, by virtue of their own weight and their inherent elasticity, together with the pull exerted by the recoil of the elastic lungs and

bronchi (see Fig 69), return to their original (preinspiratory) posi-tion The relaxed diaphram ascends, being drawn up by the "nega-tive" intrathoracic pressure and also by the elastic recoil of the lungs

Accessory muscles of respiration. In forced or difficult breathing, certain muscles come into play to facilitate the respiratory movements, in inspiration, the nostrils and the opening between the vocal cords (glottis) are enlarged by muscular action, thus permitting a freer entrance of air into the lungs. The muscles of the neck (anterior and middle scalenus muscles and the sternomastoids) contract and raise the first rib, lifting the sternum upward and forward, and increasing the inspiratory movement of the succeeding ribs. Thus the capacity of the thorax is materially increased. During a forced or difficult expiration, the abdominal muscles contract and, increasing the intra-abdominal pressure, and in the ascent of the diaphragm. The internal intercostal muscles exert a weak expiratory action; their fibers run in a direction the reverse of that of the external intercostals; their contraction depresses the ribs. Thus, expiration becomes a more active movement than normal.

Artificial respiration. When the respiratory mechanism has failed, as in the apparently drowned or as a result of carbon monoxide poisoning, electrocution, or other cause, some artificial means must be employed to bring air into the lungs until natural breathing is re-sumed or until all hope of this occurring has been abandoned. In any method of artificial respiration, promptness in starting the treatment is of first importance. So long as the heart beats, the tissues for a time are able to gain a small amount of oxygen from the circulating blood. But the heart is peculiarly susceptible to oxygen deprivation and, unless the air in the lungs can be renewed and the blood oxygenated, it will soon cease to beat. Nervous tissue also, especially the higher centers of the brain, does not survive for more



Fig 6 14 Holger Nielsen method of artificial respiration. (From First Aid, Metropolitan Life Insurance Co )

than a few minutes after its oxygen supply has been arrested When the heart ceases to beat and can no longer maintain the circulation, methods of resuscitation are usually of no avail. However, it is often impossible for one who is not medically qualified to detect the beat of the heart when it is very weak. For this reason artificial respiration should be continued until natural breathing has been restored or until a physician has pronounced the death of the patient. Not until then should hope be abandoned.

Most methods of artificial respiration are designed to increase and reduce alternately the capacity of the thorax in such a way as to draw air into the lungs and expel it again. Thus the blood is oxy-

<sup>1</sup> If a physician is not available, efforts at resuscitation should be persisted in until rigor mortu (stiffness of the muscles after death) is clearly evident.

240

apparatus A flexible collar is fitted around his neck, so as to prevent the leakage of air (Fig 6 15). A motor driven pump causes alternate variations in the air pressure within the chamber. Reduction in air pressure from 3 to 4 mm below that of the atmosphere causes expan sion of the chest (inspiration). The next moment the pressure within the chamber becomes atmospheric, the chest returns to the resting position and expiration occurs. Portholes for tending the patient are provided in the walls of the apparatus. The patient is kept continuously in the chamber until respiratory muscles which have not been paralyzed, or unparalyzed fibers within the affected muscles (e.g., diaphragm or intercostals), have developed sufficient strength to carry out, unaided, the movements required for adequate ventilation of the

A rocking bed, operated by a motor, has recently been designed to carry out artificial respiration by the Eve method over long periods It is claimed to induce better ventilation of the lungs than the Drinker method

In cases of apparent drowning or gas poisoning, it is usually of great benefit to add oxygen to the inspired air. Carbon dioxide in a concentration of about 7 percent is also frequently employed the object being to stimulate the respiratory center (p. 253) and bring on natural breathing. Oxygen and carbon dioxide are also used in patients suffering from pneumonia and other conditions in which though the patient is able to breather the blood is being imperfectly oxygenated, the gases being passed into a tent covering the patients. head and trunk

### The Subdivisions of the Lung Air

Only about one tenth of the total volume of air that the lungs are able to inspire and expire in a single respiration (see vital capacity, p 242) moves into and out of the lungs during ordinary quiet breathing this, which amounts to about 500 cc, is called the tidal volume. In an inspiration of maximum depth six times this amount (namely, 3000 cc) can be inhaled. This is termed the inspiratory capacity (it in cludes tidal volume). If one empties his lungs as completely as possible (forced expiration) after an ordinary expiration, about 1000 cc of air can be expelled from the lungs. This air, which can be expelled by an extra effort is called the expiratory reserve volume. The lungs cannot, of course be emptied completely for they are kept in the partially expanded state by the subatmospheric pressure within

the pleural cavities Even after the most forcible expiration, from 1000 to 1500 cc of air remain in the lungs. This large volume of air can be expelled only after death by opening the thoracic cavity, thus establishing communication between the pleural cavities and the

Table 6-1

Tidal volume		500 cc
Inspiratory capacity (which includes		
tidal volume)*	3000 cc	
Expiratory reserve volume	1000 cc	
Vital capacity	4000 cc	
Residual volume	1200 cc	
Total lung capacity	5200 cc	

atmosphere, it is called the residual volume (see Table 6-1 and Fig 6-16) But, the lungs, after the chest has been opened or even after they have been removed from the body, contain air entrapped within the air sacs. This part of the residual air is termed the minimal air. To it the lung tissue owes its buoyancy when put into water. The lungs of a stillborn child, since they have not been expanded by

If the tidal volume is not included it is called the inspiratory reserve volume




Fig 6 16 Showing subdivisions of the lung air.

breathing and contain no air, sink in water. This fact is the basis of an important medicolegal test in cases of suspected infanticide, by means of which it is possible to determine whether or not an infant had been born alive

The maximum quantity of air which can be expired after the deepest possible inspiration-that is the sum of the inspiratory capacity-



Fig 617 Spirometer A, graduated glass bell B water d mouthpiece g counter weight r pul lev (After Reichert )

and the expiratory reserve volume is called the sital capacity. Athletes and other per sons in good physical condition have, in general greater vital capacities than those of the same size who lead a sedentary life or are debilitated. The vital capacity is therefore used as one of the criteria of physical fitness. In health it bears a relationship to the surface area of the body, amounting to 2500 cc per square meter of body surface for the average man, 2000 cc for women, and 2800 cc for athletes For a man of average build (17 square meters of body surface), it amounts, therefore, to (1 7 × 2500 -) 4250 cc Certain diseases of the lungs chest, and heart reduce the vital capacity below the normal standard for the individual

The vital capacity is measured by having the subject inspire fully, and then empty his lungs as completely as possible into an instrument called a spirometer (see Fig 6 17)

The average values of the main subdivisions of the lung air in round figures are tabulated in Table 6-1 on page 241

Alyeolar air and dead space air The air in the air sacs -that is, the air in contact

with the true respiratory part of the lungs-is called the alveolar air The space enclosed by the nonrespiratory part of the lungs-namely, the bronchi and muscular bronchioles, trachea larynx, pharynx and nasal cavities-is called the dead space it has a capacity of about 150 cc The air within this space is called the dead space air

#### The Chemistry of Respiration

The exchange of the respiratory gases (oxygen and carbon dioxide) between the blood and the air in the lungs, and between the tissue cells and the blood, follows those physical laws which govern the behavior of gases in general. Some space will be devoted, therefore, to an account of the properties of gases.




Fig 6 18 Illustrating the diffusion of gases. The two gas storage cylinders contain the same kind of gas at unequal pressures. In the cylinder on the left, the gas is at the lower pressure. When samples of gas from each cylinder are allowed to enter the glass chamber A, the molecules, as indicated by the small circles, intermingle freely and become evenly distributed (as in B) with consequent equalization of pressure throughout.

The kinetic theory. The behavior of a gas is explained upon the theory that its molecules are in ceaseless motion. They move at high velocity and if unobstructed in any way become separated from one another by immeasurably great distances A gas, therefore, is capable of expanding to infinite dimensions, its molecular concentration diminishing proportionately as it expands. When confined, the gas molecules strike numberless blows upon the walls of the container. The pressure exerted by a gas is attributed to this ceaseless molecular bombardment. It follows, therefore, that the greater the number of molecules within a given space (i.e., the higher their concentration), the higher will be the gas pressure.

If two samples of the same gas but of different molecular concen-

244

trations be brought together rapid and even distribution of the mol ecules takes place between the two samples, with consequent equal ization of the pressure throughout any space in which the gas is confined (see Fig 6 18) The final pressure has a value somewhere between the pressures of the separate samples Gas molecules are therefore said to move from a point of higher to one of lower pres cure

When two dissimilar gases either at the same or at different pressures are brought together, each behaves as though it were the only kind present. The molecules of each gas become evenly dispersed throughout the space its molecular concentration and in consequence its pressure becoming everywhere equal. After mixing the molecular concentrations and therefore, the pressures of the individual gases may differ widely from one another for, as just mentioned, they be have quite independently of one another their final concentrations depending upon the original concentrations of the respective samples. In order to realize the speed with which dispersion of the molecules occurs one has only to recall how quickly some odorous or pungent gas permeates the atmosphere. The process whereby samples of the same gas at different pressures or of different gases at the same or unequal pressures become uniformly mixed is called diffusion (Fig. 618). When two dissimilar gases either at the same or at different pres 6 18)

Though the molecular movements are very much slower, liquids and solids in solution behave in the same way. For example, a basin of water in which a bottle of red ink is opened carefully, so as to avoid mechanical mixing becomes after a time of a uniform color through out. Molecules of gases also diffuse into liquids which are in contact with them until the gas pressures within and without the liquid are

The properties of gases are stated more specifically in the following

laws

The gas laws. Boyle's law The temperature remaining constant reduction in the dimensions of the space in which a gas is confined increases the pressure of the gas and vice versa. In other words at constant temperature the pressure of a gas varies inversely with its volume—the product of the pressure and the volume is constant te doubling the pressure reduces the volume by half, or conversely doubling the volume of the container reduces the pressure by half.

Charles' (or Ga) Lussaes) law For each rise in temperature of 1° C a gas expands by approximately \$\frac{1}{278}\$ of its volume at 0° C provided that the pressure is kept constant, or, its volume remaining con

stand, the pressure of a gas increases by  $\frac{1}{2}$  for each degree (Centigrade) rise in temperature.

Dalton's law of partial pressures. A quantity of gas when mixed with other gases exerts the same pressure as it would exert were the other gases not present. The total pressure of a mixture of gases is, therefore, the sum of the pressures of the individual gases in the mixture, each gas exerting a pressure proportional to its percentage in the mixture. For example, the atmosphere (dry) at sea level has a pressure of 760 mm Hg; it contains 20.96 percent of oxygen. The partial pressure exerted by oxygen, in the air at sea level is, therefore,

$$\frac{20.96}{100} \times 760 = 159.2 \text{ mm Hg}$$

or approximately \frac{1}{5} the pressure of the atmosphere at sea level.

Henry's law of the solution of gases. The temperature remaining constant, the quantity of gas which goes into solution in any given liquid is proportional to the partial pressure of the gas For example, water when exposed to the atmosphere absorbs the constituent gases—oxygen, nitrogen, and carbon dioxide—in quantities proportional to the pressure which each exerts in the air mixture. The partial pressure or tension of each gas in the water will be the same as its partial pressure in the atmosphere. The liquid is then said to be equilibrated with the gas.

The compositions and partial pressures of the gases in atmospheric, expired, and alveolar airs. Atmospheric air contains 20.96 volumes percent of oxygen, 79.00 percent of nitrogen, and 0 04 percent of carbon dioxide. It also contains minute amounts of argon and other rare gases. The approximate partial pressures of these gases in the atmosphere (that is, in inspired air) are, therefore, 158, 596 and 0,30 mm Hg, respectively. In expired air the pressure (tension) of oxygen is lower (since its percentage is lower) and that of carbon dioxide higher, than in inspired (atmospheric) air. In the air of the pulmonary alveoli (alveolar air), the oxygen pressure is lower again, and the carbon dioxide pressure higher than in the inspired air (see Tables 6-2 and 6-3, p. 246). Thus, the pressure of oxygen declines from inspired air to alveolar air, and that of carbon dioxide from alveolar air to inspired air. Nitrogen is an inert gas so far as respiration is concerned—that is, it is neither used in the body (like oxygen) nor produced (like carbon dioxide). The difference between the pressure of nitrogen in the expired and alveolar airs, on the one hand, and in the inspired air, on the other (see Table 6-3), is due simply to changes

Table 6-2

Volumes percent of gases in dry, inspired, expired, and alveolar airs (average flagres)

	Volumes percent		
	Inspired (atmospheric) air	Expired air	Alveolar air
Oxygen Carbon dioxide Nitrogen (with argon and other rare gases in minute quan	20 96 0 04	163 40	14 2 5 5
tities)	78 00	79 <i>7</i>	803
Totals	100 00	100 0	100 0

Table 6-3

Partial pressures of gases in inspired, expired, and alveolar airs (average figures) Barometer reading 760 mm Hg

		Partial pressures		
	Inspired	Expired	Alveolar	
	air	air	aır	
	mm Hg	mm Hg	mm Hg	
Oxygen	158 2	116 2	101 0	
Carbon dioxide	0 3	28 5	40 0	
Nitrogen	596 5	760 0	760 0	
Water vapor	5 0	47 0	47 0	
Totals	760 0	760 0	760 0	

in the pressures of the other gases and to the water vapor added to the air in the lungs, for it should be remembered that the total pressure of expired air or of alveolar air must be the same as that of the atmosphere

The partial pressure of water vapor in the air of the lungs amounts to about 47 mm Hg. The value for the barometric pressure, less 47, must therefore be used in calculating the partial pressure of one or other gas in the expired or the alveolar air from its percentage. For example, if the expired air contains 16.3 percent of oxygen and the barometer registers 760 mm, then the partial pressure of oxygen is

$$\frac{16.3 \times (760 - 47)}{100} = 116.2 \text{ mm Hg}$$

'The diffusion of gases between the atmosphere and the air in the alveoli. As has been pointed out, the lungs are not filled with fresh air during inspiration nor completely empited during expiration (p 240) The tidal air amounts to only about one sixth of the total quantity of The tidal air amounts to only about one sixth of the total quantity of air contained in the lungs, under ordinary circumstances. The ventilation of the air sacs is brought about largely through diffusion—that is, the movement of gas molecules from a point of higher to one of lower pressure. Diffusion is very materially aided by mechanical mixing caused by the respiratory movements, especially during the elastic recoil of the lungs at the end of inspiration. Oxygen diffuses into the alveolar air from the fresh air drawn into the respiratory passages. (dead space) during inspiration, and carbon dioxide diffuses from the dueate space) during inspiration, and carbon dioxide aimuses from the alveolar air into the air of the dead space (See Table 6 3 and compare figures for partial pressures of oxygen and carbon dioxide in inspired, expired, and alveolar airs.) During expiration a part of the air (about 500 cc of the total 2500 or 3000 cc in the lungs) is forced air (about 500 cc of the total 2500 or 3000 cc in the lungs) is forced from the air sacs. About 350 cc of this is expelled to the outside to gether with about 150 cc which had filled the respiratory passages. The last 150 cc of alveolar air replaces that which has been swept from the dead space. At the next inspiration the column of air which was in the dead space is drawn back again into the air sacs, together with 350 cc or so of atmospheric air. Another 150 cc of atmospheric air fills the dead space

The depth to which fresh air is drawn into the air sacs varies, of course, with the quantity of air inspired In very shallow breathing, pure atmospheric air may not fill the dead space, whereas in deep breathing it penetrates beyond the respiratory bronchioles (p 229)

The exchange of gases between the alveolar air and the blood. Each

The exchange of gases between the alveolar air and the blood. Each minute during bodily rest from 4 to 6 liters of blood are delivered to the lungs by the pulmonary artery. In traversing the capillaries of the lungs, the red blood corpuscles are separated from the alveolar air by the exquisitely thin and highly permeable membranes forming the alveolar and capillary walls. In their passage through the capillaries which surround the alveoli, the red cells are for the most part in single file. Thus, a thin film of blood, having an area calculated at about 1000 square feet, is exposed to the alveolar air. The blood coming to the lungs (i.e., the mixed venous blood from all parts of the body), having given up a part of its oxygen load to the tissues

218

and having received therefrom a somewhat smaller volume of carbon dioxide, has therefore a relatively low pressure of the former gas and a relatively high pressure of the latter. The capillary blood and the alveolar air (which at the end of inspiration has a relatively high pressure of oxygen and a low pressure of carbon dioxide) come rapidly into gaseous equilibrium, oxygen diffusing from alveolar air to blood and carbon dioxide from blood to alveolar air. The blood leaving the lungs (arterial blood) therefore contains more oxygen and less carbon dioxide than does venous blood; the oxygen and carbon dioxide pressures show corresponding differences, the oxygen pressure being around 100 mm Hg in arterial and 40 mm in venous blood; the carbon dioxide pressures are about 40 to 46 mm Hg, respectively (see Pl. 6B).

The volumes percent (i.e., the number of cubic centimeters of gas in 100 cc of blood) and the pressures of oxygen and carbon dioxide in arterial and in mixed venous blood are given in Tables 6-4 and 6-5.

Table 6-4

Pressures of oxygen and carbon dioxide in arterial and in mixed venous blood (averages). Compare with Table 6-3.

	Pressures		
	Arterial blood	Mixed venous blood	Difference *
Oxygen	100 40	40 46	60 6

Table 6-5


Volumes percent of oxygen and carbon dioxide in arterial and in mixed venous blood (averages)

	Volumes percent		
	Arterial blood	Mixed venous blood	Difference
Oxygen	19 50	12.5 56	6.5 6.0

Carbon dioxide is much more diffusible (30 times) than oxygen, so that rapid diffusion of the former gas takes place at a smaller pressure difference.

The gaseous exchanges in the tissues. The gaseous exchanges taking place in the tissues (internal respiration) are the reverse of those in the lungs (external respiration). The oxygen pressure in the tissues is low, the earbon dioxide pressure high. The blood in passing through the capillaries of the systemic circulation (e.g., of muscles, skin, abdominal organs, etc.) gives up from 5 to 7 volumes percent of oxygen (depending upon the particular type of tissue and its activity at the time) and absorbs from 4 to 7 volumes percent of carbon dioxide. The gases diffuse across the capillary membrane from plasma to tissue

Fig 6 19 The oxygen dissociation curve of hemoglobin Heavy shading shows reduced hemoglobin and light shading oxyde ingolobin Percentage saturation of hemoglobin with oxygen along vertical lines (ordinates), oxygen presure (tension) in mm Hg along the horizon tal lines (abcessae)



fluid (oxygen), and from tissue fluid to plasma (carbon dioxide) (See Fig 6 19)

The transport of the respiratory gases in the blood. Oxygen transport When human blood has been fully oxygenated (as after exposing it, at body temperature, to the atmosphere at sea level which has an oxygen pressure of about 160 mm Hg), each 100 cc contains about 20 cc of oxygen This is the maximum quantity of oxygen which the blood will absorb, and it is then said to be fully saturated with the gas Twenty volumes percent is, therefore, the oxygen capacity of normal human blood The blood does not, however, become fully saturated with oxygen in passing through the lungs Arterial blood contains only about 19 4 volumes percent of oxygen In other words, it is only 97 5 percent saturated Of the 19 4 volumes in every 100 cc of blood all but 0 24 volumes are in combination with hemoglobin This 0 24 volume percent is in simple solution in the plasma Whole


250

blood, owing to its content of hemoglobin, can absorb, therefore, 80

times more oxygen than can an equivalent quantity of plasma
Hemoglobin when saturated with oxygen, or nearly so, is called
oxyhemoglobin. The oxygen is bound very loosely to the hemoglobin,
the latter is said to be oxygenated rather than oxidized, for a true oxthe latter is said to be experiment rained intain counter, for the two dies is not formed. Hemoglobin that has given up part of its oxygen is called reduced hemoglobin. If hemoglobin formed a stable compound with oxygen—that is, if it formed an oxide which did not readily part with its oxygen—it would serve no useful purpose in respiration, for the unloading of oxygen to the tissues is quite as important as its rapid absorption by the blood in the lungs. The combination of oxygen with hemoglobin in the lungs and the dissociation of oxygen in the tissues proceed at nearly equal velocities. The quantity of oxygen which will combine or be held in combination with hemoglobin in the erythrocytes is dependent upon the partial pressure of oxygen in the plasma (i.e., the oxygen in simple solution). The pressure of oxygen in the plasma varies in turn with the pressure of oxygen in the alveolar air, on the one hand, and in the tissue fluids and cells, on the other (Henry's law, p 245) For example, in passing through the lungs the plasma comes into equilibrium with the alveolar air which has an oxygen pressure of around 100 mm Hg The hemoglobin therefore, becomes nearly saturated with oxygen In the tissue fluids the oxygen pressure is relatively low, and still lower in the cells Oxygen diffuses, therefore, from the plasma to the tissue fluids and from the latter to the cells. The lowered oxygen pressure of the plasma thus occasioned causes the liberation of oxygen from the hemoglobin. Thus, through this gradient of pressure, a steady stream of oxygen from the red cells to the tissue cells is assured.

The oxygen dissociation curve of hemoglobin The relationship The Origin dissociation curve of hemoglobin and the partial pressure of oxygen to which it is exposed can be shown by placing samples of blood in a series of cylindrical glass containers (called tonometers) and introducing air mixtures of known oxygen and carbon dioxide pressures. The glass vessels are rotated in a water bath at body temperature, the blood is thus spread in a thin film over their interiors, perature, the blood is thus spread in a thin him over their memory, and allowed to come into gaseous equilibrium with the atmosphere to which it is exposed. The percentage oxygen saturation of each of the blood samples is then determined and plotted as shown in Figure 6.19. Percentage oxygen saturation is indicated along the upright lines (ordinates), oxygen pressures along the horizontal lines (abscissae). The curve plotted in this way is called the oxygen dissociation curve of hemoglobin. It is important to notice the peculiar shape of this curve. It tends to flatten out at oxygen pressures above 70 mm Hg. It has been described as S shaped. At 100 mm Hg, which is the oxygen pressure in the alweolar air, the hemoglobin is over 97 percent saturated. It would become fully saturated at about 160 Hg. Raising the oxygen pressure above 100 mm Hg can therefore cause little increase in the quantity of oxygen absorbed; and a fall in oxygen pressure down to 60 or 70 mm Hg causes a minor reduction in the oxygen saturation.

Fig 6 20 Showing effect of increasing pressures (tensions) of CO, upon the oxygen dissociation curve of hemoglobin (Bohr effect at constant temp. 37º C). (After Barcroft.) Percentage saturation of hemoglobin along upright lines (ordinates); oxygen pressure in mm Hg along horizontal lines (abscissae).



The slope of the lower part of the curve (i.e., at the lower oxygen pressures) is much steeper. This means that any given reduction in oxygen pressure causes the release of a relatively large quantity of oxygen from the hemoglobin. In other words, in passing through the tissues, where the oxygen pressure is low, very free dissociation of oxyhemoglobin occurs and a relatively large quantity of oxygen is unloaded for delivery to the tissue cells.

Two factors—namely, a rise in temperature and an increase in hydrogen ion concentration—cause a shift to the right of the oxygen dissociation curve of hemoglobin. That is to say, each of these factors reduces the quantity of oxygen which hemoglobin will hold at any given oxygen pressure. Therefore, during increased tissue activity (e.g., muscular contraction) the local rise in temperature and the increase in hydrogen ion concentration, resulting from carbon dioxide and lactic acid production, cause a greater load of oxygen (per unit quantity of blood) to be delivered to the tissues. The influence of carbon dioxide pressure variations upon the form of the oxygen disso-

ciation curve of hemoglobin is illustrated in Figure 6.20. This is known as the Bohr effect.

known as the Bohr effect.

Carbon dioxide transport. Arterial blood contains from 44 to 52 volumes percent of carbon dioxide, venous blood from 50 to 60 volumes percent. A small part of this—namely, about 2.5 volumes percent (or from 4 to 5 percent of the total—is in simple solution in the plasma. A further 4 or 5 volumes percent (8 to 10 percent of the total) is combined directly with hemoglobin. The compound is called carbohemoglobin The rest of the carbon dioxide is carried in the plasma as sodium bicarbonate. The chloride shift mechanism, through which the sodium bicarbonate is formed when carbon dioxide enters the the sodium bicarbonate is formed when carbon dioxide enters the blood from the tissues, has already been explained (p. 112). After a study of this mechanism it will be realized that, though only a small part of the carbon dioxide is carried in the blood actually combined with hemoglobin, the latter serves in an indurect way—that is, by giving up its alkali—for the carriage of over 85 percent of the total carbon dioxide. For this reason the quantity of carbon dioxide which can be absorbed by plasma, from which the red cells have been separated by centrifuging (so-called separated plasma), is only a small fraction of that which the plasma in whole blood ("true plasma") will about the plasma in whole blood ("true plasma") will absorb.

### The Control of Respiration

The Control of Respiration

Respiration is essentially an involuntary act—that is, it is carried out automatically and without thought Nevertheless, it is to a certain extent under voluntary control. One can, for example, cease breathing for a time, or can vary the rate or the rhythm of the respirations (as in speaking or singing). The respirations are also controlled through an effort of the will or semiautomatically (reflexty) during swallowing, coughing, sucking, etc., and are altered by emotional states. But the respiratory movements are essentially automatic in nature. The breathing cannot be suspended voluntarily for more than 45 seconds or so. At the end of this time one is compelled to take a breath.

The automatic, involuntary character of the respiratory movements is due to the rhythmical discharge of impulses from a group of nerve cells situated in the lower part of the floor of the fourth ventricle in the medulla oblongata. These constitute the respiratory center. Destruction of this region causes complete and permanent arrest of respiration. The impulses originating in the center are conducted down the spinal cord to the anterior horn cells of the third, fourth,

and fifth cervical segments The phrenic nerve which supplies the chief muscle of respiration—namely, the diaphragm—arises from these spinal segments. Impulses also descend from the center to the motor neurons in the third fourth fifth and sixth thoracic segments, supplying fibers to the intercostal muscles.




Fig 6 21 Diagram showing the nervous connections in the respiratory mechanism

The complete central nervous mechanism in the brain stem con trolling respiration does not occupy an area as circumscribed as was at one time believed. Though we still speak of the respiratory center, and stimulation of a small spot in the lower part of the medulla ob longata will cause pronounced respiratory responses, modern research has revealed that the regulation of the respiratory rhythm is de pendent upon a much more complex and extensive neural mechanism. There is a group of nerve cells or center in the medulla which when stimulated electrically during inspiration, causes the immediate arrest

254

of this movement and brings on expiration, it is called the expiratory center. An inspiratory center lies in close proximity to the expiratory center and when stimulated causes strong contractions of all the inspiratory muscles. In the living body, the expiratory center appears to discharge an intermittent stream of inhibitory impulses to the in spiratory center, interrupting its activity at regular intervals. Expiration then follows automatically and in a passive manner. In other words, expiration is the result of the cessation of impulses from the inspiratory or primary center to the diaphragm and other muscles of inspiratory and the group of nerve cells, known as the pneumotaxic center, situated in the lower part of the pons, is connected to the in spiratory center by nerve tracts. The function of this center is not altogether clear, but it is believed by most physiologists to receive excitatory impulses, discharged rhythmically from the inspiratory center which it relays to the expiratory center, the latter then exerts, intermittently, an inhibitory influence upon the inspiratory center as described above. All three centers are bilaterally represented (see Fig. 221).

The frequency of the impulses discharged along the phrenic nerves, and other nerves to the inspiratory muscles, varies rhythmically with the inspiratory movements, rising to a maximum toward the end of inspiration and falling to a minimum, or ceasing entirely during expiration

The respiratory nervous mechanism, though fundamentally automatic in its action, is influenced by *chemical* and *reflex* factors. These will be considered separately.

will be considered separately

Chemical factors. Carbon dioxide stimulates the respirations, any rise in the partial pressure of this gas in the arterial blood results in increased frequency and depth of the respirations. Under ordinary circumstances, the carbon dioxide pressure in the alveolar air, as in the arterial blood remains remarkably constant at around 40 mm Hg, for the slightest rise suffices to increase the pulmonary ventilation. The excess gas is thus removed and the pressure restored, automatically, to the normal level Professor J. S. Haldane and his associates found that an increase of only 0.2 percent in the carbon dioxide of the alveolar air (a rise in partial pressure of 1.5 mm Hg) doubled the volume of the air breathed per minute. The effect of carbon dioxide upon the respirations may be demonstrated by having a subject breathe an air mixture containing a high percentage (5 or 6 percent of carbon dioxide (Fig. 6.22). The respirations become deeper, more rapid, and even violent, though the mixture contains over 90 percent

of oxygen. Conversely, if a subject voluntarily increases his pulmonary ventilation by breathing quickly and as deeply as possible for a minute or so, a greater than ordinary quantity of carbon dioxide is "pumped" out of, or "blown" from, his blood. When the forced voluntary effort (forced breathing) ceases, spontaneous respiratory movements are not resumed for a period of from 40 to 60 seconds. This interval of suspended respiration, during which there is no desire to breathe, is called apnea (literally, no breathingsee Fig. 6.23). With longer peri-




Fig 6 22 Showing effect upon pulmonary ventilation of increasing the CO<sub>2</sub> percentage in the inspired air. (After Scott, redrawn and modified.)

ods of forced breathing, the apneic period may be extended to 2 minutes or more. If the subject repeats the experiment, but instead of breathing ordinary air, breathes an air mixture containing a higher concentration (4 to 5 percent) of carbon dioxide and a normal percentage of oxygen, apnea does not occur. The cessation of breathing following hyperventilation of the lungs is due, therefore, to the removal of earbon dioxide from the blood, and not to its being overcharged with oxygen. A reduction in the carbon dioxide pressure of the alveolar air by about 0.2 percent is sufficient to induce a short period of apnea.

Effects of a low pressure of oxygen in the arterial blood. The effect upon the respirations of oxygen want is seen at altitudes of 10,000 feet or more, where the oxygen pressure of the atmosphere is greatly reduced (p. 261). An increase in pulmonary ventilation then occurs. The respiratory stimulus, in this instance, cannot be carbon dioxide.



Fig 6 23 Forced breathing for a period of 2 minutes is followed by apnea lasting for 3 minutes. This is succeeded by periodic breathing of the Cheyne-Stokes type (p 259 and Fig. 6 25), which persists for a minute or so before normal breathing is resumed (After Douglas and Haldane)

for the pressure of this gas in the alveolar air at high altitudes may be only 30 mm Hg, instead of the usual 40 mm. The first return of the respirations at the end of a long apneic period caused by forced breathing is also due to oxygen want, for at this time the partial pressure of carbon dioxide in the blood is still below the normal level. For this reason, forced breathing of an air mixture containing a high




Fig 6.24 Diagram summarizing the nervous factors in the control of respiration (Modified from Best and Taylor, The Physiological Basis of Medical Practice.)

percentage of oxygen results in an extraordinarily long period of apnea (7 to 8 minutes), the large store of oxygen held in the lungs postponing the onset of oxygen want.

postponing the onset of oxygen want.

The stimulating effect which oxygen want exerts upon the respiratory center is now believed to be an indirect one—namely, through the stimulation of the chemoreceptors in the carotid body (p. 258). The direct effect of anoxia upon the center itself is of a depressing rather than of a stimulating nature. Under the ordinary circumstances of health, however, oxygen want is not even an indirect factor in the control of respiration, for it is not until the oxygen percentage in the atmosphere reaches a very low level (12 to 14 percent) that any noticeable effect upon respiration is produced. When one holds the

breath, for example, the uncontrollable desire to breathe is due to the accumulation of carbon dioxide, and not to lack of oxygen. The intense urge to breathe occurs before the oxygen content of the blood has fallen to the point where the respirations are stimulated (through the carotid body) by oxygen want.

Reflex and chemoreflex factors. Afferent impulses originating in almost any part of the body may influence the activity of the respiratory center. The sharp inspiration which follows a painful stimulus, the inhibition of inspiration during swallowing (p. 301) or when sensory endings in the nasal mucosa are stimulated (as by some pungent odor), and the increased rate and depth of breathing caused by the stimulation of cutaneous nerves by a cold or a hot bath are a few familiar examples illustrating the effects of afferent nerve stimulation upon the respiratory movements.

The activity of the respiratory center is also influenced by proprioceptive impulses (p. 514) arising in the respiratory muscles, as well as by impulses discharged from the higher (psychic) centers of the brain. Various changes in respiratory rate, depth, and rhythm may be brought about through emotional and mental factors. The spasmodic breathing of laughter, the sobbing respirations in grief, the slow shallow breathing during attention, suspense, or apprehension, and the rapid respirations in excitement or fear may be cited (Fig. 6.24).

An outstanding reflex effect upon the rate and depth of breathing is brought about through impulses originating in afferent endings of the vagus situated in the pulmonary tissue itself. The adequate stimulus for the vagal endings is stretching caused by inflation of the lungs. Thus, at the end of inspiration, distension of the air sacs causes a stream of impulses to be discharged to the respiratory center. These depress the activity of the center; that is, the frequency of the impulses discharged down the phrenic and other efferent respiratory nerves is reduced; or the efferent discharges may cease entirely. Expiration, which, as we have seen, is brought about mainly in a passive manner, therefore follows. The arrest of inspiration brought about in this way, with the consequent onset of expiration, is called the Hering-Breuer reflex. Section of the vagus nerves in experimental animals, by interrupting the afferent impulses from the lungs, results in slowing and deepening of the respirations—the inspiratory phase being then prolonged.

The carotid sinuses (p. 199) and carotid bodies under certain special circumstances, but probably not in ordinary conditions, play a role in respiratory control. The walls of the sinus contain receptors

which, when stimulated mechanically (pressure or stretch), discharge impulses to the respiratory center as well as to the cardio-inhibitory and vasomotor centers. Thus, distension of the sinus by a rise in blood pressure causes slowing of the respirations, or even complete respiratory arrest (apnea) for a short time. A fall in blood pressure has the reverse effect—namely, increased pulmonary ventilation. These effects come into play only under very unusual and special circumstances.

The carotid body is a small structure attached to a twig of a small branch of the external carotid artery close to the carotid sinus. It contains cells which have the specialized function of responding to chemical stimuli. These chemoreceptors, as they are called, are stimulated by a rise in the carbon dioxide tension of the arterial blood, by an increase in hydrogen ion concentration, or by oxygen lack. The nerve impulses set up by the stimulation of these chemoreceptors and discharged along the sinus nerve to the respiratory center result in an increase in the rate and depth of breathing. A rise in the carbon dioxide pressure exerts a similar, though less pronounced, effect. The carotid body plays no role, or a very minor one, in the control of the circulation (see p. 2011).

As mentioned on page 257, it is only when oxygen want reaches an extreme degree that the chemoreceptors of the carotid body are stimulated. The same applies to carbon dioxide excess. The sinus mechanism, therefore, plays no part in the control of respiration under ordinary physiological conditions.

# Physiological Modifications of Respiration

Laughing. Laughing consists of a deep inspiration followed by a series of short spasmodic expirations. The characteristic sound is caused by the vibrations of the vocal cords which are held tense in the path of the outgoing air. In crying, the respiratory movements are very similar.

Coughing. This is a reflex act usually caused by stimulation of sensory endings of the vagus nerve in the mucosa of the larynx or trachea, but it may result from the excitation of afferent vagal endings in the lungs or pleura. The act comprises a short inspiration, followed immediately by closure of the opening of the larynx (glottis),

and a forcible expiratory effort A high pressure is thus created within the lungs and lower air passages. The glottis then opens suddenly, allowing the air to escape in a blast. Thus, any irritating material which may be present in the laryax is expelled.

Sneezing. Like coughing, sneezing consists of a short inspiration followed by a forcible expiration, but the glotts remains open During the first part of the expiratory effort the way into the mouth is blocked by the elevation of the tongue against the soft palate, the blast of air being thus directed through the nose Later, the resistance offered by the tongue is, as a rule, removed, the air then escaping through the mouth. The sneezing reflex is initiated by irritation of the nasal endings of the trigeminal nerve


Yawning. Usually an indication of fatigue, sleepiness, or simply of boredom, yawning may also be induced by seeing someone else yawn. It is primarily of psychic origin. A yawn comprises a deep inspiration with the mouth fully open. Sighing consists of a prolonged expiration.

Hyperpnea. This term is applied to any increase in pulmonary ventilation, due to an increase in either in the rate or in the depth of breathing, or in both

# **Abnormal Types of Respiration**

Periodic breathing. This term is given to certain types of abnormal respiratory rhythm (see Fig 6 25) The best known of these is called Cheyne-Stokes breathing, after two physicians of the last century who described it as a characteristic feature of certain diseases. This type of breathing is marked by periods of rapid and deep respiration (hy-

Fig 625 Cheyne Stokes breathing f, hyperpheic periods, 2, apneic periods (After Waller)



perpneic periods) alternating with intervals of complete cessation of respiration (apneic periods). At the beginning of the hyperpneic periods, the respirations are slow and shallow, but they quickly increase to a maximum rate and depth, and then, becoming smaller and smaller again, cease. The apneic periods are due to the excessive elimination of carbon dioxide from the blood during the preceding periods of hyperpnea. The latter are due, in turn, to oxygen lack,

acting through the carotid body reflex, and the accumulation of carbon dioxide during the apnete periods. The respiratory center, apparently, has lost some stabilizing factor which normally prevents if from responding too vigorously to changes in the gaseous composition of the blood. Like a machine without a flywheel, the action of the center consists of uneven, relatively small changes in gas pressures causing a too sudden and unduly errait respiratory effect.

causing a too sudden and unduly great respiratory effect
Cheyne Stokes breathing is seen in advanced kidney disease, in
cases of raised intracranial pressure, and in poisoning by narcotices
In these conditions it is a symptom of very grave portent But periodic
breathing of a similar type occurs under certain physiological con
ditions—e.g., in animals during hibernation and sometimes in children during sleep. It is also seen when the respirations are returning
after the period of apnea induced by forced breathing, or it may occur
at high altitudes

Dyspnea. Difficult breathing, or dyspnea, is seen in a number of diseased conditions. The breathlessness in heart disease or in acidosis and the respiratory distress of asthma, pneumonia, and other pul monary conditions, and at high altitudes are examples. In acidosis the respirations are increased in an effort to maintain the HisCO<sub>2</sub>.

ratio at its normal value by reducing the numerator of the equation—that is, by "blowing off carbon dioxide from the blood (p 255) The dyspines of heart disease, anemia, and pneumonia, and at high altitudes is caused by an inadequate supply of oxygen to the tissues, in cluding the carotid body which is stimulated thereby

A rise in metabolism, from whatever cause, brings about an in-

A rise in metabolism, from whatever cause, brings about an increased production of carbon dioxide Muscular exercise increases the metabolism and, therefore, stimulates the respirations, and even a healthy person suffers from dyspine after a bout of strenuous muscular effort when he puffs, gasps, or pants and feels that he will suffocate if he does not get more air into his lungs. If, as in hyper-thyroidism (p. 411), the resting metabolism is already above normal, dyspine as experienced with a grade of muscular exertion which would not cause any respiratory distress in a normal person. In anemia, also, breathlessness occurs upon any slight muscular exertion, because the hemoglobin concentration of the patient's blood, though capable of carrying the required quantity of oxygen to the tissues when the

•

Whenever breathing is distressing or a person is unduly aware of the respiratory act the term dyspace may be applied

body is at rest, is unable to satisfy the demands of even mild physical effort.

The degree of hyperpnea at which respiratory distress is experienced varies widely in different individuals according to their physical state. As compared with untrained persons, athletes are able to increase their pulmonary ventilation to a much greater extent before any difficulty in breathing (i.e., dyspnea) is experienced.

# Oxygen Want-Anoxia or Hypoxia

Any failure of the tissues to secure a supply of oxygen sufficient for their needs is referred to as oxygen want or anoxia Anoxia is classified according to its cause into the following four groups: anoxic, stagmant, anemic, and histotoxic.

Anoxic anoxia. This term applies to any type of anoxia in which the blood in its passage through the lungs does not reach normal saturation (about 97.5 percent), whether as a result of a low oxygen tension in the atmosphere as at high altitudes and in vitiated atmospheres (e.g., in mines), or to some interference with breathing (e.g., obstruction of the respiratory passages, asthma, pneumonia, etc.).

Mountain sickness. The unpleasant and sometimes serious effects of the anoxia of high altitudes are referred to as mountain sickness. The first signs of mountain sickness appear at an altitude of between 8000 and 10,000 feet above sea level. At this height the percentage oxygen saturation of the hemoglobin is considerably below the normal level. For example, at a height of 14,000 feet above sea level, the atmospheric pressure is around 450 mm Hg. The partial pressure of oxygen, being reduced proportionately, amounts to  $\left(\frac{20.96}{100} \times 450 = \right)$  94 mm Hg, as compared with 160 mm at sea level. The oxygen pressure in the alveolar air and arterial blood is, therefore, only from 55 to 60 mm Hg. The oxygen saturation of the hemoglobin is around 82 percent as compared with the normal of about 97.5 percent, and the symptoms and signs of anoxia are apparent. These include headache. nausea and vomiting, dyspnea, and cyanosis (p. 263). Emotional outbursts (e g, laughing or crying, quarrelsomeness or hilarity) are common There is often a sense of exhilaration, or an exhibition of foolhardiness, boisterousness, or stubbornness. If a person remains for a

<sup>&</sup>lt;sup>a</sup> Hypoxia, which means a diminished oxygen supply, is a better term than anoxia (literally no oxygen) and is, therefore, more correct etymologically. In the types of oxygen deficiency to be immediately described, the older word (anoxia) will be retained, since in this connection it has the sanction of long usage.

time at the high altitude, these effects gradually disappear, he becomes acclimatized The acclimatization process consists of a rise in the red cell count (see Fig 3 5, p 62), and adjustments in the circulatory and respiratory mechanisms. Natives of high mountains have a red cell count of from 6,500,000 to 8,500,000, depending upon the altitude at which they live. The volume of the lungs and the size of the chest is greater than the normal for a man of the same height residing at sea level. In rapid ascents to very high altitudes (40,000 to 50 000 feet), as in an airplane, consciousness is lost rapidly unless the flyer can receive oxygen inhalations from a storage cylinder, or the plane is provided with an airtight cabin within which the partial pressure of oxygen is equal to that at an altitude of 8000 feet or less. Stagnant anovia. This anoxia is due to slowing of the circulation through the tissues as occurs in heart failure. The réduced velocity of the blood in the capillaries permits the hemoglobin to give up a greater fraction of its total oxygen load than normally. The venous blood contains therefore a proportionately larger amount of reduced hemoglobin. When the blood remains long in the capillaries, though a larger quantity of oxygen is abstracted from it, the greater part of the gas is delivered at low pressure. The itsues, therefore, receive less oxygen in a given time than when the blood flow is rapid. Anemic anoxia. This type of oxygen want results from a reduced oxygen-tarrying capacity of the blood. It includes that due to carbon monoxide poisoning as well as that resulting from anemia.

\*\*Carbon monoxide poisoning Carbon monoxide combines with hemoglobin to form COHb, a relatively stable compound—that is, one which dissociates very slowly. Any hemoglobin in this form cannot take up oxygen and is therefore dispossessed of any respiratory function. Carbon monoxide causes death for this reason only. The subject poisoned by the gas is deprived of the use of part of his hemoglobin just as surely as i

As in the anoxia of mountain sickness, in that due to carbon monoxide poisoning, the subject often becomes stubborn, perverse, or unruly Though he is not unaware of his danger he may, owing to some fanciful idea, make no attempt to escape, and may even resist efforts of others to bring him into fresh air. This is another dangerous feature associated with exposure to this gas.

The most effective means of resuscitating a subject of carbon monoxide poisoning is artificial respiration and, when possible, using an air mixture of about 40 percent oxygen and 7 percent carbon dioxide. The oxygen tends to hasten the liberation of carbon monoxide from the hemoglobin while the carbon dioxide stimulates the respirations.

Histotoxic anoxia. In this type of oxygen deficiency, the oxidative processes of the tissues are depressed or abolished by a poison such as cyanide. The oxygen-carrying capacity of the blood is not affected, but, since cellular oxidations are greatly reduced or in abeyance, the usual amount of oxygen is not removed from the blood in the capillaries. The venous blood, therefore, has a high oxygen saturation, and is arterial in color. Cyanide poisoning is treated by the intravenous injection of a solution of sodium intrite followed by the injection of sodium throsulfate. However, since cyanide poisoning is so rapidly fatal, there is rarely an opportunity for applying this or any other method of resuscitation.

Cyanosis. The name cyanosis is given to the blue tint of the skin caused by an unusually high concentration of reduced hemoglobin in the blood of the capillaries composing the subpapillary venous plexus (p 35). The blueness is usually most pronounced in the skin of the lips, ears, and finger tips. It is the visible sign of hypoxia, either of the anoxic or stagnant type. That is, the blood either does not receive a full load of oxygen in the lungs or it gives up an unusually large proportion of its oxygen to the tissues. In both these instances, the capillary blood will have an abnormally high concentration of reduced hemoglobin, which is darker in color than oxyhemoglobin. Blood contains some 15 grams of hemoglobin per 100 cc. In the cutaneous capillaries of a normal person, about 2.5 grams of this is reduced hemoglobin, the remaining 12.5 grams is oxyhemoglobin. When, as a result of either of the two types of hypoxia just mentioned, the concentration of reduced hemoglobin in the capillaries of the skin of reduced hemoglobin per 100 cc, cyanosis appears. In other words, 5 grams of reduced hemoglobin per 100 cc of blood in the capillaries of the skin and mucous membranes is necessary for the production of cyanosis.

#### 264 RESPIRATION


The anemic and histotoxic types of anoxia do not cause cyanosis. In the first of these, though the hemoglobin is diminished in amount, what there is of it is as fully saturated with oxygen as in health. Furthermore, a person with a very severe anemia (hemoglobin less than 5 grams per 100 cc of blood) obviously cannot become cyanosed under any circumstances, for even though all his hemoglobin should be in the reduced form, it would not be in sufficiently high concentration to cause any discoloration of the skin. On the other hand, a person whose blood had an abnormally high concentration of hemoglobin (25 to 30 grams per 100 cc, as in polycythemia, instead of the normal of about 15 grams) becomes cyanosed with a relatively mild degree of hypoxia. In the histotoxic type, the capillary blood actually contains a lower concentration of reduced hemoglobin than in health, for, as a result of the action of the poison on the respiratory function of the cells of the tissues, the oxygen in the blood is not utilized.

Oxygen administration. In certain acute respiratory conditions accompanied by hypoxia the administration of oxygen in high concentration (50 to 100 percent) is of great value. Also, in heart disease in which the stagnant type of anoxia may exist, oxygen inhalations are of benefit. It must be remembered, however, that there is a limit to the amount of oxygen which the blood will take up from the lungs in health the blood coming from the lungs is over 97.5 percent saturated with oxygen so, increasing the oxygen in the inspired air five times does not mean that the blood will take up five times as much oxygen as when ordinary air is breathed. The most that can be effected is to increase the oxygen saturation of the blood to 100 percent. In the stagnant type of anoxia when the arterial blood is at the normal of about 97.5 percent, this would mean an increase of only 2.5 percent or so Nevertheless, this extra 2.5 percent in heart disease is often highly salutary.

### The Voice

The mode of production of the human voice—with its varied tones, its range of pitch, and its volume—has long aroused the interest of physiologists. As an instrument of sound, the voice box or larynx, with the cavities of the mouth, throat, trachea, and lungs, may be compared to the pipe of an organ. A reed and the column of air in the organ pipe are set into vibration by an air blast. In a somewhat similar manner the vocal cords within the larynx are thrown into vibration by air expelled from the lungs. The nose, mouth, throat, and chest

Fig 6 26 The larynx and upper rings of the trachea. The hyoid is a small bone lying near the root of the tongue, The thyroid cartilage forms the front and side walls of the larynx. The cricoid is a circle of cartilage shaped like a signet ring, lying between the larynx above and the trachea below.



serve as resonating chambers. Certain notes can be played by the organist which imitate the human voice in a truly remarkable way.

The larynx (Fig 6 26) lies at the upper end of the trachea, its walls being formed of cartilage and lined with mucous membrane; the vocal cords are two thin-edged bands or membranes lying within it. The cords run from before backward, being attached behind to two small cartilagnous bodies (the arytenoid cartilages) and fixed in front to the wall of the larynx. By the contraction of small muscles attached to




Fig. 6 27 The vocal cords viewed from above (diagrammatic) in different degrees of approximation A, in midposition, B, in full addiction; C, in full abduction.

them, the arytenoid cartilages can be rotated when necessary By this means the vocal cords can be swung away from one another—that is, outward against the walls of the larynx (Figs 6 27 and 6 28), or in ward toward one another until only a small chink remains between them. At ordinary times the cords he against the wall of the larynx, the gap separating them 4s wide, and no sound is produced during the pas sage of the breath During speech they are brought toward one another and into the current of air expelled from the lungs, which, being un



Fig 6 28 The interior of the larynx as seen with the laryngoscope

able to escape except through the narrow opening, sets the edges of the cords into vibration

Sound possesses three properties—loudness quality or timbre, and pitch. The loudness of the voice depends upon the energy with which the vocal cords vibrate, the greater the pressure under which the air. is expired, and the greater the movements made by the cords, the louder will be the sound The pitch is determined, primarily, by the length and tightness of the cords, and, therefore, by the frequency of their vibrations. In children and women the cords are short, and the their vibrations In children and women the cords are short, and the voice is high pitched In men the cords are longer, and the voice is deeper All of us can adjust the tension and to some extent the length of our vocal cords and so alter the pitch of the voice, but trained sing ers have developed this ability to the greatest degree. The quality or timbre of our voices depends on the number and intensity of the overtones or harmonics which are produced, and these in turn depend upon the shape and capacity of the resonating chambers—the mouth, the trachea, and the chest. The training of the voice consists very largely in modification of the mouth and throat cavities so that the sound produced in the largery receives the greater possible numbers of these harduced in the larynx receives the greatest possible number of these har-monies or supplementary tones

In speech the musical sounds produced by the vibration of the vo-




Fig 6 29 Showing the positions of the lips and tongue and the shape of the mouth cavity when pronouncing a 1 and oo

cal cords are modified by the numerous changes which may be made in the size and shape of the air passages—the pharynx and the mouth The vowel sounds a e, i, o u, are formed in the lower air passages and molded as it were by the mouth the latter cavity assuming positions which are characteristic for each vowel (Fig 629) When we whisper, these sounds may be produced simply by placing the mouth in the required position—that is, without vibration of the vocal cords. The consonants are formed by interrupting to different degrees the expired air in various parts of the vocal pathway

# The Physiology of the Kidney. Micturition

The structure of the kidney. The kidney is composed of a large number of microscopical structures called nephrons. They are the excreting units of the kidney. Their function is the separation from the blood of water and certain materials (urea, salts, etc.) in solution which are then excreted as urine. In man, there are some 2,000,000 nephrons-1,000,000 in each kidney.

The nephron comprises (1) a spherical, vascular structure called the renal glomerulus or Malpighian corpuscle, and (2) the renal tu-

bule (Figs 7.1, 7 2, 7.4, and 7.5).

The renal glomerulus consists of some fifty separate capillaries bent into short loops to form a compact mass called the glomerular tuft. The upper end of each renal tubule is expanded into a structure known as Bowman's capsule. The cavity of Bowman's capsule is crescentic in cross section, owing to the encroachment of the capillary tuft which is, as it were, pushed into one side of the capsule as a ball might be pressed into the side of a partially inflated balloon. The walls of the capsule are formed of a thin membrane composed of a single layer of flat cells which, therefore, serves also as a covering for the glomerulus. This membrane and the equally thin walls of the capillaries alone separate the blood from the cavity of Bowman's capsule.

The renal tubule drains Bowman's capsule and, for purposes of description, can be divided into three parts. The first third or so is tortuous, and lies in close relation to the glomerulus; it is known as the proximal convoluted tubule. The walls of this portion of the tubule are composed of a single layer of columnar epithelial cells. The free




Fig. 7.1 Showing a renal unit or nephron of the cortex of the kidney, with its blood supply, and a collecting tubule. The thin part of Henle's loop varies in length; it often extends into the ascending limb of the loop or may even be confined to this limb.

270




Fig 72 Diagram of a glomerulus of the kidney Note that capillanes pass from the afferent to the efferent arteriole in separate loops There are many more capil lary loops than are shown here each glomerulus contains 50 or more

borders of the cells show fine perpendicular strations which give them a brushlike appearance. This so called brush border is not seen else where in the tubule, though the cells lining the small intestine have a similar appearance (Fig. 7.3). After forming the proximal convoluted tubule, the nephron straightens out and descends for a distance through the substance of the kidney, then turning upon itself, it ascends again to the region of the glomerulus. The hairpinlike bend, so formed, is called Henle's loop the straight parts of the tubule leading to and



Fig 73 Cross section of a con voluted tubule, highly magnified.

away from the bend are called, respectively, the descending and as cending limbs of Henle's loop The tubule composing the loop for half or more of its length is much nar rower than either the proximal or distal convoluted tubule, it is usually referred to as the thin portion of Henle's loop (Fig. 7.1) The ascend ing limb passes into a second tortuous section lying in close relation to the glomerulus, this is called the distal convoluted tubule The distal convoluted tubules empty the tubular fluid (now completely formed urine) into larger channels, called collect




Fig 7.4 Drawn from a photograph of portion of a kidney after injecting its vessels with fluid that sol dified and then destroying the renal tissue with acid to reveal the glomeruli



Fig. 7.5 Microscopical appearance of normal renal tis sue G glomerulus T tubule

ing tubules. These latter tubes serve merely to conduct the urine to the pelvis of the kidney (p. 285 and Fig. 7.7), playing little or no part in the actual formation of the urine.

The blood and nerve supply of the kidney. Blood is delivered to the kidney by the renal artery, a short thick vessel which arises from the





Fig 7.6 Diagram of renal circulation. Veins, black; arteries, white; hatching indicates conversion of arterial to venous blood.

abdominal aorta. All the blood which enters the kidney, except a small fraction for the nourishment of the covering (capsule) and connective tissue of the organ, passes through the glomeruli The capillaries of each glomerular tuft are derived from a single arteriole called the afferent vessel. Converging again, the capillaries form a single effluent channel, known as the efferent vessel (Fig. 7.6). This is also of arteriolar construction. There is considerable disparity in size between the diameters of the two glomerular arterioles, the efferent vessel being the smaller. This arrangement makes for a relatively high pressure within the capillary loops. Furthermore, variations in glo-

merular capillary pressure can be brought about through changes in the relative diameters of the two vessels. Dilatation of the afferent arteriole or constriction of the efferent, the caliber of the other vessel in each instance remaining the same, causes a rise in the blood pressure in the capillaries of the renal glomerulus. Constriction of the afferent or dilatation of the efferent vessel will produce the reverse effect.

The blood, after traversing the glomerulus, is distributed to the rest of the nephron through branches of the efferent vessels which terminate in a capillary network surrounding the tubules.

Fig 7.7 Interior of the kidney sectioned lengthwise Urine enters pelvis of the kidney through openings in the tips of the pyramids in the medulla



The nerves of the kidney are derived from the vagus and splanchnic (sympathetic) nerves. These nerves have no secretory action, for the kidney continues to perform its functions in a normal manner after it has been completely denervated.

The renal nerves act solely upon the renal vessels controlling the volume of blood entering the organ It is probable that they also cause changes in the distribution of blood within the kidney itself, diverting blood from one part of the renal tissue to another. It has been found within recent years that there are actually two pathways which the blood may take in its passage through the kidney—namely, through the glomeruli in the superficial part of the renal cortex, or through other glomeruli situated more deeply in the cortex near the renal medulia (Figs. 7 6 and 7.7). For example, stimulation of the central end of a cut sciatic nerve in an animal such as the rabbit causes the blood to "by-pass" the first-mentioned vessels, which is the course taken by the blood under ordinary circumstances, and to travel through

the more deeply placed glomerul: A renal vascular reflex of this sort probably occurs in surgical shock due to severe tissue injury

The composition of the urine. The chief urinary constituents in grams per liter of urine are given in the following table

Urea is derived mainly from the breakdown in the body of food pro tein it therefore, varies considerably with the quantity of protein in the diet. On a low protein intake, it may be less than a sixth of the

Table 7-1

### Urinary constituents I Inorganic constituents

	Grams per liter
Chloride expressed as NaCl	9 Ó 🗸
Phosphorus expressed as P2O3	20
Total sulfur expressed as SO <sub>3</sub>	15
Sodium expressed as Na O	40
Potassium expressed as k O	20
Calcium expressed as CaO	02
Magnesium expressed as MgO	02
fron	0.003

II Nitrogenous constitu	ents
Urea	
25 0 grams containing approximately	10 0 grams nitrogen
Ammonia	
0 6 gram containing approximately	0 4 gram nitrogen
Une acid	
0.6 gram containing approximately	0 2 gram nitrogen
Creatinine	
1 5 gram containing approximately	0 5 gram nitrogen
Undetermined nitrogen	0 6 gram
Total	11 7 gram nitrogen

value given in Table 7 1 The creatinine, on the other hand, is derived mainly from the disintegration of body tissue, its excretion is, therefore, almost uninfluenced by the protein level of the diet

The uric acid is derived from purines (p 366) taken in the food, as well as from the purines of body tissue Ammonia is formed in the kidney itself from glutamine carried to it in the blood. The color of the urine is due to a pigment called urochrome. Normal urine does not contain glucose, except occasionally after meals, minute amounts may then appear

The reaction and specific gravity of the urine. Under ordinary circumstances, the urine is definitely acid. The average pH is around 6.0, but the reaction varies considerably with the nature of the diet. Most fruits reduce the acid its oxidized, leaving the alkali, which is excreted. Starvation increases the urinary acidity, because sulfuric and phosphoric acids are formed from the breakdown of body protein, and acetoacetic and  $\beta$ -hydroxybutyric acids from the metabolism of body fat. A high protein diet also increases the quantity of acid excretion. Herbivorous animals, on the other hand, excrete an alkaline urine except when fasting, the urine then becoming acid In diseases associated with the production of large amounts of organic acids in the body—e g., acetoacetic and  $\beta$ -hydroxybutyric acids in diabetes—the urine is more strongly acid than normally.

The normal acidity of the urine is a manifestation of one of the most important functions performed by the kidney—namely, to aid in maintaining the alkalinity of the blood and body fluids. Blood coming to the kidney has a pH of around 7.4; the urine, as just mentioned, has an average pH of 6.0. The kidney separates acid from basic substances, retaining the latter within the body and excreting the former. This conservation of the body's base is brought about in part by the conversion in the kidney of the alkaline phosphate of the blood (Na<sub>2</sub>HPO<sub>4</sub>) into the acid phosphate (NaHPO<sub>4</sub>). For example, the concentrations of the alkaline and acid phosphates in plasma are as 4 to 1, whereas, in urine they are as 1 to 9. Another way in which the kidney conserves fixed base (chiefly sodium) to the body is by the formation of ammonia. The ammonia then combines with acids brought to the kidney, which excretes them as ammonium salts.

The specific gravity of the urine of adults ranges under ordinary circumstances between 1.020 and 1.032; it is much lower in infants. In health, the specific gravity swings readily in one or other direction, in response to variations in the intake of fluid. For example, when a normal person drinks a large quantity of water, the urine which he passes within the next couple of hours is very dilute; the specific gravity may be as low as 1.001 or 1.002. On the other hand, after abstaining from fluids for some time, the specific gravity of the urine rises to 1.030 or higher. In the later stages of kidney disease, the specific gravity of the urine is low under all circumstances. It remains almost fixed around a value of 1.010, in spite of wide variations in the fluid intake.

The volume of the urine. In man, from 1000 to 1500 cc of urine are formed in 24 hours. The volume varies, of course, with the amount of

276

fluid drunk and with that lost through other channels—skin, lungs, and bowels. It is reduced, therefore, in hot weather or as a result of diarrhea or vomiting. The products of protein metabolism, especially urea, increase urine production; the volume of the urine is greater, therefore, upon a high than upon a low protein diet. Urine formation is reduced during sleep, but the total excretion of urinary solids (e.g., urea, uric acids, phosphate, etc.) is about the same during the night as during a day period of equal length, the kidney merely excreting less water—that is, more water is reabsorbed from the tubules. The urine passed after rising in the morning is therefore more concentrated than ordinary day urine. A decided rise in the volume of the night urine, as a persistent phenomenon, is called nycturia. It is an early sign of chronic renal disease. Muscular exercise, especially of a strenuous nature, reduces the volume of the urine, as a result of the liberation of the antidiuretic hormone of the pituitary body as described on page 281

The formation of urine. Though the renal corpuscle was described by the Italian anatomist, Marcello Malpighi, in the seventeenth century (1666), it was not until nearly 200 years later (1842) that the English anatomist and surgeon, Sir William Bowman, recognized the functional significance of this structure, and of the capsule which is known by his name. From his studies of the structure of the glomerulus, Bowman concluded that it served simply as a filter to remove water and salts from the plasma Though others have contended that the layer of Bowman's capsule covering the capillaries of the glomerular tuit actually secreted these urinary constituents, the former view is now universally accepted. The glomerular membrane—that is, the walls of the capillaries and the layer of Bowman's capsule in contact with them—acts in a purely passive manner, its cells, unlike those of a true gland, possessing no secretory function.

Filtration through the glomerulus. Let us now consider the filtration process in greater detail. Filtration through a membrane may be
defined as the separation, by gravity or pressure, of liquid and such
materials as can pass through the membrane from other materials to
which it is impervious. Water, salts, and other substances of relatively small molecular size pass through the capillary walls of the glomerulus into Bowman's capsule, but the capillary membrane offers an
impassable barsier to the blood cells and the plasma colloids (proteins). If there were no proteins or other colloids of comparable molecular size in the blood plasma, the force driving water and salts from
the capillaries of the tuft into the cavity of Bowman's capsule would

be equal to about 65 mm Hg—that is, the difference between the capillary blood pressure (70 mm Hg or so) and the pressure in Bowman's capsule (about 5 mm Hg). But the plasma proteins, since the glomerular membrane is impermeable to them, exert an osmotic pressure (see p. 64) which opposes the blood pressure This force acts to hold water and dissolved substances within the vessels; it, and the pressure in Bowman's capsule, must be overcome by the hydraulic force of the blood in the glomerular capillaries. The capillary blood pressure must be higher, therefore, than the osmotic pressure of the




Fig 78 Diagram to illustrate the forces responsible for the excretion of water and dissolved substances from the blood flowing through the glomerular capillaries, and the reabsorption of threshold substances from the tubules BP, blood pressure, OP, comotic pressure, FP, filtration pressure, CP, capsular pressure. All pressures in mm of mercury.

plasma colloids (proteins) and the capsular pressure combined, in order for filtration to occur. The osmotic pressure of the plasma proteins is around 30 mm Hg. Therefore, the actual driving force, or effective filtration pressure, in millimeters of mercury is as follows (see also Fig. 7.8):

### THE PHYSIOLOGY OF THE KIDNEY, MICTURITION

70	-	30	 5	=	45
Blood		Osmotic pressure	Pressure in capsule		Effective filtration

278

The filtration theory of glomerular function just outlined has been confirmed in every detail by experiment. For example, a rise in blood pressure within the glomerular capillaries, without any change occurring in osmotic or capsular pressure, causes an increase in the amount of filtrate formed. Lowering the osmotic pressure while the blood pressure is kept constant, as by the intravenous injection of a saline solution (which, of course, dilutes the proteins), also increases the flow of urine. On the other hand, raising the pressure in Bowman's capsule by blocking the ureter reduces the volume of the filtrate and, when the capsular pressure just about equals the amount by which the




Fig. 7.9 Illustration Richards' method of obtaining a sample of glomerular filtrate. When the bulb of mercury is raised, air is expelled from the tubing; upon lowering the bulb to the position indicated by the dotted outline, fluid is sucked from Bowman's capsule.

blood pressure exceeds the plasma osmotic pressure (which would be 40 mm Hg in the example given above), urine formation ceases.

A crucial experiment was performed by Dr. Richards of the University of Pennsylvania which has finally dispelled all doubt that the glomerulus acts as a filter. He obtained a sample of fluid as it was produced by the glomerulus of the kidney of a living frog, by inserting a fine glass pipette into Bowman's capsule and sucking off the filtrate by means of a flask of mercury and a length of tubing as depicted in Figure 7.9. This fluid was found to contain glucose, chloride, and all the other constituents of plasma except the proteins. It was alkaline in reaction. In other words, it was a filtrate (or ultrafiltrate)—simply deproteinized plasma.

It is scarcely conceivable that the fluid of Bowman's capsule would be of exactly the same composition as that of plasma if it were produced by a secretory process.

The concentrating function of the tubules. Normal urine is much more concentrated than the filtrate in Bowman's capsule—that is, it contains a higher percentage of total solids; it is hypertonic with respect to the plasma, and has a much smaller volume. The quantity of

Table 7-2
(After Cushney)

	Blood plasma	Urine	Times increase in concentration in the renal tubules
Water . Proteins, fats, and other colloids . Glucose . Na . Cl . Urea . Urica acid . K	percent 90–93 7–9 0 1 0 30 0 37 0 03 0 004 0 020 0 001 (?) 0 008 0 0025	percent 95  0 35 0 6 2 0 05 0 15 0 04 0 015	little or none 2 60 12 7 40 2
PO, SO, Creatmine	0 0023 0 009 0 002 0 001 (?)	0 006 0 15 0 18 0 075	2 16 90 75

filtrate formed by the human kidneys in 24 hours amounts to from 75 to 150 liters, whereas the quantity of urine secreted in the same time is only from 1 to 1½ liters. Obviously, a large volume of water—from 98 to 99 percent of that filtered through the glomerulus—must have been reabsorbed from the tubules into the blood stream Furthermore, glucose is present in the filtrate, but is absent, in health, from the urine. Glucose, then, must be reabsorbed; in the frog sodium chloride must also be returned to the blood, for Richards, in the experiment upon the frog's kidney mentioned above, found it in the filtrate but not in the urine. Materials such as urea, phosphates, and uric acid are reabsorbed to only a small extent, while other waste products, such as creatinine and sulfates, are reabsorbed in traces or not at all. Glucose and the essential salts, since they do not escape into the urine in considerable quantities unless their concentrations in the plasma are abnormally high, are called high-threshold substances. Those substances which are reabsorbed to a minor extent (urea, phosphate, and uric acid) are called low-threshold substances, while those (creatmine and sulfates) which are not absorbed at all are called nonthreshold substances (see Table 7-2).

Glucose, sodium chloride, vitamin C, amino acids, and probably other essential materials are reabsorbed from the filtrate in the protimal convoluted tubule. This part of the nephron is also responsible for from 80 to 90 percent of the water reabsorbed from the filtrate. The remaining 10 to 20 percent of the reabsorbed water is returned to the renal blood from more distal portions of the tubules—probably the thin part of Henle's loop. Little or no concentration of the filtrate occurs in the proximal tubule, the water reabsorbed containing solids in about the same concentration as does the filtrate. The concentrating function is effected by the reabsorption of water in the more distal portions of the nephrons.

The change in reaction of the urine, from alkaline to acid, occurs in the distal convoluted tubule.

As already stated (p. 272), the urine after leaving the distal convoluted tubule and entering the collecting tubules undergoes little or no further change

The process of filtration across the glomerular membrane (capillary walls and covering layer of Bowman's capsule) is purely passive, in so far as the membrane itself us concerned. That is to say, this membrane interposed between the capillary blood and the cavity of Bowman's capsule plays no active part in the separation of the water, salts, and other materials present in true solution in the plasma from the

281

plasma proteins. It exhibits no selective action; no energy is expended; no oxygen is consumed in the process. On the contrary, the cells of the tubules exercise a selective action; some constituents of the glomerular filtrate are reabsorbed; others are discarded (excreted in the urine); energy is expended by these cells in the reabsorption process: oxygen is consumed. The process may be looked upon as a reverse secretion, selected materials being passed back into the blood rather than as in the secretion of an ordinary gland which, after separating certain constituents from the blood by a specific action of its cells, discharges them into the alveoli of the gland.1

The tubules, in some species at any rate, also have an ordinary secretory function, adding urea, uric acid, and other substances to the filtrate

The action of hormones upon renal function. In health, the proportion of the filtered water which is reabsorbed by the proximal convoluted tubule is fixed at a value somewhere between the values already given-namely, 80 to 90 percent. That is to say, if the volume of the filtrate is 100 liters and the total volume of water reabsorbed is 98 liters, then 85 percent of this, or about 83 liters, is absorbed by the proximal convoluted tubule. On the other hand, the amount of water reabsorbed by the more distal part of the tubule varies very widely under different physiological and pathological conditions. It is solely through this variability that the urine is made less or more concentrated with corresponding changes in its volume. The foremost and most potent factor in altering the amount of water reabsorbed by the tubules distal to their proximal convoluted portions is the antidiuretic hormone of the pituitary body (see Chap. 10). This principle increases the reabsorption of water and reduces the reabsorption of sodium chloride; when it is lacking, reabsorption of water is reduced and the reabsorption of sodium chloride increased. In the absence of this hormone, the urine is, therefore, of large volume and very dilute (see diabetes insipidus, p. 444). The antidiuretic hormone appears to exert its effect upon the thin section of Henle's loop. One reason for believing in the truth of this statement is the fact that in animals such as frogs, whose kidneys do not have this segment, pituitary extract (containing the antidiurctic hormone) exerts no renal effect. The kidneys

<sup>1</sup> The cells of the tubules in reabsorbing water from the filtrate expend a progressively greater amount of energy as the concentration of this fluid increases and, in gessively greater and an advanced services are said to perform osmotic work.

When the osmotic pressure reaches a point at which it exceeds the power of the cells not perform the required osmotic work, the reabsorption of water ceases, the urine having attained its maximal concentration.

of such animals have no concentrating power, their urines are actually

hypotonic with respect to the plasma

The hormones of the adrenal cortex (p 426) also exert an effect
upon the excretion of water and sodium chloride by the kidney They
increase the reabsorption of sodium and reduce the reabsorption of increase the reabsorption of sodium and reduce the reassorption of potassium. Therefore, when there is insufficiency of the adrenal cortex, the reabsorption of sodium is reduced and that of potassium is increased, with consequent loss of the former from, and undue retention of the latter in, the tissues and fluids of the body (see Addison's disease, Chap 10)

Diuresis. Increased flow of urine is called diuresis, and drugs which cause such an effect are known as durettes. This group of drugs in cludes caffeine (of coffee and tea), theobromine (of cocoa), theophylline (of tea), urea, and various salts (e g, potassium nitrate, po tassium citrate, ammonium chloride, sodium sulfate, and certain mercurv compounds)

It is evident from the outline of the mechanism of renal excretion It is evident from the outline of the mechanism of renal excretion just given that there are two ways in which a greater flow of urine could be produced—namely, by increasing glomerular filtration or by diminishing tubular reabsorption. Caffeine, theobromine, and theophylline cause more capillary loops to open up and, thus to increase the total filtering surface of the kidney. Sodium sulfate and the mercurial diureties diminish reabsorption—the former by its osmotic effect after its passage into the renal tubules, the latter by its depressing action upon the readsorptive function of the cells of the tubules. Caffeine also exerts its diuretic effect by reducing reabsorption as well as by causing renal vasodilatation

The renal response to water drinking—water diuresis. The drinking of a large quantity (1 to 3 pints) of water is followed within 60 min utes or so by the passage of a large volume of very dilute urine (specific gravity around 1001) The renal effect lasts for 2 or 3 hours The resemblance of this response, as well as the very dilute nature of the urine, to the characteristic renal features of diabetes insipidus (due to urine, to the characteristic renal features of diabetes insipidus (due to the lack of antidurette principle of the pituitary body) has long been recognized. The suspicion grew that in the mechanism of water dure sis the antidurette hormone of the pituitary was in some way involved. The results of recent experiments have shown how well founded was this suspicion. Without going into details, the course of events will be briefly described. The ingested water after absorption from the intestine causes, through dilution, a slight fall in the osmotic pressure of the blood. The blood dilution exerts an effect upon certain nerve cells of the brain (their exact location is unknown) which are extremely sensitive to any change in the osmotic pressure of the blood with which they are supplied. The nerve cells—a type of receptor organ (p. 514)—are called osmoreceptors. In response to a lowering of the osmotic pressure of the plasma, the osmoreceptors exert an influence, either nervous or hormonal in nature, upon the cells of the posterior lobe of the hypophysis which causes a reduction or suppression of its anti-diuretic hormone. This hormone, under ordinary circumstances, is continually being secreted into the circulation and exerting its characteristic effect upon the formation of urine. The withdrawal of the antidiuretic action, therefore, is followed after a short interval by an increased production of urine, which is of very low specific gravity.

During a period of water deprivation, the kidney conserves its water stores. The volume of urine is reduced due to the liberation of an excess of the antidiuretic hormone. Again, the effect upon the pituitary is brought about by a change in the osmotic pressure of the blood, but now the effect is in the opposite direction for the deprivation of water has caused some slight increase in blood concentration.

The reduction in urine volume caused by the injection of a small quantity (0 001 cc) of pituitary extract into the internal carotid artery can be duplicated exactly by 10 cc of a 2.5 percent (hypertonic) solution of sodium chloride or of sugar, introduced in the same manner into the cerebral circulation; such hypertonic solutions cause a rise in the osmotic pressure of the plasma of sufficient magnitude to affect the osmoreceptors.

Summary of the chief functions of the kidney. 1. The kidney maintains the normal composition of the plasma through the elimination of excess water and the waste products of protein and purine metabolism (e.g., urea, creatinine, sulfates, uric acid, etc.), retaining at the same time the essential constituents of the blood (e.g., cells, proteins, glucose, and inorganic salts).

2. The kidney plays a prominent role in the regulation of the acid base balance of the body, through (a) the production of ammonia which, being then excreted in combination with fixed acids (ammonium salts), conserves the body's store of alkali, and (b) through the conversion of alkaline phosphate to the acid salt (Na<sub>2</sub>HPO<sub>4</sub> to NaH<sub>2</sub>PO<sub>4</sub>).

Impairment of renal function. In nephritis, the commonest type of kidney disease, a proportion of the renal glomeruli are the seat of an

acute or chronic inflammatory process. Some or all of the capillary loops in the diseased glomeruli are destroyed and, as a consequence, the filtering surface of the kidney is reduced. When the glomerular destruction is extensive, serious impairment of renal function results. The reader will recall that the tubules are supplied with blood which has previously traversed the glomerular capillaries. It follows, therefore, that glomerular damage must inevitably interfere with the nourishment of the tubules, with consequent impairment of their power of selective reabsorption and, thus, to concentrate the filtrate. The urine is, therefore, much more dilute, as a rule, than normal. It has been mentioned that the specific gravity of the urine in chronic nephritis cannot be raised above 1 010 or so. Nor can it be lowered much below this level by drinking a large volume of water. The changes in urine concentration (as indicated by the specific gravity) which can be accomplished so easily by the healthy kidney, in response to alterations in fluid intake, are quite beyond the power of the severely diseased kidney.

The chief manifestations of kidney disease are: (1) the retention in the blood of nitrogenous waste products (e.g., urea, uric acid, creatine, as well as of phosphates and sulfates); (2) impaired extion of water, or failure to concentrate the urine; (3) protein, casts, and perhaps blood in the urine; (4) edema; (5) high blood pressure (see p 143); and (6) acidosis, in the later stages of the disease.

In severe nephritis the concentration in the plasma of urea and the other wastes mentioned above may be several times greater than in health. The difficulty in the excretion of water is shown by giving the subject a quart or so of water to drink and measuring the urine output hourly for a few hours thereafter. The normal person excretes all the ingested water within from 2 to 4 hours, but when the kidneys are seriously damaged only a part of the water is excreted by the end of 4 hours.

Normal urine is free from protein, but in kidney disease large quantities may appear. It is derived from the plasma. The diseased glomerular membrane permits the passage of serum albumin and smaller quantities of serum globulin (whose molecule is larger than that of albumin); fibrinogen, on the other hand, which has a larger molecule than those of the other plasma proteins, does not escape into the urine. When the loss of protein in the urine is excessive, its concentration in the plasma is reduced The osmotic pressure of the latter falls in consequence, and edema results (see p. 101).

## Micturition or the Voiding of Urine

The tubules of the kidney empty through a number of short ducts into a funnel-shaped membranous structure called the pelvis (basin) of the kidney. A narrow duct, containing smooth muscle in its walls and called the ureter, conveys the urine from the kidney pelvis to the urinary bladder. The urine is propelled along the ureter by peristaltic contractions; it enters the bladder in jets which occur at the rate of from one to five per minute.

The urinary bladder serves as a reservoir for the urine. It is a hollow muscular organ lined by modified stratified (transitional) epithelium. It has a maximum capacity of from a half to one pint. The most dependent part of the bladder leads into the urethra, through which the





Fig. 7.10 Sketch of the urinary system from in front.

urine is passed to the exterior (Fig. 7.10). This junction of the bladder with the urethra is guarded by a collection of circularly arranged, smooth muscle fibers, called the internal vesical sphinter. A little above and behind the urethral opening, the bladder wall is pierced obliquely by the ureters. The triangular area of the bladder wall marked off by the urethra and the two ureteral openings is called the trigone. The muscle forming the rest of the bladder wall is called the trigone. The caliber of the first inch or so of the male urethra is controlled by struated muscle, which is generally referred to as the external vesical sphincter.

Evacuation of the bladder. As urine gradually collects within the bladder, the pressure within it shows very little change for a time, the tone of the detrusor muscle adapting readily to the increased volume



አጵሮ

Fig. 7.11 Curve of pressure changes in the human bladder during filing (upper curve) and emptying (lower curve). Note that the internal pressure shows little sustained increase until the volume of fluid amounts to more than 200 cc. (Denny-Brown and Robertson)

(see postural tone, p. 294). Not until the urine volume is between 200 and 300 cc does any noticeable rise in internal pressure result (Fig. 7.11). Rhythmical contractions of the detrusor muscle then occur in response to the tension exerted upon the muscle fibers. The contractions are weak at first but, gradually gaining strength as the urine volume increases, culminate in a strong reflex contraction of the bladder wall, accompanied by relaxation of the internal sphincter and trigone.

The act of micturition, though dependent upon reflex mechanisms, is, in the adult, very largely under voluntary control. The reflex contractions of the detrusor can be inhibited, and contractions of the internal sphincter induced, by an effort of the will. Thus, the reflex mechanisms can be restrained until an opportunity for voiding the urine presents itself. The restraint is then removed, and the contractions of the bladder wall which ensue are reinforced by voluntary nerve impulses transmitted along the motor nerves supplying the

detrusor muscle. A rise in intra-abdominal pressure, due to contractions of the abdominal muscles and fixation of the diaphragm in the inspiratory position, usually precedes and accompanies the act of micturition. However, a voluntary movement of this character is not essential, the bladder mechanism being capable itself of expelling the urine.

The motor fibers to the detrusor muscle, and the inhibitory fibers to the internal sphincter and trigone, are derived from the parasympathetic division of the autonomic nervous system through the pelvic nerves (second, third, and fourth sacral segments). These fibers are peculiar in that, as indicated above, they transmit voluntary as well as involuntary impulses to the detrusor muscle.

The sympathetic division of the autonomic nervous system exerts an opposite effect upon the movements of the bladder. Through the hypogastric nerves it transmits inhibitory impulses to the detrusor and motor impulses to the internal sphincter and trigone (Pl. 7 B).

The external sphincter of the male, which is composed of striated muscle, is supplied with motor fibers through the pudendal nerves. The afferent nerves of the reflex arc governing micturition run mainly in the pelvic nerves.

The centers of the micturation reflex are situated in the midbrain, pons, and sacral part of the spinal cord (second, third, and fourth sacral segments). Micturition is carried out through the spinal center when the cord is severed above the sacral region Even after destruction of the spinal center, the bladder empties automatically, its movements being then brought about through the nerve plexuses in close relation to the bladder wall

The nervous mechanisms through which restraint is exercised in the control of micturition—that is, the inhibition of the micturition reflex—is late in developing to the state of efficiency where, as in a healthy adult, the urine can be retained in even an uncomfortably distended bladder. In infants, urine is voided involuntarily with no effort being made at restraint. In young children, though there may be complete control of the act of urination during the day, the emptying of the bladder during sleep, "bed-wetting" or nocturnal enuresis, as it is called, is of common occurrence. The age at which restraint is fully developed—at night as well as during the day—varies with the nervous constitution of the child and with his training, but is established as a rule, except for an occasional lapse under unusual circumstances, at 3 to 4 years. In persons of neurotic disposition, nocturnal enuresis may persist into adult life and, in the feeble-minded, restraint may fail to be exercised either during sleep or in the waking hours.

# Digestion

The alimentary or digestive tract comprises the mouth, pharynx, sophagus (or gullet), stomach, and intestines. The pharynx receives he food from the mouth and transfers it to the esophagus, along which is conveyed to the stomach. The stomach evacuates its contents into ne small intestine which discharges, in turn, into the large intestine see Fig. 8.1). A descriptive outline of the structure of each part of the igestive tract, in so far as a knowledge of structure is necessary for an inderstanding of function, will be given in the section dealing with the physiology of that particular part.

The various processes carried on in the alimentary tract which bring bout the physical and chemical changes in the food—mechanical distegration, the decomposition of large molecules into simpler and adily soluble compounds, and the absorption of these into the blood are all embraced under the word digestion. Digestion involves, erefore, mechanical factors (mastication, swallowing, and the movements of the gastrointestinal tract) as well as chemical. Digestion in e different parts of the alimentary tract will be described in order, immencing with the mouth. In each instance, a description of the temical changes which the food undergoes during the digestive procs, and of the nervous and hormonal mechanisms controlling secretion the digestive juices, will be followed by an account of the movements the part under discussion. First of all, a general account of enzymes ferments and of the properties of smooth muscle, of which the alls of the digestive tube are largely composed, must be given.

## Lazymes (G. zymē = yeast) or Ferments and Their Action

These substances—e.g., the ptyalin of saliva, the pepsin of gastric juice, and the trypsin of pancreatic juice—are responsible for the 288




Fig. 8.1 Diagram of the gastrointestinal tract. Most of the small intestine (ileum) excised.

chemical changes which the food undergoes during digestion. The chemical changes comprise the breaking up of the large molecules of carbohydrates, proteins, or fats into smaller molecules which are able to pass through the intestinal mucosa into the blood stream. The chemical process is called hydrolysis (Gr. hudor, water + lysis, breaking up) since, as a preparatory step to the splitting of the larger molecules, a molecule of water is taken up. Thus, the cleavage of a molecule of maltose into two of glucose is represented by the following equation:

 $C_{12}H_{22}O_{11} + H_2O = C_6H_{12}O_6 + C_6H_{12}O_6$ maltose water glucose glucose Such reactions, in order to be accomplished by the ordinary chemical methods of the laboratory, require the use of strong reagents, and usually a high temperature. The digestive ferments act at a relatively low temperature (that of the body), and without the aid of harsh chemicals.

It is not only the chemical reactions involved in digestion that are dependent upon enzyme action. These substances are concerned in the great majority of chemical reactions which occur in the tissues of both animals and plants and upon which the properties of living matter depend Enzymes or at least the great majority of them, are protein in nature. Though produced by living cells, they do not all require the presence of cells in order to act. Certain enzymes of yeast, for example, when expressed from the yeast cells, are capable of exerting their usual effect—the conversion of sugar to alcohol.

Several theories have been put forward to explain the manner in which enzymes act, but none can be said to have been definitely established. However, many facts are known concerning the action of enzymes which indicate that they belong to the group of substances known in chemistry as catalysts. Now, a catalyst increases the velocity of a chemical reaction but does not actually initiate the reaction. Nor loss a catalyst become altered or form a part of the product of the reaction. Catalytic action has been compared by Sir William Bayliss to the effect which a drop of oil exerts upon the speed of a metal weight moving down a glass incline. The oil hastens but does not initiate, the movement of the weight, nor does it combine with the metal or the glass or undergo any chemical change whatever. To take a specific instance of catalytic action when acetic acid and ethyl alcohol are mixed ethyl acetate is formed slowly, but the velocity of the reaction is greatly increased by the addition of a very small amount of hydrochloric acid. Thus

It will be seen from this equation that the hydrochloric acid does not enter into the reaction. The latter, as shown by the arrows, is revers ible. That is, the combination of acetic acid with ethyl alcohol to form ethyl acetate is accompanied by a reaction in the opposite direction—namely, the breakdown of ethyl acetate into acetic acid and ethyl alcohol. When the velocities of these two opposing reactions are equal, a state of equilibrium has been reached. If the concentration of ethyl

acetate is raised, the velocity of the reaction on the left is increased—that is, the reaction shifts to the left until equilibrium is reached again, if the concentrations of acetic acid and ethyl alcohol are raised, the reaction shifts to the right, or the concentration of ethyl acetate is reduced by removing it as it is formed. The hydrochloric acid increases the velocity of the reaction in either direction, it is an inorganic cata lysts—that is, catalysts formed by living organisms. It should be pointed out that both types of catalyst exert their effects in very low concentrations, a small quantity of ptyalin, for example, will convert a relatively large quantity of starch to maltose, 'a little leaven leaven eth the whole lump"

Enzymes differ in one important respect from most of the inorganic catalysts in being absolutely specific in their actions. Ptyalin, for ex ample, acts upon starch, but has no effect whatsoever upon proteins or fats, pepsin converts protein to proteoses and peptones, but has no action upon the other two types of foodstuff.

Reversibility of enzyme action. The actions of most enzymes are reversible. That is to say, after catalyzing a certain reaction, they can drive that reaction in the opposite direction and cause the reverse effect. This is indicated in the equation shown above, and has already been noted for carbonic anhydrase (p. 112). But in the case of the digestive enzymes and many others, reversibility of action does not occur in the body to any appreciable degree, though it can be demon strated in vitro. It has been shown, for example, that pepsin (p. 307) can synthesize a proteinlike substance from a solution containing a mixture of aminon acids.

Conditions which influence the action of enzymes. 1 Optimum temperature Enzymes act with maximum efficiency at a certain tem perature (optimum temperature) Lowering the temperature below, or raising it above, this level slows the reaction There is a certain definite temperature for each enzyme at which it acts at maximal efficiency. The digestive and other enzymes of animals act best around the temperature of the body A high degree of heat (above 60° C) perma nently abolishes their action

2 Hydrogen ion concentration The optimum pH for the different enzymes varies widely, for pepsin it is around 15, for pancreatic amy plase about 70 The former enzyme is mactive in an alkaline medium, the latter in a strongly acid medium

<sup>&</sup>lt;sup>2</sup> These facts are embodied in the law of mass action which states that "the velocity of a chemical reaction is proportional to the concentrations of the reacting substances"

- 3. Inorganic salts. Enzyme action is profoundly affected by certain ions, notably chloride which is necessary for the activation of the amylases. Magnesium greatly accelerates the action of phosphatase, and trypsin is activated by calcium. Iron is essential for the action of certain respiratory enzymes in the tissues. Substances such as these, which activate or enhance the action of enzymes, are called coenzymes.
- 4. Poisons. Mercury, gold, copper, and several other heavy metals exert a strongly toxic effect owing, apparently, to their combining with the enzyme. Cyanudes abolish the action of oxidizing enzymes of the tissues (p. 263). Fluorides, bromine, iodine, and chloroform are also toxic for some enzymes.
- 5. The products of the enzyme reaction. It has been mentioned (p. 290) that an enzyme reaction comprises two processes—decomposition and synthesis—proceeding simultaneously. As the equilibrium point is approached the velocity of the reaction gradually diminishes. For example, in the conversion of starch to maltose by ptyalin, the reaction proceeds rapidly at first, but as the concentration of maltose increases, the velocity of the opposing or synthetic reaction—maltose to starch—increases. At a certain stage the quantity of starch broken down in a given time just balances the quantity resynthesized. In other words, the enzyme reaction, to all intents and purposes, ceases as a result of the accumulation of its own products. If more starch (substrate) is added, the reaction starts again and continues until equilibrium is again reached.

Nomenclature. The material upon which an enzyme acts is called the substrate, and it is now the practice to derive the name of the enzyme by adding the suffix "ase" to the name of its substrate. For example, an enzyme which hydrolyzes starch is called an amylase (L. amylum, starch); one which acts upon protein, a proteinase; and on fat, a lipase (Gk. lipos, fat). Amylolytic (starch-splitting), proteolytic (protein-splitting), and lipolytic (fat-splitting) are corresponding terms used to denote the respective actions of these three types of enzyme in some instances the name of the enzyme is coined by suffixing "ase" to the name of the reaction which it brings about. Thus an enzyme which causes deaminations (p. 363) is called a deaminase and an oxidizing enzyme an oxidase.

Names were given to certain digestive enzymes before the system of terminology just outlined had come into use. The ferment of the saliva, for example, has long been known as pryalin. Pepsin of the gastric juice, and trypsin (protein-splitting enzymes), amylopsin (starch-splitting enzyme), and steapsin (fat-splitting enzyme) given to

the ferments of pancreatic juice are also old-established names. Corresponding terms in the new system are salivary amylase, gastric proteinase, and pancreatic proteinase, amylase, and lipase.

The physiological properties of smooth (nonstriated) muscle. Smooth muscle differs from skeletal muscle in the following respects: (a) slower, more sluggish contraction; (b) greater extensibility; (c) the exhibition of a sustained contraction or tonus for long periods, even though separated from the central nervous system; (d) the power of rhythmical contraction; (e) the possession of a double autonomic innervation (parasympathetic and sympathetic); (f) greater sensitivity



Fig. 8.2 Diagram to illustrate the manner in which muscle fibers may slide over one another and increase the capacity of a hollow organ. The upper group of four muscle cells is from a hollow organ whose walls are contracted and its cavity almost obliterated, the lower group are the same muscle fibers when the organ is full. (Redrawn from Gruttner.)

to thermal and chemical influences and to certain types of mechanical stimulation, such as stretching, but a lower excitability to electrical stimulation.


Tonus of smooth muscle may be defined as the steady sustained contraction through which the muscle offers resistance to a stretching force. The rhythmical contractions are superimposed upon the tonus state which may vary independently of the rhythmical contractions themselves. The tonic contraction of smooth muscle is associated with a negligible expenditure of energy. It is relatively insusceptible to fatigue; heat production and electrical changes are not detectable; and a rise or fall in the degree of tonus is not accompanied by a corresponding change in oxygen consumption.

Certain hollow organs, such as the stomach and urinary bladder and, to a less degree, the intestines, have the remarkable ability to enlarge when their contents are considerably increased without showing any rise in internal pressure. The tone of the muscle composing the walls of these organs becomes adjusted automatically to the distending 294

force This postural tone, as it is called, is very difficult to explain. It is possible that, in the stomach at any rate, the individual fibers which are disposed in layers, instead of being lengthened by the stretching force, simply slide over one another, the wall of the organ thus being increased in area but reduced in thickness. This could occur with little strain being placed upon the fibers themselves. In Figure 8.2 will be found a diagram illustrating the manner in which this might result

## Digestion in the Mouth

The structure of the salivary glands. Saliva is secreted by three pairs of glands—the submavillary, sublingual (or submandibular), and parotid glands. The submaxillary glands are situated beneath the floor of the mouth in close relation, one on either side, with the inner aspect of the lower jaw (mandible) The submaxiliary secretion reaches the cavity of the mouth through the duct of Wharton, which opens be neath the tongue The sublingual glands also he below the floor of the mouth, but nearer the mid line Each sublingual gland pours its



one side

secretion into the mouth be neath the tongue through a number of small ducts-ducts of Rivinus and of Bartholm The parotid gland lies below the ear, behind and overlapping the vertical part of the mandible (jawbone) Its duct -the duct of Stenson-opens upon the inner aspect of the cheek opposite the second up per molar tooth (see Fig. 83)

The salivary glands are composed of cells arranged in small groups around a central Showing the salivary glands of globular cavity called the al veolus or acinus 'The alveoli

are drained by fine ducts which join with those from neighboring alveolt to form larger channels. Through the successive junctions of smaller ducts and the formation of larger ones, the secretion is directed into the main secretory duct or ducts (Fig 8 4) This alveolar arrangement of the cells and the system of ducts of the salivary glands

Fig 8.4 Showing the plan of a racemose gland such as the salivary (diagrammatic). The small globular structures are alveoli or acini, some of which are represented as having been cut across



suggests a bunch of grapes To these and other glands (e g, the pancreas), showing a similar pattern, the term racemose (L racemus, a bunch of grapes) is therefore applied

The cells of the salivary glands are of two types, serous and mucous. The secretion of the former type is thin and watery, that of the latter type contains mucin, and is therefore thicker, and more slimy and viscous. The parotid gland is composed entirely of serous cells, whereas the cells of the sublingual gland are predominantly of the mucous type, serous cells being scarce. The submaxillary gland contains both serous and mucous cells in about equal proportions, the two types being seen in many instances in the wall of the same alveolus. The serous cells are then found on the outer side of the mucous cells which, therefore, he between the former and the alveolar cavity. Here they appear as crescentic bodies known as the demilunes of Gianuzzi (Fig. 8.5).




Fig 8.5 Microscopical section of a submaxillary salivary gland, showing mucous cells and demilunes of Gianuzzi




Fig 8.6 Diagram of the nerve supply of the salivary glands S-S-N superior salivatory nucleus I-S-N inferior salivatory nucleus S-G submaxillary ganglion, T P tympanic plexus

The control of salvary secretion The salvary glands are under the control of the autonomic (involuntary) nervous system, receiving fibers from both parasympathetic and sympathetic divisions of this system. The parasympathetic fibers to the submaxillary and sublingual glands arise from a center—the superior salvatory center—in the pons, they leave the brain in the facial nerve and are conveyed in the chorda tympani branch of the latter nerve to the cavity of the mouth. Here they join the lingual nerve which transmits them to the floor of the mouth where, separating again from the lingual fibers they make connections with ganglion cells situated in close relationship with the

sublingual gland, or within the substance of the submaxillary gland (see Fig. 8.6). Postganglionic fibers—the axons of the ganglion cells—pass to the secreting cells.

—pass to the secreting cells.

The chorda tympani nerve also carries vasodilator fibers (p. 196) to the glands and taste fibers to the anterior two thirds of the tongue. The posterior third of the tongue receives taste fibers through the glossopharyngeal nerve. The submaxillary and sublingual glands receive, as well, secretory and vasoconstrictor fibers from the sympathetic plexus investing the neighboring branches of the external carotid artery. The parotid gland derives secretory and vasodilator parasympathetic fibers through the tympanic branch of the glossopharyngeal nerve. These fibers arise from a group of cells in the upper part of the medulla oblongata called the inferior salivatory nucleus. Vasoconstrictor fibers to the parotid gland are derived from the sympathetic.

Stimulation of either the parasympathetic (chorda tympani) fibers or of the sympathetic fibers to the submaxillary or sublingual gland causes a secretion of salva. The secretion resulting from parasympathetic stimulation is, in most animals, profuse and watery in consistency; sympathetic stimulation, on the contrary, causes a scanty secretion of a thick and slimy mucinous juice. Apparently, then, the parasympathetic fibers innervate the serous cells, and the sympathetic fibers the mucous cells. Stimulation of the parasympathetic fibers to the parotid gland also causes a profuse watery secretion, whereas no obvious secretion follows stimulation of the sympathetic.

Under natural conditions, the secretion of saliva is a reflex phenomenon usually brought about by the stimulation of the taste fibers in the mouth. But stimulation of the ordinary sensory nerves in the mucosa of the mouth—that is, those fibers conveying sensations of touch, pain, and temperature—evokes a flow of saliva. We are all familiar with the salivation which results from moving some tasteless object around in the mouth, or from the manipulations of the dentist. Indeed, secretion of saliva may follow stimulation of almost any afterent nerve in the body. In animals, stimulation of the central cut end of the sciatic nerve excites salivary secretion and, in man, irritation of the nerves of the stomach or esophagus, in diseased states, is not uncommonly a cause of troublesome salivation.

Salivary reflexes are of two types. The salivary response evoked by the introduction of material into the mouth is called an unconditional reflex. But it is well known that salivation very often results from the sight, smell, or even the thought of food. We often hear a person say that his "mouth waters" when he sees or smells appetizing

food. The response following such forms of stimulation is called a conditioned reflex. The unconditioned reflex is inborn; it can be elicited immediately after birth. The conditioned reflex, on the contrary, depends upon experience. In order for it to become established, an association, upon some previous occasion, must have been formed in the brain between sensations received through the nerves of taste and those gained through some other sense organ (e g., eye, nose, ear, etc.). We need say no more here of this type of reflex; it will be considered in more detail in Chapter 12.

The characters, quantity, and composition of saliva. Normal human saliva as collected from the mouth (i.e., mixed saliva, containing secretions of all six glands, as well as the secretion of the small mucous glands scattered diffusely over the mucosa of the mouth) is slightly acid in reaction, varying between pH 6.35 and 6 85. Its specific gravity is between 1.002 and 1.008. The average adult secretes from 1200 cc to 1500 ce of saliva in 24 hours. The composition of saliva is given in the following table.

#### Table 8-1

### I. Salts (approximately 0 2 percent)

Sodium and potassium chloride Sodium bicarbonate Acid and alkaline sodium phosphates Calcium carbonate and calcium phosphates

Potassium sulphocyanate (thiocyanate)

#### II. Gases

Carbon dioxide and oxygen

#### III. Organic substances

Ptyalin (salivary amylase) Maltace

Serum albumin Serum globulin


Urea

Mucin, mainly from the submaxillary and sublingual secretions

The functions of saliva, 1. Digestive. The enzyme of the salivaptyalin-acts upon starch. The starch molecule is split into smaller molecules of the disaccharide, maltose. The rapid passage of the food through the mouth precludes the possibility that starch is acted upon here to any appreciable extent by the saliva. Whether starchy food, after its thorough impregnation with saliva, undergoes any important degree of digestion in the stomach has been debated. Ptyalin is active

in an alkaline, neutral, or faintly acid medium, but is inactivated by strong acid. It was thought, therefore, that the highly acid gastric juice would prevent or soon terminate salivary digestion. It has been shown, however, that the latter part of the meal, which usually contains the carbohydrate, may remain in the upper part (fundus) of the stomach, and thus be protected for some time, from the acidifying action of the gastric juice, by the mass of food ingested previously. For this reason it is likely that, under favorable circumstances, the salivary digestion of starch may proceed for a considerable period after food has reached the stomach.

If boiled starch is placed in a test tube with mixed human saliva and kept at body temperature, a slow conversion of the starch into maltose takes place The chemical change occurs in a series of stages which may be distinguished by the manner in which the product of each stage reacts with iodine Iodine gives a characteristic blue color with boiled starch. A short time after the saliva has commenced to act, a physical change may be seen to have occurred in the starch. It loses its opalescent appearance and becomes soluble, though it still gives the blue color with jodine. After a short time, this soluble starch becomes partially broken down, being converted into a dextrin which, since it gives a red color with iodine, is known as erythrodextrin Small amounts of maltose may also be detected at this stage Still later, no color reaction occurs upon the addition of iodine, a colorless product -achroodextrin-has been formed Finally, the starch is entirely con verted into maltose and isomaltose. In the final stage, traces of glucose may also appear, due to the presence of an enzyme maltase present in saliva in low concentration which splits the disaccharide maltose (see p 367) into two molecules of glucose The following scheme illustrates the action of ptyalin



Ptyalin has no action upon cellulose, and for this reason the starch must be cooked in order to rupture the cellulose envelope surround ing the starch grains, and thus allow the enzyme to come into contact with the starch

- 2. Preparation of the food for swallowing by altering its consistency. This is one of the most important functions of the saliva; the dood moistened, thus enabling it to be rolled into a plastic mass, which, by virtue of the slimy nature of the mucin which the saliva contains, is given a lubricant coating. Claude Bernard showed that a horse with a parotid fistula (the saliva draining to the outer surface of the cheek) had the utmost difficulty in swallowing dry hay or oats.
- 3. Solvent action. Taste is a chemical sense (Chap. 13). All solid substances, therefore, in order that they shall stimulate the taste buds, must be dissolved in the salva.
- 4. Cleansing action. The constant flow of saliva exerts a very necessary cleansing effect. The mouth and teeth are rinsed and kept comparatively free from food debris, shed epithelial cells, foreign particles, etc.; in this way the saliva tends to inhibit the growth of bacteria by removing material which may serve as culture media. In order to realize how important its function is in this regard, one has but to consider the foul condition of the mouth that occurs in certain fevers (such as typhoid), when the salivary secretion is suppressed. Then, decomposing organic material, swarming with bacteria, collects upon the teeth and lips, and must be removed by artificial means (e.g., by wiping with moistened eauze).
- 5. Moistening and lubricating action. The saliva, by moistening and lubricating the soft parts of the mouth and lips, keeps them pliable and resilient for the purposes of articulation. Frequent sips of water are almost essential for some public speakers in whom, as a result of evaporation from the mouth during speech, or owing to nervousness with consequent inhibition of salivary secretion, the lips and mouth become dry.
- 6. Excretory. Many substances, both organic and inorganic, are excreted in the saliva. Drugs such as mercury, potassium iodide, lead, etc., when introduced into the body, are excreted, in part, by the saliva. Severe inflammation of the oral mucosa (stomatitis) may be caused by the excretion through this route of excessive amounts of mercury. The blue line on the gum margins, in lead poisoning, is due to the metal having been excreted in the saliva and deposited as the sulfide. The sulfur is provided by organic material contained in the tartar formed on the bases of the teeth. For this reason the discoloration of the gum does not occur where teeth are absent. In chronic nephritis the saliva contains a high percentage of urea; sugar sometimes appears in severe diabetes; and in parathyroid overdosage the calcium concentration of the saliva is elevated. Severed types of microorganisms, some

intensely virulent—e.g., the virus of hydrophobia (rabies)—are excreted in the saliva.

Mastication. This act comprises the movements of the lower jaw, tongue, lips, and cheeks, whereby the food is reduced to a soft mass which can be easily swallowed. The incisor and canine teeth are used chiefly for tearing the food and breaking it up into small pieces, the




Fig. 8 7 The mouth, nose, pharynx, larynx, and the commencement of the esophagus, as exposed by a section a little to the left of the median plane of the head.

molars for grinding it. The movements of the tongue, lips, and cheeks move the food about in the mouth, apply pressure or suction to it, and introduce fresh material between the jaws. The whole mass thus becomes thoroughly impregnated with saliva. The jaw movements are brought about by the contractions of temporal, masseter, and pterygold muscles, all of which are supplied by the third division of the trigeminal (fifth cranial) nerve.

Swallowing or deglutition. The act of swallowing is usually described in three stages. In the first stage, the masticated food mass or bolus, as it is called, is moved to the upper surface of the tongue, which is then drawn sharply backward by the action of muscles, chiefly the mylohyoid, attached to its base. The food is thus projected into the

302

pharynx The opening into the pharynx, which is called the fauces, is limited on either side by two muscular pillars (the pillars of the fauces) As a result of the backward movement of the tongue, which at the same time presses against the hard palate, a pressure of 20 cm of water is created in the posterior part of the mouth and pharynx. The first stage of swallowing is a voluntary act

In the second stage, the food passes through the pharynx This stage is very brief. The muscles of the pharyngeal wall (constrictors of the pharynx) contract and, gripping or squeezing the food, force it into the esophagus There are, however, three other paths which the food might take—into the nose, into the larynx, or back again into the mouth (see Fig 8 7) The food is prevented from entering the nose by the elevation of the soft palate, the posterior edge of the latter is brought against the posterior pharyngeal wall, thus closing off the pharynx from the posterior entrance to the nasal cavities. This move ment rarely fails to take place at the proper time in normal persons, but in the paralysis of the elevators of the palate, which sometimes follows diphtheria, the passage of liquids into the nose during swal lowing is a common occurrence. The base of the tongue, which con tinues to be held in position against the fauces, and the approximation of the muscles forming the pillars of the fauces effectively block com munication with the cavity of the mouth Elevation of the larynx, which brings its opening under the shelter of the epiglottis and base of the tongue, prevents the food from entering the air passages This upward movement of the larynx is an essential part of the swallowing act When as a result of disease, the larynx is fixed swallowing becomes very difficult or impossible When the movement is not accurately timed with the pharyngeal movements, food enters the larynx and excites the coughing reflex Everyone at some time or another has experienced this accident

A short inspiration (inspiration of swallowing) occurs at the commencement of the first stage, this is followed by complete inhibition of respiration, which persists until the end of the second stage

or respiration, which persists until the end of the second stage. The third stage is occupied in passing the food down the esophagus into the stomach. In man, the upper two thirds of the esophagus is composed of striated muscle, the lower third of smooth muscle. The food is carried through the tube by a peristaltic contraction (p. 336) of the muscular wall, the passage of food being much more rapid through the upper striated portion of the tube than through the lower part. The esophagus of the dog is composed throughout of striated muscle, which accounts for the amazing rapidity with which this animal can swallow

food In the lower end of the esophagus—that is, where it opens into the stomach—the muscle fibers are condensed to form the cardiac sphincter. The sphincter at ordinary times is contracted, but it relaxes upon the approach of the food, which is then swept into the stomach by the peristalite waves. The second and third stages of the swallowing act are entirely reflex in nature—i e, involuntary. The reflex is initiated by the passage of the food through the fauces. Afferent nerve endings are stimulated by the contact of the food with the mucosa in the region of the fauces and pharynx. Of course, we can swallow at will, even though no food is in the mouth. It might be thought, therefore, that it was not correct to speak of the second and third stages of swallowing as being involuntary. But when one wishes to swallow, food or (if the mouth is empty) a little saliva, by a movement of the tongue, is passed backwards, to act as a stimulus. Swallowing then occurs automatically

Innervation of the swallowing reflex. The afferent fibers of the reflex are contained in the branches of the trigeminal, glossopharyngeal, and vagus nerves supplying the mucous membrane in the region of the fauces, tonsil, and pharynx. Temporarily abolishing the function of the nerve endings in these regions, by the application of a local anesthetic (e.g., cocame) to the mucous membrane, renders swallowing impossible for a time. The chief swallowing center is situated in the medulla oblongata, in the neighborhood of the nucleus of the vagus. The efferent fibers travel to the various muscles taking part in the act by the hypoglossal (to the muscles of the tongue), by the trigeminal (to the mylohyoid muscle in the floor of the mouth), and by the glossopharyngeal and vagus nerves to the muscles of the pharynx and esophagus.

The nervous control of the cardiac sphincter has for years been a question of some dispute, but in man, at any rate, the vagus appears to exert an inhibitory effect upon the muscle, while excitatory fibers are of sympathetic origin. It sometimes happens, though rarely, that the innervation of the sphincter becomes disordered, the sympathetic being overactive (or the vagus underactive). The muscle, as a result, remains tonically contracted during swallowing and thus prevents the easy entrance of food into the stomach. Difficulty in swallowing is experienced, and the esophagus for some distance above the obstruction becomes distended by retained food. This condition is called cardiospasm.

<sup>2</sup> A sphincter is a circular band of muscle which acting like a purse string when it contracts serves to narrow the entrance to or outlet from certain hollow organs




Fig 88 Interior of the stomach showing rugae of the mucous membrane




Fig 8.9 Diagrams of the stomach empty on left, full on right

## Digestion in the Stomach

General description of the stomach. The walls of the stomach are composed of smooth muscle fibers arranged in three layers—longitudinal, circular, and oblique—in this order from without inward, and a lining of mucous membrane. The mucous membrane contains the gastric glands; its surface epithelium is of the columnar nonciliated type. When empty, the stomach is a narrow tubular organ; its mucosa is thrown into numerous longitudinal folds called rugae. Its contour varies somewhat in different persons, but most commonly bears a general resemblance to the letter J (see Figs. 8.8, 8.9, and 8.11). The vertical part of the J is called the body; the upper part of the body is expanded into a domed chamber by a bubble of gas, and is called the fundus. The fundus lies in contact with the diaphragm which inter-

venes between it and the heart. The region of the stomach corresponding to the book of the I is named the pyloric part (pars pylorica). An annular band of muscle, the pyloric sphincter, surrounds the opening between the pyloric part and the duodenum (first 10 inches or so of the small intestine). The wails of the empty stomach, except in the region of the fundus, are in contact, but the food after passing through the cardia presses steadily downward, separating the gastric walls. As the organ distends it becomes pear




Fig. 8.10 Mucosa of the stomach highly magnified to show foveolae.

shaped; its rugae become flatter, and, with increasing distension, disappear. The capacity of the human stomach is from 1 to 1½ quarts.

The gastrie glands. The gastric juice is secreted by the gastric glands which are scattered diffusely in large numbers throughout the gastric mucosa. The total number of glands in the human stomach is around 35,000,000. They are minute tubular or flask-shaped structures which




Fig. 8.11 (Left) X-ray photograph of a normal J-shaped stomach. (Kindness of Dr. A. C. Singleton.)
Fig. 8.12 (Right) Glands from fundus of stomach. 1, pit or foveola on mucous surface of stomach; 2, neck of gland containing mucin-secreting cells; 3, bottom (fundus) of eland, 4, parietal cells: 5, chief cells.

open, each one separately, into a smaller, narrow pit called a *foveola* upon the inner (epithelial) surface of the stomach. The gastric mucosa is dotted with immense numbers of such foveolae (Fig. 8.10).

The cells forming the walls of the glands are continuous with the epithelium covering the gastric mucosa, but differ from the latter both structurally and functionally. The glands in the fundus and the body of the stomach are also different in their histological appearance and in their secretory functions from those in the lower part of the stomach (pyloric region). The cells of the glands in the fundus and body of the

stomach are of three types and are called, respectively, (a) mucous neck cells or chief cells of the neck, (b) chief or zymogenic cells, (c) parietal or border cells. The mucous neck cells are situated in the narrow superficial part of the gland tubule called the neck. They secrete mucus (Fig. 8.12). The chief or zymogenic cells form a continuous lining for the deeper part or body of the gland tubule. These cells secrete the enzymes of the gastric juice. The parietal or border cells do not form a continuous layer, but lie here and there along the tubule to the outer side of the chief cells. They are entirely responsible for the production of the hydrochloric acid of the juice.

The glands of the pyloric region contain only mucus-secreting cells; chief cells and parietal cells are absent; their secretion is alkaline in reaction. The mucosa for a small area near the cardia also contains

glands of this type.

The composition and actions of gastric juice. The gastric juice contains enzymes, mucin, and hydrochloric acid. An analysis of human gastric juice is given in Table 8-2.

#### Table 8-2

#### Human gastric juice

Acidity Free HCl, 0.50 to 0.60 percent Total acidity, 0.45 to 0.60 percent pH, 0.9 to 1.5

Solids Organic, including mucin and the various ferments, 0.42 to 0.46 percent

Inorganic, 0.13 to 0.14 percent

Specific gravity, 1.006 to 1.009

Total nitrogen, 0.051 to 0.075 percent

The enzymes are pepsin, rennin, and lipase. The pepsin acts upon protein. The protein molecule, as we shall see (p. 360), is constituted of large groups of amino acids linked together. The first important action of the gastric juice upon the food is the combination of the hydrochloric acid with protein material to form acid metaprotein. The pepsin then acts upon the latter, splitting the protein structure into smaller molecules called proteoses and peptones. The attack made upon protein by pepsin ceases at this stage.

When the mucosa of the fundus or body of the stomach is extracted with an alkaline solution, the extract possesses little or no proteolytic activity, but becomes active upon acidification. It is, therefore, sometimes stated that pepsin is secreted as an inactive precursor which differs chemically from active pepsin, and is referred to as pepsinogen. A simpler and more correct explanation of the observation that an alkaline extract of the gastric mucosa is inactive is based upon the fact that pepsin, like other enzymes, acts best at a certain pH (optimum pH), and becomes inactive if any great change from this reaction occurs. The optimum pH for the action of pepsin is around 1.5, which is the reaction of pure gastric juice.

The concentration of hydrochloric acid in gastric juice, as the latter is secreted by the glands, is from 0.5 percent to 0.6 percent. The contents of the stomach have rarely, however, an acidity as high as this, for the juice, after it reaches the cavity of the stomach, becomes diluted by other fluids—e.g., saliva, mucus from the gastric epithelium, the alka-



Fig. 8.13 High magnification of a portion of gastric gland to show intercellular and intracellular canaheuli.

line secretion of the pyloric glands, and food residues. The hydrochloric acid is formed by the parietal cells and is apparently secreted as such into the cavity of the tubules passing along fine canals (canaliculi) running between the chief cells. These intercellular canaliculi drain a network of still finer channels within the parietal cells (Fig. 8 13). The acid is not, as was at one time believed, secreted in the form of a precursor which is converted to hydrochloric acid after reaching the lumina of the tubules, or the cavity of the stomach. Indeed, it is a truly remarkable fact that the cells do form and secrete a mineral acid of such high concentration. Also, it may well be asked, why do not the pepsin and hydrochloric acid of the gastric juice digest the stomach itself? This is a conundrum which antil recently has received no satisfactory solution. It is no answer to this question to say that the stomach is hving tissue and so can protect itself, for, if the leg of a living frog is inserted into the stomach of a dog, it undergoes digestion about as readily as if it were a piece of beef. It has been discovered, however, that

Digestion of the mucous membranes by pepsin hydrochloric acid is a potent factor in the development of gastric or duodenal ulcer (p 315).




PLATE 1

Left sample of blood immed stely after centrifuging showing the separation of cells (about 45 per cent) from plasma (about 55 per cent). Right uncentrifuged sample of blood about one hour after it has coagulated and the clot has condensed or strated.





PLATE 2



1 corpuscles mixed with compatible serum - agglut nation of corpuscles caused by mempatible serum



Section of a lobule of the apleen 1 capsule 2 branch of spienic artery 3 Maly ghian corpusele 4 vein 5 spienic pulp showing blood annues (semi diagrammatic)






PLATE 4

Diagram of the circulation 1 systemic results on pulmonary circulation 3 hepatic artery 4 attents to gastro-intestinal tract and spicen 5 portal sein 6 hepatic circulation 7 renal streadation 8 rapidlary bed of lower limbs 9 capillary bed of head and neck. RA right surrice RV right ventrice LA left aurice LV left sentricle



#### PLATE 5A

Dagram of the efferent cardae nerves. I, vaguis: 2, pregangionic fibers of the accelerator nerves (sympathetic): 3, post-gangionic fibers of the accelerators constituting the superior, middle and inferior cardiac nerves. Note that the post-gangionic fibers of the vagus originate from nerve cells situated in the wall of the heart.





#### PLATE 5B

Sinus and aortic referes. Diagram I, cardonabilitory center; 2, vasoconstrictor center; 3, carotid sinus and sinus nerve, 4, spanual cord; 5, arterole; 0, heart; 7, efferent fibers of vagus; 6, arterole; 8, afferent fibers of vagus; 6, arterole; 200 constrictor fibers; 9, portagaljonic vasoconstrictor fibers; 9, portagaljonic vasoconstrictor fibers; Wasodiastor center and fibers not shown. Tracing below, taken from Bayliss, shows effect upon the blood pressure of stimulating the cardiac depressor herve. Arrow indicates direction of tracing.




PLATE 61

A g am of he reulation through the skin 1 epidern s 2 papilse 3 eap a 3 loot 4 term nai ar er ole 5 subpapilary enous ylexus. The venous 1 ub of he cay lary loop and the planus are shown the con entional blue color b set aly the blool in these bannels a heal his title darker than arrienal blood.





PLATE GB

Dagram Bust at ng the exchange of gases between the limit the alveolur #



#### PLATE 7A

Showing the coronary artery 2 right coronary artery 3 aorta 4 pulmonary artery 5 superior vens cava 6 in ferior vens cava.

#### PLATE 7B

Diagram of the innervation of the unnary bladder B bladder E external sphincter G hypogastric ganglion I in ternal sphincter K kidney U ureter L 1 2 3 4 refer to the ganglia of the lumbar sympathetic chain 9 1 2 3 indicate the first second and third sacral segments of the spinal cord.






PLATE SA Dagram I the nerve supply of the gastro-intest nal tract 1 ? vagus nerves Ali vagal fibers pregangione a d post ganglone nrel sympathete in blue pels c nerve black 3 4 an i 5 splanchn e nerves (sym tathet c) 6 7 and 8 prevertebral sympathet e gangl a 9 pelve nerve 10 to I ver C gangi ated cord of sympa tlete T 5-12 thoracc seg ments of sp nal cori L 15 I mlar segments S 1 3 sa cral segn e ta





#### PLATE 8B

Nerve plexuses of the nestinal wall. Per gangl on a fibers of vagua (1) in red po gang on a fibers of sympathetic (2) in blue plexuses (3 and 1) in values 3 intest nal will: 4 muscular an core 5 submucous (Me senera) plexus 6 crc lar muscular coat fibers shown in cross sect on 7 mysenters (Auerbache) plexus 7 longitud nal m ecular coat




PLATE 9A

Diagram of receptor and effector organs. 1 exteroceptors 2 interoceptors 3 proprioceptors (muscle spindles) 4 effector organs smooth muscle gland and skeletal muscle 5 central nervous system.





PLATE 9B

Diagram of a cross section of the spinal cord to show the main nerve tracts. Ascending tracts in blue descending in red 1 lateral (crossed) corticospinal tract 2 rubrospinal tract 3 lateral vertibulospinal tract 4 antenor vestibulospinal tract 5 faccilus cruestus gracing 7 posterior genocerbellar tract 8 faccilus gracing 7 posterior genocerbellar tract 8 antenor spinocerbellar tract 9 posterior spinothalamic tract 10 antenor spinothalamic tract 110 antenor formation from the first posterior spinal prevent 12 antenor horn cell (motor neuron) 13 fibers of posterior spinal nerve root connecting with Meccondary neurons in spinal gray matter




#### PLATE 10A

Disprain of the descending tracts of the core in Best cell of motor of the core in Best cell of motor area of the core in Best cell of motor area of the core in t

#### PLATE 10B

Diagram of the ascending tracts of tectud 1 opte thalamus filations of left aide not able with in order that the connections may be seen more a constant of the control of






PLATE 11

Plan of the autonomic nervous system Parasympathetic fibers on left, sympathetic on right, pregnaphone in Moch, postganglouen in Tod Parasympathetic enters in stippled shading. The Roman numerals refer to the respective cranic nerves A, cerebral cortex, B, hypothalamus, C, pitutary gland, D, mid-brain, E, pons, F, medulla oblongata, G, citary ganglion, H, sphenoplatine ganglion, postganglion, fibers to lachrymal gland. I, submanillary gainglon, postganglion, p







PLATE 12B

The interior of the posterior half of the left cycholl showing the entrance of the optic nerve (optic papilla or disk) retinal vessels macula later and fover centralis

PLATE 12 V


Diagram of the connections of the sympatric fibers in Effects in black preganglionic solid lines poet ganglionic interrupted lines afferent vasceral fiber in red Collateral ganglion prevertebral ganglion (Alter Best and Taylor The Physiological Basis of Virtual Parties)



### PLATE 12C


Dagram to illustrate the division of the autonome nervous system into cholmerare to discrete P programmer of the sympathetic 1 pregametric fibers 2 postganghous fibers should be supposed for fibers 2 postganghous fibers should be supposed for fibers postganghous fibers should be supposed for fibers for fibers fibers and fibers fibers from the fiber fibers f

ETC



#### PLATE 13A

Illustrating the course of the yield fibers from the tetum to the occupital cortex. I left yield field 2 right yould field 3 left yield a left yield cotton. T = temporal N = nazal 5 option change in the properties of the propert






PLATE 14A

Discrete of the three primary color sensations (Neug-Heimholts theory) I represents the red 2 the green and 3 the vicinity of the properties of the properties of the vicinity of the curve at which it is cut by the vertical has the extent to which the several primary sensations of color are extited by whereholds of different wave lengths. (Mittel Heimholts)






PLATE 11B

Chart showing color fields of the right retina. The entire area of the retina circumscribed by the periphery of a given colored band is sensitive to that color Area for yellow not shown. (After Howell.)



## PLATI 15 %

Map of tongue slowing detribut on of the fundamental taste censations. Yellow sweet Use ac I and salty green bitter I tonul "circu vallate pap like 3 function pag like



### PLATI 15B


The nerves of the ans. Con stretor part way, (affected and of ferent) o red dilators in Mersus affects duck interrupted nest is san le lary muscle "e clary gangt on 3 nucleus of oculo notor nerve 4 superior coll cult as the color of the co

## PLATE 15C

Very early hun an o un em bedded in uterine walf. Note the inte locking of the t-suc of tle ovum (chornon c vill) will the uterine itssue (dec dua). Tle red areas are blood a nuses a aminot ceavity 9 yolk sac 3 embryo (After Bryce)



Diagram showing a series of stages in the growth of the ovum and development of the embryo (Redrawn and modified from Gray) 1 ammion 2 yolk sac 3 chor on 4 chorionic vill 5 embryo 6 heart 7 umbil cal cord



Petal ere lation. Red oxygenated blood. Mue reduced blood purple mixed blood. C card dartey T1.0 engant services and a services and after services and after services and after services and after services and services and after services and ammonia is formed in the gastric mucosa which, being alkaline, protects the gastric tissue from the action of the acid.

The hydrochloric acid is formed from the sodium chloride of the blood, but, though several theories have been advanced, the manner in which the gland cells accomplish this chemical feat is not clearly understood.

Rennin coagulates milk, producing a flocculent mass called the curd, and a clear fluid known as whey. This action has been made use of from time immemorial in cheese making, the rennin being extracted from calves' stomachs. The curdling or clotting of milk presents a superficial chemical resemblance to the clotting of blood. As a result of the action of rennin, the soluble milk protein, casemogen, is converted into an insoluble product, and, as in the conversion of fibrinogen to fibrin by thrombin, calcium is necessary for this reaction. The role played by calcium is not, however, the same in both instances—whey protein and paracasem—both of which are soluble. The latter protein then combines with calcium to form calcium paracasein which is insoluble. The whey protein and the calcium paracasein are then digested by pepsin.

The lipase of gastric juice has a very weak, fat-splitting action and in the adult is of little practical importance. It acts only upon fats which are in very fine emulsion (e.g., cream and egg yolk). It may be

of more value to the infant.

Mucin belongs to the class of glycoproteins (p. 361). It is present in pure gastric juice. It is secreted by the cells in the necks of the glands in the body of the stomach, as well as by the pyloric glands. The secretion of the surface epithelium of the stomach (goblet cells, p. 37) also contains a high percentage of mucin. This latter secretion is a slimy fluid which tends to cling to the surface of the mucosa. It thus serves to protect the mucosa from injury by coarse particles of food and, to a certain extent, from the action of the pepsin hydrochloric acid. It may be mentioned in this regard that mucin is capable of combining with a relatively large amount of hydrochloric acid, and of inhibiting peptic activity. Preparations of mucin obtained from pigs' stomachs have been used recently in the treatment of gastric and duodenal ulcers with the object of protecting the surface of the ulcer from the action of the highly acid gastric juice.

The secretion of gastric juice. It is now customary to speak of three phases of gastric digestion: (1) psychic or cephalic, (2) gastric, and

(3) intestinal.

The psychic or cephalic phase. It was first shown by the Russian physiologist, Ivan Pavlov (1849–1936), that gastric juice is secreted while food is being chewed, and before any has entered the stomach The more appetizing the food, the greater is its effect upon gastric secretion. Pavlov, therefore, spoke of the secretion caused in this man-



Fig 8.14 Dog prepared with an phageal fistula for "sham feeding"

is, spoke of the secretion caused in this manner as the appetite puice. The secretion under such circumstances is, of course, purely reflex in nature. The afferent fibers of the reflex are in the nerves of taste, the effect annot be produced after the vagus nerves have been sectioned.

Pavlov demonstrated this reflex in dogs by what he termed "sham feeding." An animal was first prepared by dividing the esophagus in the neck, bringing the lower end of the upper segment through the neck wound and

fixing it there by sutures (Fig. 8.14). The animal, prepared in this way, relishes its food just as does any normal dog, but the food cannot enter the stomach; it simply falls from the open esophagus after it has been swallowed. Nevertheless, a flow of gastric juice results. The juice secreted by the stomach can be collected from an opening made through the abdominal wall into the stomach, and measured. It can then be analyzed for its peptic and acid concentrations, if desired; or a minature stomach can be fashioned.

The miniature stomach, as fashioned by Pavlov's method, is shown in Figure 8.15. It offers a means of obtaining pure gastric juice, uncontaminated by food or saliva, under various physiological conditions. Since it is in connection by nerves and blood vessels with the main part of the stomach, it is therefore under the same nervous and hormonal influences as the latter.

The psychic secretion of gastric juice has been clearly demonstrated several times in man. In Dr. Carlson's laboratory in Chicago, a man was studied whose esophagus was impassable as a result of an injury received in childhood. A permanent opening had been made through the wall of the abdomen into the stomach. The man could be fed only through this opening, which is called a gastric fistula. Through it samples of gastric juice were withdrawn, measured, and analyzed under various physiological conditions. He could also be "sham fed," though in his case the food was chewed and then spat out. It was found that food for which this subject had a particular fondness caused the




Fig 8 15 Upper drawings show Pavlor's method of fashioning a gastic pouch or ministure stomach. A horizontal incision a b is made which causes minimal injury to the gastine nerves and blood supply D vagus nerves. The flap C is turned down and the pouch S solated from the main cavity of the stomach V as shown in the right hand sketch. A, abdominal wall. The lower drawing illustrates the manner in which pure gastine junce can be collected from the ministure stomach while digestion is proceeding in the main part of the stomach.

greatest secretion of juice (Fig. 8.16) In some instances the mere sight or smell of food called forth a secretion. That is to say, a conditioned reflex (pp. 297 and 567) can be established for gastric secretion, as well as for the secretion of saliva.

The psychic secretion of gastric juice has also been demonstrated in persons during hypnosis A suggestion made to the subject that he was cating some tempting piece of food stimulated gastric secretion. On the other hand, psychic influences—e.g., worry, fear, anger, and pain—are powerfully inhibitory to the gastric glands, as are also bad odors, an unsavory character of the meal, or even an unattractive appearance of its general appointments. Pleasant surroundings, con-

tentment, delight for the eyes and ears—e g, flowers on the table cleanliness, and good music—tend toward an opposite effect The importance of appetite in aiding digestion has been pithily stated by Pavlov in the words "Appetite spells gastine juice." Macbeth at the banquet expresses a similar thought when he says, Let good digestion wart on appetite and health on both. It is not wise, therefore, to force a child to eat food that he detests

Variations in the secretory activity of the gastric glands are accompanied by changes in the blood supply to the gastric mucous mem




Fig. 8.16. Typical curve of secretion of gastne juice collected at 5 minute intervals during mastication of palatable food for 20 min utes. The rise in secretion during the last 5 minutes of mastication is due to chewing the dessert (fruit) for which the person had an especial fondness (After Carlson).

brane When the interior of the stomach is viewed through a gastric fistula or by means of a specially devised instrument known as a gas troscope the mucosa is seen to blanch in response to the inhibitory influences mentioned above and to become flushed when secretion occurs

The gastric phase. This refers to the secretion of gastric juice that occurs after the food has entered the stomach, it continues for a much longer period than does the psychic secretion. Let us now consider the mechanisms controlling the secretion at this time. The stretching of the stomach wall by the food has only a slight secretory effect. The secretion during this phase is due mainly to the action of a chemical material—a hormone—formed in the mucosa of the stomach. This material is absorbed into the blood stream and reaching the cells of the gastric tubules excites them to secretion. The hormone is produced in turn through the action upon the pylonic mucosa of certain substances in the food. Such substances are present in aqueous extracts.

of meats and vegetables (e.g., soups and beef extracts). For this reason they are referred to as extractives. On account of their stimulant action on secretion they are also known, from the physiological point of view, as eastric secretogogues.

The products of protein digestion—proteoses and peptones—and certain other materials act in a similar fashion.

When evidence for its existence was first obtained, the gastric hormone was named gastrin; its chemical nature was unknown. Extracts of the pyloric mucosa, obtained from an animal after a meal of meat or of other substances rich in extractives, contained the hormone. This was shown by the fact that, when such extracts were injected into the veins of an animal, a profuse secretion of gastric juice resulted.

It is now generally conceded, however, that the hormone in pyloric extracts—i.e., gastrin—is not a specific and hitherto unknown substance, but is simply histamine, which is found in other tissues as well as in the gastric mucous membrane. The discovery that histamine and gastrin are identical does not necessarily lead to the conclusion that there is no other gastric hormone. On the contrary, a considerable body of evidence suggests that a gastric hormone quite distinct from histamine is formed in the gastric mucosa during digestion. But, so far, this hormone has not been obtained by extraction.

The intestinal phase. After the partially digested food materials have reached the intestine, certain of their constituents (split products of protein digestion, meat extracts, etc.) continue to exert an excitatory effect upon gastric secretion. Little is known concerning the manner in which this effect is brought about; it may be dependent upon the formation or liberation of a hormone from the intestinal mucosa, for an extract of the latter, when injected intravenously, has an excitatory effect upon gastric secretion. Another possibility is that certain constituents of the food are absorbed into the blood stream and act directly as excitants of the gland cells.

The effects of different foods and of certain chemicals and drugs upon the secretion of gastric juice. The hydrochloric acid in the gastric contents is in part combined with food, especially in acid metaprotein and with mucus; the remainder is free. The combined acid and the free acid together are referred to as the total acid.

In man the activity of the gastric glands is investigated by giving a meal of toast and weak tea—the so-called test meal—and removing samples of the contents of the stomach every 15 minutes during a subsequent 2-hour period. The samples are analyzed, the percentages of total and free acid determined, and curves plotted (see Fig. 8.16).

Meat, since it is tich in extractives, stimulates the glands more power fully than does carbohydrate. Fats have an inhibitory effect upon gas trie secretion, as well as upon the movements of the stomach. A meal containing a high proportion of fat takes a relatively long time to leave the stomach. Fat also exerts an inhibitory effect upon both gas trie secretion and motility, after it has reached the duodenum. The latter effect is attributed to a chemical substance (a chalone, p. 403) formed in the intestinal mucosa, this substance has been named enterogastrone.

Among chemicals which stimulate gastric secretion, the most pow erful are histamine and alcohol. Atropine by its paralyzing action upon the vagus endings temporarily suppresses secretion. Alkalies such as sodium bicarbonate in repeated small doses excite the gastric glands, but a single large dose has, as a rule, an inhibitory effect

Abnormalities in gastric secretion. When no secretion of acid occurs after a test meal, the condition is called gastric anacidity of achlorhydra. About 4 percent of perfectly healthy and otherwise normal persons show this peculiarity. There may be a complete absence of peptic secretion as well, when such is the case, the condition is called achyling gastrica. Gastric anacidity is also seen in a number of diseased states—e.g., cancer of the stomach pernicious anemia chronic inflammation of the stomach chronic aritimits gall bladder disease, etc.

In some instances, the gastric glands though failing to secrete when stimulated by a test meal, respond to histamine the anacidity is then called false or apparent. In other instances even a powerful stimulus such as histamine or alcohol, is ineffective. This absolute failure of acid secretion is called true anacidity. An abnormally high acidity of the gastric contents is called gastric hyperacidity or hyperchlorhydria. In this condition, the acidity of the gastric juice as it leaves the gland intuities, is not higher than usual (0.5 to 0.6 percent) but owing to a high rate of secretion or to failure of the juice to be neutralized after it has reached the cavity of the stomach, the percentages of total and free acid in the gastric contents are much above the normal values. Under normal circumstances, the acidity of the gastric contents rises after a test meal, reaching a maximum in an hour or so and then declining (see chart Fig. 8.17). The return of the curve to the resting level is due to several factors—reduced rate of secretion by the glands, emptying of the stomach, neutralization by mucus and probably also, to some extent, by the regurgitation of alkaline juices from the duodenum. An abnormally high acidity of the gastric contents is due to

failure of one or another or all of these factors; the curve of gastric acidity rises above the normal maximum, and in many instances does not return to the resting level within the normal time.

Gastric hyperacidity is the commonest of all digestive complaints in otherwise normal and healthy persons. It causes "heartburn" and a feeling of discomfort in the chest. The burning sensation is due to the highly acid gastric contents being carried into the esophagus by contractions of the stomach, and stimulating sensitive nerve endines in the




Fig 8.17 Normal curve of free hydrochloric acid in the gastric contents following a test meal.

lating sensitive nerve endings in the esophageal mucous membrane. Ascending waves of contraction may carry material into the throat or the mouth. The gastric mucosa itself is insensitive to acid. Hyperacidity of the contents of the stomach is seen in a large proportion of cases of gastric and duodenal ulcer; indeed it is now generally agreed that it is the chief cause of these conditions.

In duodenal ulcer, for example, ulceration occurs where the gastric juice, after issuing from the pylorus, first comes into contact with the intestinal mucosa. Treatment of gastric or duodenal ulcer is therefore directed toward reducing the acidity of the gastric contents. This is accomplished by giving alkalies, or other substances which combine with the excess acid, such as preparations of mucin; by restricting the diet in respect to meat and other materials rich in extractives which stimulate gastric secretion; and by substituting foods (e.g., milk, cream, and carbohydrates) which either inhibit or exert no pronounced stimulating action upon the glands. Alcoholic drinks are forbidden, or allowed only in minimal amounts and well diluted Excessive smoking should also be avoided. Sometimes a preparation of enterogastrone is given for its inhibitory effect on gastric secretion and motility.

A high gastric acidity is also seen when the pylorus is obstructed and, as a consequence, gastric evacuation is delayed. The distension of the stomach which results from pyloric obstruction acts further as a stimulus to secretion.

The nerves which influence gastric secretion and motility. The gastric glands receive secretory fibers through the vagus nerves. When

these nerves are stimulated, the glands secrete a highly acid juice, rich in pepsin. Sympathetic fibers can also be traced to the gastric glands, but their action is less well known; it seems, however, that glands, but then action is less well known, it seems, nowers, and they innervate the glands of the pyloric region, causing a secretion of an alkaline juice consisting largely of mucus.

The movements of the stomach are under the control of both the

vagus (parasympathetic) and sympathetic nerves. The vagus is motor



in action—that is, it raises the tone and increases the force of the contractions of the gastric muscle, the sympathetic is inhibitory. The vagus and sympathetic nerves are also motor and inhibitory, respectively, to the pyloric sphincter.

Movements and emptying of the stomach. The upper half or so of the body of the stomach does not show active movements, but is the seat of a steady tonic contraction. When digestion is proceeding, peristaltic contractions commence about Fig 8.18 Drawing ceeding, peristaltic contractions commence about the middle of the body of the stomach showing peristalsis.

showing peristaliss, downward (i.e., toward the pylorus), become progressively more powerful (Fig. 8.18). The pyloruc part is, therefore, the more actively motile region, the motility becoming more and more marked as the digestive processes progress. These movements serve to mix the food thoroughly with the gastric juice and to break up food masses that have been softened by the digestive juices. When the food has reached a consistency suitable for discharge into the duodenum, it is propelled through the pylorus by the descending waves of peristalsis.

The pyloric sphincter is relaxed most of the time. It is contracted only when a peristaltic wave passes over it and closes the pyloric orifice for a moment The latter varies in diameter from time to time, however, being wider during the later stages of digestion than at the beginning. Thus the stronger peristaltic contractions and the greater diameter of the pyloric opening, when gastric digestion is nearing completion, facilitate the evacuation of the stomach. It should be emphasized that the consistency of the food is one of the most important factors determining the time after swallowing at which it leaves the stomach Fluids, for example, commence to leave it almost immediately after entering. Solid food in the stomach must first be reduced to a fluid or semifluid state, and those articles of diet which are most readily softened and liquefied (e.g., carbohydrates) leave before meats, which require a longer time to become reduced to a suitable consistency. Fats are evacuated most slowly, for, as already mentioned, they inhibit gastric secretion and motility. They also, for this reason, retard the evacuation of other types of food. An ordinary mixed meal is usually completely evacuated within from 3 to 44 hours.

The stomach contents, after they have become softened and form a nearly homogeneous semifluid or plastic mass, are referred to as the chyme. In this state, the food commences to pass into the duodenum. Chyme is an old word meaning juice, and arose from the belief that digestion in the

stomach was essentially a mechanical process, the nutritive juices being expressed from the food through the pressure exerted by the contractions of the stomach. That the food underwent chemical changes in the stomach was not fully realized until the French scientist René Réaumur (1683-1757), and the Italian priest and scientist, Lazaro Spallanzani (1729-1799) (Fig. 8.19), carried out their ingenious experiments. Réaumur gave a kite and other animals small perforated metal tubes filled with meat.



Fig. 8.19 Lazaro Spallanzani.

He found, upon recovering the tubes, that the meat, which could be acted upon by the gastric juice though protected from any mechanical action of the stomach, was partially dissolved. He also showed that gastric juice would digest meat in a test tube—that is, outside the body. Spallanzani performed similar experiments upon animals and upon himself. He swallowed perforated wooden tubes filled with food and showed that, though the containers were unbroken, after passing through the alimentary tract, the food had become dissolved. He also found that gastric juice, obtained by making himself vomit, had the power to dissolve meat in vitro.

In the early part of the nineteenth century (1819) Dr. William Beaumont (Fig. 8.20), an American army surgeon, was offered an opportunity, unique at that time, to study the digestive processes in

the human stomach. A French Canadian hunter, Alexis St. Martin, while at Fort Mackinac on Lake Michigan, received a severe wound in the abdomen from the accidental discharge of a musket. Part of the wall of the stomach was destroyed. Under Beaumont's care the man recovered but the large wound in the abdominal wall and the stomach failed to close completely. Through this opening Beaumont was able to observe the changes taking place in the food during the



Fig 8 20 William Beaumont,

different stages of digestion, and performed a number of simple experiments which clarified many obscure aspects of gastric physiology.

The experiments of these pioneers upon the chemical features of digestion had led men to forget for a time the importance of mechanical factors. The pendulum had swung too far. But today, the mechanical factors are recognized as playing a prominent role in gastric digestion, though this should not be taken to mean that nutriment is expressed from the food by any mechanical action.

Abnormalities of the motor mechanisms of the stom-

ach are much more often the immediate cause of digestive disorders than is a failure, or even the entire absence, of the digestive power of the gastric juice. Persons in which gastric secretion is entirely lacking (achylia gastrica) may enjoy perfect health, and suffer no digestive discomfort whatsoever. On the other hand, disorders of the movements of the stomach—e.g., too slow or too rapid evacuation of the food, or spasm of the pylorus or of the cardiac sphincter—frequently give rise to severe gastric symptoms

Vomiting. This is usually preceded by nausea. The ejection of the stomach contents is accomplished in the following manner. A strong contraction occurs in the pyloric part of the stomach, together with relaxation of the body of the stomach and the cardiac sphincter. Ac-

companying or immediately following these movements, the abdominal muscles contract forcibly, and the diaphragm descends. Thus the stomach is compressed and its contents, being prevented from passing downward by the firm contraction of the pyloric region, are forced through the relaxed cardia into the esophagus. The latter relaxes throughout its length, which permits the free passage upward of the stomach contents. The larynx is raised at the same time and its opening closed, thus preventing the passage of material into the air passages. The stomach, it will be noted, is emptied passively; that is, it is emptied from pressure applied from without. There is no evidence, in the adult at any rate, that an ascending peristaltic contraction of the gastric wall is instrumental in ejecting the stomach contents into the esophagus. In the young infant, however, a reverse peristaltic movement of this nature probably does occur. The purely passive role played by the stomach in vomiting was shown clearly by the French physiologist, Magendie. He excised the stomach of an animal, replaced it by a pig's bladder, and then gave an emetic which acted on the vomiting center; a typical vomiting act followed, fluid being forcibly expelled from the grafted bladder.

Vomiting is a reflex act and may follow irritation of neare and

Vomiting is a reflex act and may follow irritation of nerve endings (vagus or sympathetic) in the stomach or duodenum. But the stimulation of afferent fibers, especially those transmitting pain impulses, in almost any organ of the body may induce vomiting. The vomiting due to disease of the appendix or gall bladder, or to some painful injury, is well known. Afferent impulses arising in a failing heart may also cause vomiting, or the act may be induced by psychic influences—anxiety, fear, or disgust. Seasickness and other forms of motion sickness are due to the stimulation of nerve endings in the utricle of the ear (Chap. 13). Nausea or vomiting not uncommonly results from eye strain, the afferent impulses initiating the reflex arising in the eye muscles.

Drugs and other substances used in medicine for the induction of vomiting are called emetics. Among those which act upon the nerve endings in the gastric or duodenal wall are antimony tartrate (tartar emetic), copper and zinc sulphates, and salt or mustard and water. Certain other emetics, such as apomorphine, act upon the vomiting center in the medulla oblongata. It is probable, however, that the vomiting which follows the administration of emetics of the latter type is also essentially reflex in nature. The drug acts, apparently, by raising the excitability of the center; that is, the threshold of the center to stimulation is lowered, so that impulses from various parts of

the body, which normally make no impression upon it, become effective. The vomiting resulting from metabolic disturbances (e.g., in pregnancy or nephritis), or that associated with general bodily fatigue, is explained in a similar way.

# Digestion in the Intestine

The intestinal tract is divided for the purpose of description into two parts—the small intestine and the large intestine. The small intestine, though of smaller diameter, is much longer than the large intestine (see Fig 8.1). In man its length is about 21 feet. The first 10 inches or so (twelve finger breadths) are called the duodenum. The rest of the small intestine is divided rather arbitrarily into an upper portion (about two fifths) called the jejunum, and a lower somewhat narrower portion (about three fifths) called the ileum. The small intestine from





Fig 8.21 X-ray photograph of the large intestine.

the duodenum onward is attached to the posterior abdominal wall by a fan-shaped membrane called the *mesentery* which carries vessels and nerves to the intestinal wall.

All those processes, through which the food is finally prepared for absorption, are carried out in the small intestine. Amino acids and certain other products which have escaped absorption from this part of the alimentary tract undergo bacterial decomposition in the large intestine (see p. 345), but nothing of nutritional value is prepared in this part of the digestive tract. Large quantities of water are, however, absorbed from the large intestine; it converts the fluid contents of the small intestine into the pasty consistency of normal feces.

The large intestine extends from the end of the ileum to the anus. It is about 5 feet long, in man, and consists of the cecum, colon, rec-



tum, and anal canal. The cecum is the saclike dilatation at the commencement of the large intestine; it lies in the lower right-hand part of the abdominal cavity. The ileum opens into its upper end and the vermiform appendix arises 2 inches or so lower down. The colon is described in four parts—the ascending, transverse, descending, and pelvic colons. These are shown in Figure 8.1, p. 288 and Figure 8.21. The ascending colon occupies the right side of the abdomen and is continuous below with the cecum. The transverse colon crosses the abdomen and is continued into the descending colon which occupies the left side of the abdomen. The lower part of the large intestine occupies the pelvis and consists of the pelvic colon, the rectum, and the anal canal, in this order from above downwards. The interior of the cecum is shown in Figure 8.22. The bends of the colon in the regions of the liver (i.e., at the junction of the ascending and transverse colons) and spleen (at the junction of the transverse and descending colons) are called, respectively, the hepatic and splenic flexures. The internal anal sphincter is situated at the lower end of the rectum. The external anal sphincter guards the lower aperture of the anal canal. The latter opening is called the anal orifice or anus.

The pancreas is a racemose gland (p. 40). It is an elongated




Fig 8.23 Diagram to show the relation of the pancreas to the duodenum, stomach, and left kidney.

structure lying across the posterior wall of the abdomen; its right-hand broad part, called the head, fits into the curve of the duodenum (Fig 8 23). Its body and tail are directed toward the left. The cells which form its external secretion—the pancrealic juice—have an alveolar arrangement resembling that seen in the salivary gland. The alveoli or actin are drained by a system of ducts. In the human subject, the main pancreatic duct (duct of Wirsung) pierces the duodenal wall obliquely, and joins the common bile duct to form a small chamber—the ampulla of Vater. The ampulla opens into the duodenum about 31 inches below the pylorus. The pancreas contains, besides the cells which form the pancreatic juice, other rounded groups of

cells called the *islands* or *islets of Langerhans* (Fig 8.24). The islet cells are of an entirely different character; they are surrounded by the acinar (external secreting) cells, but separated from them by a narrow clear space. They are responsible for the production of insulin (p. 370).

When the gland is resting (as in a fasting animal), the alveolar cells, like the serous cells of the salivary glands, contain numerous granules




Fig. 8 24 Microscopic section of pancreatic tissue AC, alveoli of glands which secrete pancreatic juice, IN, one of the islands of Langerhans, which produce insulin

(p. 40); these furnish the enzymes of the juice. During secretion, the cells discharge their zymogen granules into the alveolar cavity, their cytoplasm, except perhaps for a narrow zone bordering the alveolar cavity, then appearing quite clear.

The pancreatic juice. Composition. The pancreatic juice is alkaline in reaction, due to its content of sodium carbonate and bicarbonate. Its principal ferments are trypsin (pancreatic protease), amylopsin (pancreatic amylase), steapsin (pancreatic lipase), and rennin.

Trypsin acts upon the proteoses and peptones formed in the peptic digestion of protein This enzyme is also capable of attacking any protein which has escaped gastric digestion, and converting it to proteoses and peptones. Trypsin breaks the peptone molecule into smaller groups of amino acids, named peptids. When the amino acid groups are relatively large, they are called polypeptids. The final

stage in the digestion of protein—that is, the complete disruption of the molecule into its individual amino acids—is brought about by the erepsin (peptidases) of the intestinal juice. But the protein molecule can be broken down into groups of four amino acids (tetrapeptids) and three (trupeptids), or even two (dipeptids) amino acids, during trytte digestion.

Juice collected directly from the pancreatic duct has very little power to digest protein, but, if allowed to come into contact with the intestinal mucosa, it at once becomes active. The inactive material in the juice is called trypsinogen; its activation to trypsin is dependent upon a constituent of the intestinal juice, named enterokinase (o. 327). Trypsinogen is also activated by calcium.

Trypsin can act only in an alkaline medium, the optimum pH being around 8.5.

Amylopsin or pancreatic amylase (also called pancreatic diastase) has an action similar to that of the ptyalin in saliva (p. 298), converting starch into maltose But pancreatic juice has a more powerful starch-splitting action than has saliva; it is also afforded a longer time in which to act. As compared with saliva, pancreatic juice is, therefore, of much greater importance in the digestion of starch. Pancreatic juice also contains some maltase which converts a part of the maltose to glucose

Steapsin, or pancreatic lipase, splits fats into their constituents—fatty acids and glycerin (glycerol). If the intestinal contents are alkaline in reaction, the liberated fatty acids may combine with alkali to form soaps The action of the steapsin upon the fat, tristearin, is shown in the following equation:

 $C_1H_1(C_1)H_{20}COO)_1 + 3H_2O = C_1H_1(OH)_2 + 3C_{11}H_{20}COOH$ testes and given stears acid  $C_1H_{10}COOH + NaOH = C_1H_{20}COONa + H_2O$ stears acid sodium stearste

Soap formation in the intestine probably occurs to a very limited extent, for soaps can form freely only in an alkaline medium; any which may have formed become precipitated again when the medium is acidified. The intestinal contents are slightly acid in reaction except when the pancreatic secretion is at its height. Even then, it is only the duodenal contents which are rendered slightly alkaline, the contents of the rest of the intestinal tract remaining slightly acid.

The optimum pH for the action of pancreatic lipase is around 8.0.

The secretion of pancreatic juice. The secretion of pancreatic juice is brought about by a hormone formed in the mucosa of the small intestine. The existence of this hormone was demonstrated in 1902 by Bayliss and Starling. They found that a hydrochloric acid extract of the duodenal mucosa, when injected intravenously, caused the secretion of pancreatic juice. The intravenous injection of the acid by itself was ineffective. On the other hand, the introduction of hydrochloric acid into a loop of the bowel, the nerves of which had been sectioned and thus isolated from all connections with the pancreas except through the blood stream, was followed by a secretion of pancreatic juice. It was therefore concluded that in the normal course of digestion the acid chyme, after reaching the duodenum, acted upon the duodenal mucosa to cause the formation and absorption into the blood stream of a hormone which excited the pancreatic cells. The hormone was named secretin.4 Later work has shown that extracts of the intestinal mucosa made with solvents other than acid (namely, water or alcohol) are active. It has also been demonstrated that bile introduced into the duodenum causes secretin to be absorbed into the blood stream and induces, in consequence, a secretion of pancreatic juice.

A second hormone having a stimulating action upon the secretion of pancreatic juice has been discovered within recent years It has been named pancreozymin and, like secretin, is obtained by extraction from the duodenal mucosa But, in contrast to the latter hormone, which increases only the water and salts of the juice, pancreozymin increases the production of the enzymes It causes the discharge of zymogen granules from the gland cells and has little effect upon the volume (water) of the secretion.

The secretion of pancreatic juice is also brought about through nerve impulses. Stimulation of the vagus nerve causes the secretion of a juice very rich in digestive enzymes; rapid exhaustion of the zymogen granules results. Secretin, as just mentioned, appears to be responsible mainly for the secretion of the water and inorganic constituents of the juice. The vagal effect upon pancreatic secretion is, therefore, closely similar to that caused by pancreozymin and unlike that following the injection of secretin. It was shown several years ago by Pavlov that psychic factors—the taste and smell of food—were capable of evoking a secretion of pancreatic juice. These reflex secretory effects were particularly well shown by feeding experiments

<sup>&</sup>lt;sup>4</sup> It is interesting to recall that Bayliss and Starling coined the word hormone to express the characters of their newly discovered secretin

in dogs. However, the psychic or cephalic element plays a much less prominent role in pancreatic than in gastric secretion.

The villi and glands of the small intestine. The mucous membrane of the small intestine is beset with great numbers of minute, slender, fingerlike or tonguelike processes—the intestinal villi—which give to the naked eye the soft velvety appearance of the intestinal mucosa (Fig. 8.25). Absorption from the small intestine of water and the




Fig 8 25 Showing vills of the small intestine. One villus sectioned lengthwise to show yessels.

products of digestion occurs entirely through the specialized epithelium which covers the villi. The columnar epithelial cells in this situation are continuous with those lining the intestinal glands and show short delicate lines lying along and perpendicular to their free borders. In this respect these cells resemble those lining the proximal convoluted tubules of the kidney (p. 270). The striations appear to be associated in some way with the capacity of selective absorption which both the renal and intestinal cells possess.


Intestinal glands. The crypts of Lieberkithn or intestinal glands are tiny pockets lying between the villi. Their blind ends dip deeply into the mucous membrane, reaching as far as the muscularis mucosae (Figs. 8.25 and 8.36). They are lined by epithelial cells continuous with those covering the summits of the villi. Many of the epithelial cells lining the gland become converted into goblet cells and discharge mucus. Others (cells of Paneth) situated at the deepest or blind end of the crypts are filled with coarse granules and a believed to secrete the enzymes of the intestinal pince (Fig. 8.26).

The intestinal juice, succus entericus. The term intestinal juice

(L. succus entericus) is reserved for the secretion of the intestinal glands or crypts of Lieberkuhn described in the preceding section and does not mean merely the fluid contents of the intestinal canal.

Composition. The intestinal juice is alkaline in reaction, due to its content of sodium carbonate and bicarbonate. It contains enterokinase, the activator of trypsin, and the following enzymes.

- Peptidases (erepsin),<sup>8</sup> enzymes which effect the final breakdown of the protein molecule into its constituent amino acids
- Sucrase, maltase, and lactase, enzymes which act upon the disaccha-



3. Lipase.




Fig 8 26 Showing a crypt of Lieberkühn on left with goblet cells. On right, an enlarged sketch of bottom of crypt to show cells of Paneth loaded with granules

Peptidases do not act upon unchanged protein. The protein must first be digested to the peptid stage by trypsin. Peptidases act best at a pH of about 8. The intestinal wall itself contains these enzymes and small amounts are present in other tissues of the body, including the blood. The intestinal wall also contains enzymes—nuclease, nucleotidase, and nucleosidase—capable of digesting nucleic acid and its derivatives (p. 366).


Sucrase, maltase, and lactase. Sucrose (cane, beet, and maple sugar) is split by sucrase into a molecule each of glucose and fructose; maltose, when acted upon by maltase, yields two molecules of glucose; and lactose is converted into a molecule each of glucose and

<sup>5</sup> Erepsin (Gr ereptomar, to feed upon) is an old term applied to the enzymes of the intestinal juice which effect the final breakdown of the protein molecule to amno acids in the belief that but a single enzyme was responsible.

in dogs. However, the psychic or cephalic element plays a much less

prominent role in pancreatic than in gastric secretion.

The villi and glands of the small intestine. The mucous membrane of the small intestine is beset with great numbers of minute, slender, fingerlike or tonguelike processes-the intestinal villi-which give to the naked eye the soft velvety appearance of the intestinal mucosa (Fig. 8.25). Absorption from the small intestine of water and the



villi of the small intesune. One villus sectioned lengthwise to show vessels.

products of digestion occurs entirely through the specialized epithelium which covers the villi. The columnar epithelial cells in this situation are continuous with those lining the intestinal glands and show short delicate lines lying along and perpendicular to their free borders. In this respect these cells resemble those lining the proximal convoluted tubules of the kidney (p. 270). The striations appear to be associated in some way with the capacity of selective absorp-tion which both the renal and intestinal cells possess.

Intestinal glands. The crypts of Lieberktuhn or intestinal glands are tiny pockets lying between the villi. Their blind ends dip deeply into the mucous membrane, reaching as far as the muscularis mucosae (Figs. 8.25 and 8.36). They are lined by epithelial cells continuous with those covering the summits of the villi. Many of the epithelial cells lining the gland become converted into goblet cells and discharge mucus. Others (cells of Paneth) situated at the deepest or blind end of the crypts are filled with coarse granules and are believed to secrete the enzymes of the intestinal juice (Fig. 8.26).

The intestinal mice, succus entericus. The term intestinal juice

(L. succus entericus) is reserved for the secretion of the intestinal glands or crypts of Lieberkuhn described in the preceding section and does not mean merely the fluid contents of the intestinal canal.

Composition. The intestinal juice is alkaline in reaction. due to its content of sodium carbonate and bicarbonate. It contains enterokinase, the activator of trypsin, and the following enzymes.

- 1. Peptidases (erepsin),5 enzymes which effect the final breakdown of the protein molecule into its constituent amino acids.
- 2. Sucrase, maltase, and lactase, enzymes which act upon the disaccha-(milk sugar), respectively.
- Lipase.

COBLET CELL DISCHARGING MUCUS PANETH

Fig. 8 26 Showing a crypt of Lieberkuhn on left with goblet cells. On right, an enlarged sketch of bottom of crypt to show cells of Paneth loaded with grannifes

rides, sucrose (cane sugar), maltose (sugar of malt), lactose

Peptidases do not act upon unchanged protein The protein must first be digested to the peptid stage by trypsin. Peptidases act best at a pH of about 8. The intestinal wall itself contains these enzymes and small amounts are present in other tissues of the body, including the blood. The intestinal wall also contains enzymes-nuclease. nucleotidase, and nucleosidase-capable of digesting nucleic acid and its derivatives (p. 366).

Sucrase, maltase, and lactase. Sucrose (cane, beet, and maple sugar) is split by sucrase into a molecule each of glucose and fructose; maltose, when acted upon by maltase, yields two molecules of glucose; and lactose is converted into a molecule each of glucose and

Erepsin (Gr ereptomal, to feed upon) is an old term applied to the enzymes of the intestinal juice which effect the final breakdown of the protein molecule to amino acids in the belief that but a single enzyme was responsible

DIGESTION

galactose by lactase. The hydrolysis of sucrose is shown in the following equation:

 $C_{12}H_{22}O_{11} + H_2O = C_6H_{22}O_6 + C_4H_{12}O_6$ 

Lipase, though in lower concentration in the succus entericus than in pancreatic juice, plays an important role, nevertheless, in the digestion of fat. The importance of its action is shown after abolishing the effect of the pancreatic lipase, as by tying the pancreatic duct. Following this procedure, at least 70 percent of the fat in the food undergoes digestion.

The secretion of intestinal juice. Mechanical types of stimulation are particularly effective in causing a secretion of intestinal juice. The contact with the intestinal junces of rough indigestible constituents of the food and especially distension of the intestinal wall itself by food masses exert a pronounced excitatory effect upon the intestinal glands. The secretion is brought about reflexly through the nerve plexuses of the intestinal wall. The extrinsic nerves (vagus or sympathetic) do not appear to have any excitatory effect. If either of these nerves exert any effect at all upon secretion, it is of an inhibitory nature. The question whether or not a hormone plays a role in controlling secretion cannot be answered definitely, though there is some evidence that secretin stimulates the intestinal glands as well as the pancreas.

# The Production of Bile by the Liver. The Composition of Bile; Its Storage in, and Expulsion from, the Gall Bladder

Though it is customary to speak of the secretion of bile, this fluid, like urine, is rather in the nature of an excretion. It is produced from the constituents of the blood by the cells of the liver which are arranged in rows—the liver cords—radiating from a central vein (p. 208). A small aggregation of such cords, together with the central vein, is known as an hepatic (liver) lobule. On one side of each liver cord is a blood sinus; through these vessels, called hepatic sinusoids, courses a mixture of portal and hepatic artery blood (Figs. 8.27 and 8.28). On the other side lies a fine canal (biliary canaliculus) into which the cells of the liver cords discharge the bile formed from the constituents of the blood. The canaliculi drain into channels of progressively increasing size until a relatively large tube—the hepatic duct—is formed, along which the bile leaves the liver.

The composition of bile. Human bile is a clear yellow or orangecolored fluid secreted by the liver cells. Its chief constituents are the bile pigments, the bile salts, lecithin, cholesterol, inorganic salts, and mucin. Table 8-3 gives the average composition of human bile as it is secreted by the liver.

## Table 8-3

Water	974.80
Solids	25.20
Mucin and pigments	5.30
Bile salts	9.30
Fatty acids from soaps	1.23
Cholesterol	0 63
Lecithin )	0.00
Lecithin Fat	0.22
Inorganic salts	8.32

Bile collected from the gall bladder is much more concentrated than liver bile (see p. 332).

The bile pigments are called bilirubin and biliverdin. They are derived from the hemoglobin liberated from red blood corpuscles, which at all times are undergoing disintegration in the blood stream (p. 329). The conversion of the hemoglobin to bilirubin is effected by the reticulo-endothelial cells of the spleen, liver (Kupffer cells), bone marrow, and general connective tissues (p 94) The pigments con-

sist of the porphyrin part of the hemoglobin molecule—that is, the remnant of the molecule after the globin and iron have been removed. Bilurubin is not quite identical, however, with the porphyrin in hemoglobin, for the chemical structure of the porphyrin group is also altered. The iron liberated in the conversion of hemoglobin to bile pigment is stored chiefly in the liver and spleen.

Bilirubin (C<sub>23</sub>H<sub>26</sub>N<sub>4</sub>O<sub>4</sub>) is an orange-red pigment; it is the chief pigment in human bile and in the bile of dogs. Biliverdin




Fig 8.27 Diagram of a liver lobule. 1, interlobular vein; 2, central vein; 3, a lobule. The fine lines converging toward the central vein and surrounding the liver cells are the hepatic sinusoids.





Fig 8.28 Segment of a liver lobule (diagrammatic) White circles represent blood corpuscles Arrows indicate bile secretion (See also Fig 8.27)

(C<sub>2</sub>,H<sub>2</sub>,N<sub>2</sub>O<sub>2</sub>) is green and constitutes the greater part of the pigment in the bile of birds and of herbivorous mammals. It is simply an oxidation product of bilirubin

Upon reaching the intestine, bilirubin undergoes chemical reduction as a result of bacterial action. This reduction product, which gives the yellow brown color to feces, is called stercobilinogen (also called urobilinogen). Upon exposure to air, a part of the stercobilinogen is oxidized again, and is then called stercobilin (or urobilin). Though the greater part of the stercobilinogen is passed with the feces, a part is absorbed into the portal circulation and returned to the liver, whence it is discharged again in the bile (see Fig. 8.29). In health, only minute amounts of urobilinogen, or none at all, find their

way into the general circulation. Any which does escape from the liver into the systemic blood is excreted by the kidney, but the amount is so small that it does not stain the urine; it can be detected only by delicate chemical tests. The yellow color of healthy urine is not due, therefore, to this pigment (see p. 274).

Fig 8.29 Diagram illustrating the circulation of bite pigment. 1, liver; 2, intestine; 3, kidney. The dotted line indicates the entrance of uroblimogen into the blood and its excretion by the kidney when the liver is damaged or there is an excess of bile pigment producfion



The bile salts are sodium glycocholate and sodium taurocholate. The bile acids glycocholic and taurocholic are produced in the liver by the union of cholic acid  $(C_{21}H_{10}O_2)$  with the amino acids, glycine and taurine, respectively.

The secretion of bile. We have seen that the cells forming the liver cords are exposed on one side to the blood in the sinusoids from which materials are removed and passed as bile into the biliary canaliculi on the other side (Fig. 8.28). The bile is carried through a system of channels of progressively greater size to the large hepatic duct. The liver forms and secretes bile continuously, but, instead of passing directly into the intestine as it is formed, the bile passes into the gall bladder from which it is discharged from time to time into the duodenum.

The storage of bile and its discharge into the intestine. The gall bladder is a small pear-shaped sac which, in the human subject, has a capacity of about 50 cc. It is situated on the under (posterior) surface of the liver (Fig. 8.30). Leading from its upper and smaller end is the cystic duct. The cystic duct joins the hepatic duct at an acute angle to form the common bile duct. The latter joins the pancreatic

forms important functions in the intestine. Its main function is concerned with the digestion and absorption of fat Though the bile does not contain a fat-splitting enzyme, and, therefore, has no direct action upon the digestion of fat, the bile salts, through their power to lower surface tension, aid very greatly in the emulsification of fatty materials in the intestinal contents. The digestion of fat is materially advanced thereby, for the division of the oily materials into small globules increases by several fold surface exposed to the action of the pancreatic and intestinal lipases. The bile salts have also the property of rendering soluble, in taqueous fluids of the intestine, the fatty acids liberated during fat digestion. This is the so called hydrotropic action of the bile salts. The dissolved fatty acids, in turn, have the property of lowering surface tension, any soaps which may be formed during digestion, and the cholesterol of the bile, have a similar action These several factors combine to produce a fine emulsion of fatty material in the small intestine. The bile salts aid fat digestion in another way They serve as specific activators of the pancreatic lipase. That is, quite apart from their emulsifying effect, they enhance very greatly the fatsplitting action of the pancreatic juice. The action of the intestinal lipase is not altered in this specific way by bile

Bile, important though it is for the efficient digestion of fat, is of still greater importance for fat absorption. When bile is excluded from the intestine, over 80 percent of the fat is digested-that is, split into fatty acids and glycerin (glycerol) A large proportion of the fatty acids, however, is not absorbed, but appears in the feces. Only recently has the part played by the bile salts in the absorption of fat been elucidated They unite in the intestine with the fatty acids to form complex compounds which pass readily into the epithelial cells covering the intestinal villi. Here the bile salts are freed again from the union, and are carried in the portal blood to the liver which excretes them again in the bile This circulation of the bile salts has been known for a long time, though its significance was not apparent. The fatty acids liberated from their combination with the bile salts now combine with glycerm which has penetrated into the epithelial cells from the intestinal lumen The products of fat digestion-fatty acids and glycerin-are thus reunited within the epithelial cells to form neutral fat. The greater part (about 60 percent) of the synthesized fat passes into the small lymphatics (lacteals) running through the centers of the vill. The course taken by the remaining 40 percent of the fat is uncertain. It is probably absorbed into the blood of the portal system. There is evidence for the view that a part of the fat is absorbed directly—that is, as fine globules of unsplit fat.

Jaundice (Fr. jaune, yellow) or ieterus. In health, human plasma contains a small quantity of bilirubin. The concentration ranges in different persons from 0.1 to 0.5 mg per 100 cc. The bilirubin of the plasma is that which, having been formed by the reticulo-endothelial cells from hemoglobin derived from disintegrated erythrocytes, is on its way to the liver for excretion.

When excessive amounts of bilirubin are present in the plasma, the skin, mucous membranes, and whites of the eyes are stained yellow. The condition is called jaundice or leterus. Bilirubin then appears also in the urine and sweat, but does not pass into the saliva. An abnormally high concentration of plasma bilirubin may be due to excessive amounts of the pigment being produced, the liver and excretory channels being normal. On the other hand, the quantity of pigment formed in the body may be within normal limits, but the function of the liver is impaired as a result of some toxic or infective process, and does not remove bilirubin readily from the plasma. Again, the bile passages may be obstructed; the excretion of bile into the intestine is then prevented. There are, therefore, three types of jaundice: (1) hemolytic, (2) toxic or infective, and (3) obstructive.

Hemolytic jaundice is seen in conditions which cause increased destruction of red cells—e.g., pernicious anemia, hemolytic anemia, and poisoning with certain hemolytic agents. It is therefore due to the exaggeration of a normal process—namely, the liberation of hemoglobin from disintegrated blood cells. This type of jaundice is not uncommon in the newborn infant; it persists for 4 or 5 days after birth. The baby comes into the world with an excess of red cells, which undergo destruction during the first few days, the concentration of bilirubin in the plasma being thus increased to the point at which jaundice appears. Another factor is the functional immaturity of the liver. This type of jaundice of the new born is harmless, and indeed may be looked upon as a physiological phenomenon. In the hemolytic types of jaundice, the stools are dark, due to the increased pigment excretion: urobilin may appear in the urine.

In the toxic and infective types of jaundice, the liver cells are injured by some poison or infective process which interferes with their ability to eliminate bile pigment.

Obstructive jaundice is most often the result of blockage of the common bile duct by a gallstone within its lumen, or by a tumor pressing upon it from without. In this type, the discoloration of the skin and mucous membranes is usually intense. The feces are pale, being usually described as "clay colored," because bile pigment is, of course, excluded from the intestinal tract. The urine is usually deeply colored with bilirubin.

Movements of the small intestine. Three types of movement occur in the small intestine: (a) peristaltic, (b) segmenting, and (c) pendular



В



Fig. 8 32 Diagram of the movements of the small intestine. A, peristalsis; arrows indicate the direction of movement of the constricting band; B, segmenting movements.

Peristaltic movement (peristalsis). This kind of movement is not restricted to the small intestine, but is characteristic of hollow muscular organs in general. It has been mentioned as occurring in the esophagus during the third stage of swallowing, and in the pyloric part of the stomach. It is also seen in the large intestine, as well as in the—ureter, Fallopian tube, and common bile duct. The movement consists of a ringlike contraction of the muscular wall of the tube. The answard contraction, as it travels downward, causes a certain degree of constriction of the bowel which sweeps before it any material within

the lumen (Fig. 8.32A). The movement is readily started by a mechanical, an electrical, or a chemical stimulus. The most powerful stimulus is distension of the bowel—i.e., stretching of the muscle fibers. Filling of the bowel with food material provides just such a stimulus, the contraction occurring immediately behind the stimulated region. Peristaltic movements may follow one another at regular and fairly frequent intervals, the intestinal contents being moved steadily, but gently along the canal. At other times a brisk peristaltic contraction appears which travels rapidly along the tube but, after moving the food a considerable distance, may not be repeated for some time. This is called the peristaltic rush. Especially strong stimulation, the action of a cathartic or an irritant poison, for example, may set up a peristaltic rush which travels throughout the length of the small bowel. Such a movement may also be initiated reflexly from the esophagus during swallowing, or from the stomach.

Except in the duodenum near its commencement and at the terminal part of the ileum (see Fig. 8.1, p. 288), the peristaltic movements of the normal small intestine are always from the stomach downward—that is, in a direction toward the anus. Peristalsis in the reverse direction (antiperistalsis) is occasionally observed normally in the situations just mentioned. It may also occur anywhere in the small intestine in abnormal states of the intestinal tract, especially when the canal is completely obstructed.

Segmenting movements. These movements are rhythmical constrictions of the intestine which serve to break up and knead the food, to mix it thoroughly with the intestinal juices, and to bring the intestinal contents in contact with fresh absorptive surfaces. These movements

Fig. 8.33 Diagram showing the effect of the segmenting movements upon the food mass in the intestine. A senes of constrictions of the bowel wall suddenly divides the column of food, 1, into a number of segments, 2. Each of the latter is bisected again the next instant, the adjacent halves fusing as in 3, and being redivided as treatment with the food in the combine mix.



fusing as in 3, and being redivided as in 4. The process is repeated over and over again until the food is thoroughly mixed. (After Cannon.)

also increase the blood and lymph flow in the intestinal wall; but they exert no propulsive action upon the food. They occur at the rate of from 20 to 30 per minute (see Figs. 8.32B and 8.33).

Pendular movements. These are simple constrictions of the intestinal wall. They move up and down the bowel for short distances, giv-

ing a to and fro movement to the semifluid material in the canal, and thus lending further aid to the processes of digestion and absorption (Fig. 8.34).

The Innervation of the small intestine. The small intestine receives fibers from the vagus (parasympathetic) and the splanchnic (sympathetic) nerves. The vagus is excitatory, the sympathetic inhibitory, in action (see Pl. 8A). It will be noted that the actions of these nerves




Fig. 8.34 Pendular movement; arrows indicate the directions of the movement.

upon the intestine, as compared with their actions upon the heart, are reversed. The intestunal wall itself contains two nerve plexuses. One of these lies between the two muscular coats (circular and longitudinal) of the intestine and is called Auerbach's plexus; ganglion cells are found among the network of nerve fibers. The other plexus, known as Messner's plexus, lies in the submucosa (Pt. 8B). The vagal fibers connect with the ganglion cells of Auerbach's plexus; the impulses are thence conveyed by the axons of these cells (postganglionic fibers) to the muscle fibers The sympathetic fibers, on the other hand, do not form junctions with ganglion cells in the intestinal wall but pass directly to the muscle cells. The postganglionic sympathetic fibers arise from cells in the celac and superior mesenteric ganglia (Pt. 8A).

Though the intestinal movements are influenced by impulses traveling over the vagus and splanchnic nerves, they are not dependent upon them, for the three types of intestinal movement continue after all nerves of extrinsic origin have been sectioned. The peristaltic movements are then carried out through the intrinsic nervous mechanism (plexuses of Auerbach and Meissner). The segmenting and pendular movements are dependent upon the property of rhythmical activity of

the muscle fibers themselves, and not upon any nervous mechanism. They continue in a denervated strip of intestinal muscle, or after paralysis of the plexus by cocaine.

The ileocolic sphincter. The small intestine empties into the upper part of the cecum through a valvelike opening called the ileocolic valve (Figs. 8.35 and 8.22, p. 321). The ring of muscle surrounding the opening is called the ileocolic sphincter. It controls the passage of the contents of the ileum into the cecum and prevents or hinders the reflux of material into the ileum, thus protecting the small intestine from contamination by the putrefactive bacteria which flourish in the large intestine. The sphincter opens and closes during digestion to permit gushes of ileal contents to enter the cecum. During fasting it remains closed for long periods, but opens within a few midicates a reflex action intuited by the entra



Fig. 8 35 Diagram of the ileocolic valve in section. (See also Fig. 8 22, page 321)

the ecenth. During hasting it remains closed for long periods, but opens within a few minutes after a meal. This indicates a reflex action initiated by the entrance of food into the stomach. The ileocolic sphincter receives motor fibers through the sympathetic; inhibitory fibers are believed to be carried by the vagus.

The innervation of the ileocolic sphincter is, therefore, the reverse of that of the small intestine itself. A peristaltic wave in the small intestine, accompanied by relaxation of the sphincter, is brought about by vagal impulses; the ileal contents are passed into the large intestine. Relaxation of the wall of the small intestine, with closure of the sphincter, is caused by sympathetic impulses; the entrance of material into the large intestine is prevented. A discharge of impulses from a single nervous center (vagal or sympathetic) in this way either drives the semisolid contents of the ileum through the sphincter, or holds them back; thus incomplete digestion of the food is guarded against.

The movements and innervation of the large intestine. Food material reaches the eccum about 4½ hours after a meal. Rhythmical movements resembling the segmenting movements in the small intestine may

### 340 DIGESTION

be seen occasionally in the cecum, but peristaltic contractions are absent. Regular and frequently recurring peristaltic contractions are usually absent also from the rest of the large intestine. At certain long intervals, however, a strong peristaltic wave commencing in the upper end of the ascending colon moves swiftly over the transverse colon. This mass movement, as it is called, is analogous to the peristaltic rush in the small intestine; it empties the transverse colon, sweeping the material into the descending and pelvic colons or into the rectum. The pelvic colon and lower part of the descending colon serve as a storehouse for the feces until the desire to defectate is aroused.

The eccum, ascending colon, and the first third or so of the transverse colon are supplied with motor fibers through the vagus nerves. The remainder of the large intestine, including the rectum and the anal canal, is supplied with motor fibers through the pelvic nerves (parasympathetic from the second, third, and fourth sacral segments of the spinal cord). The inhibitory fibers to the entire large intestine are derived from the sympathetic These two sets of nerves act upon the internal anal sphincter in a reverse fashion, the sympathetic is excitatory, the pelvic nerve inhibitory. As stated previously, the sympathetic is also the motor nerve to the ileocolic sphincter. The external anal sphincter is under voluntary control through the pudendal nerve.

In health, the antagonistic effects of the parasympathetic and sympathetic nerves to the large intestine are nicely balanced. But the balance is sometimes upset Relaxation and weakness of the walls of the descending and pelvic colons result from overactivity of the sympathetic innervation. The tone of the internal anal sphinicter, on the other hand, is exaggerated. The colon is unable to empty itself effectually, and dilatation of its lumen, often to enormous proportions, follows. This condition is called megacolon (G. megas, large) or Hirschspring's disease. When, on the other hand, the petic nerve (parasympathetic) is hyperactive, the descending colon becomes firmly contracted; its lumen is obliterated. This—the so-called spastic colon—is one cause of chromic constitution (p. 342).

—is one cause of arrome consupation (p. 342).

Defectation (evacuation of the bowels) is a reflex act initiated by the passage of feces into the rectum. Normally the rectum is empty, except just before defecation; feces are forced into it from the pelvic colon by a mass movement. The latter movement is often started by the entrance of food into the stomach—gastrocolic reflex. This reflex is responsible for the desire to defecate so often experienced a short time after a meal, especially breakfast. The distension of the rectum as it becomes filled with feces acts as a stimulus to afferent nerve end-

ings in the rectal wall. The impulses set up are conveyed to a center in the sacral part of the spinal cord. The efferent nerve fibers pass to the wall of the descending and pelvic colons, rectum, and the internal anal sphincter, via the pelvic nerves, and to the external anal sphincter and the striated muscle lying in relation to the rectum, via the pudendal nerves. A higher center for defecation is situated in the medulla oblongata. This is connected to the lower center by tracts of fibers in the spinal cord. The intrarectal pressure necessary to start the defecation reflex is from 40 to 50 mm Hg.

The movement of defecation consists of a powerful peristaltic contraction of the descending and pelvic colons and rectum, assisted usually by a voluntary contraction of the abdominal muscles, and of the muscles lying in relation to the rectum (levatores ani and rectococygeus muscles). The contraction of the abdominal muscles is immediately preceded by an inspiratory movement (descent of the diaphragm and closure of the opening [glottis] of the larynx), thus raising the intra-abdominal pressure Relaxation of the anal sphincters occurs reciprocally with the contraction of the intestinal wall—a nervous mechanism which we have already dealt with.

Constipation. Persons vary considerably in the frequency with which their bowels are evacuated. In some a bowel movement occurs twice or even three times daily, while others may feel no discomfort if an interval of two days or more elapses between movements. The majority of healthy persons have one evacuation daily, usually in the morning after breakfast. Owing to this variability, it is difficult to give a precise definition of constipation However, when the interval between bowel movements is greater than 24 hours and the subject, as a result, suffers distress or discomfort (e.g., headaches, digestive disturbances, etc.), or if the feces are abnormally dry and hard and the evacuation of the bowels difficult, constipation certainly exists.

Constipation is caused most commonly by bad habits. As mentioned above, the desire to empty the bowel is aroused by the passage of feces into the rectum, and the stimulation of afferent nerve endings in the intestinal wall. The act can, however, be voluntarily restrained and, when this is practiced, the tone of the rectal wall is reduced, and the rectum thus accommodates its capacity to the bulk of the feces (postural tone, p. 293); the afferent nerve endings are no longer adequately stimulated, or, at any rate, are unresponsive, and the desire to defecate passes As a result of the absorption of water, the retained feces become dry and hard. When the habit of postponing defecation in this way is persisted in, the rectum, which normally,

Other causes of constipation are: (a) A diet which leaves too little unabsorbed residue or one which contains too little fluid. The contents of the large intestine are, therefore, of small bulk and fail to furnish a sufficiently strong stimulus (stretching of the intestinal wall) to set up vigorous peristaltic contractions (b) A colon which absorbs too readity and thus causes undue drying of the feces (c) Hypertonic state of the muscle of the colon—spastic constipation; the transverse and descending colons are the seat of a strong tonic contraction which impedes the progress of the feces.

The myth of autointoxication. The general effects of constipation upon the sense of well-being are too familiar to require description.

But their cause is a subject upon which there is much misunderstanding. We hear a great deal of poisons formed in the intestinal tract, and the dire effects which they are supposed to have upon the body. Intestinal into ucation or automtoxication (self-poisoning) is glibly spoken of, and advertisements in the daily press exhort one to irrigate the colon, in order to remove the noxious materials. There is no doubt whatever that small amounts of powerful poisons are formed in the large intestine as a result of the action of bacteria upon amino acids which have escaped absorption in the small intestine. They are intensely toxic if injected into the circulation. Some of these substances -known generally as amines-are formed by the removal of a molecule of carbon dioxide from such amino acids (p. 360) as alanine, tyrosine, histidine, etc. Among the more potent of such amines are ethylamine (from alanine), histamine (from histidine), and tyramine ethylamine (from alanine), histamine (from histidine), and pranine (from tyrosine) However, the production of such poisons is a perfectly normal process, and there is nothing more certain than that they cannot be held responsible for the headache, bad breath, furred tongue, or any of the other effects of constipation. The body is provided with mechanisms for tendering innocuous the poisons formed in the colon. The first line of defense raised against them is in the wall of the bowe tuelf; here some are destroyed or changed into harmless compounds. The second and most important detoxicating mechanism is in the liver, where the toxic bodies are combined with sulfuric acid or with glycuronic acid. The resulting relatively innocuous compounds (indoxyl sulfuric acid and glycuronates) pass from the liver to the kidney for excretion.

To what then are the symptoms of constipation due? It is now generally conceded that they are of reflex origin. Afferent impulses set up in the distended colon or rectum, though ineffectual in precipitating the defecation reflex, produce reflex effects in other parts of the body, particularly the stomach and blood vessels. The headache is probably due to the effect upon the intracranial blood vessels. The rapid relief from the ill effects of constipation which follows evacuation of the bowels is a common experience, and, in itself, argues strongly against such effects being of toxic origin, for it is inconceivable that poisons could be freed so swiftly from the blood stream. To quote the pertinent comment of Dr. Alvarez, who has devoted much time to a study of the question, "A drunken man does not at once become sober when the whisky bottle is taken from him"; one might add, "and even after his stomach has been emptied."

# Absorption from the Intestinal Tract

The absorption of food products is, for all practical purposes, confined to the small intestine. Some drugs and certain hormone preparations can be absorbed in minute quantities from the mouth, especially the mucous membrane beneath the tongue. Alcohol and a number of drugs are absorbed from the stomach fairly rapidly. The gastric absorption of water and, to some extent, of mineral salts, also occurs, but much more slowly. The absorption from the stomach of glucose and other sugars in weak solution is negligible, but may be considerable if in a concentration of over 10 percent. The greatest absorption of water relative to solids occurs from the large intestine. As a result, the intestinal contents undergo a pronounced reduction in bulk, and a change in consistency in their passage through this part of the intestinal tract. Unchanged protein is not absorbed, except occasionally in minute amounts, from any part of the gastrointestinal tract. Fat does not pass from the cavity of the stomach into the circulation, though an attempt at absorption appears to be made; for after a meal of fat, fine droplets of oil can be detected in the gastric mucosa itself.

The alimentary tract, from a physiological point of view, must be

<sup>&</sup>lt;sup>6</sup> The ingestion of a large quantity of raw white of egg may be followed, even in a healthy person, by its appearance, unchanged, in small amounts in the urine.

considered as lying outside of the body proper. Food material ingested and held within the digestive tube cannot be said truly to have entered the body until it has passed across the intestinal mucosa and been absorbed into the blood. Just as the skin covers the outer surface of the




Fig 8 36 Interior of villus A, arteriole, C, blood vessels, D, lymph vessel, E, epithelium, F, crypt of Lieberkühn, L, lacteal, V, venule.

body, so the mucosa of the gastrointestinal tract constitutes its inner surface. An examination of the interior of the small intestine shows how well this part of the lining of the body has been fashioned for increasing the area exposed for the absorption of food materials. The mucosa is raised into circular folds (plicae circulares) which, in the upper part of the intestine, may be nearly a third of an inch in depth. These and the intestinal villi (p. 327), especially the latter, increase enormously the total absorbing surface (see Fig. 8.36). It has been estimated that the number of villi in the human small intestine is around 5,000,000 and the absorbing surface not far short of 10 square meters-more than five times the skin surface. The center of each villus is occupied by an arteriole, a venule, and a lymph vessel or lacteal. The arteriole leads into a network of capillaries lying just beneath the epithelial covering of the villus. The central vessels are surrounded by

areolar tissue together with bundles of smooth muscle fibers. If the intestinal mucosa of a living animal is examined with the low power of the microscope, the villi will be found to be in ceaseless motion, swaying or lashing from side to side, and lengthening and shortening alternately. These movements, by their constant agitation of the intestinal fluids in the immediate neighborhood of the villi, cause thorough mixing with the digestive enzymes and bring the food material into contact with fresh absorbing surfaces; they thus aid very materially the digestive and absorptive processes.

It will be recalled that the intestinal mucosa is impermeable, or nearly so, to the large molecules of the three foodstuffs—carbohydrates (starch and the disaccharides, cane sugar, maltose, and lactose), proteins, and fats—but permits the free passage of the smaller molecules—glucose, amino acids, fatty acids, and glycerol into which the food is split by enzyme action. The absorption of these products is not, however, a simple process of diffusion, in which the intestinal mucosa

acts merely as a passive membrane. On the contrary, the epithelial cells covering the villi take an active part in absorption, as is evidenced by the observation that an increase in oxygen consumption and in carbon dioxide production accompanies the process. Furthermore, the absorption, as shown by the following experiment, is selective. When the three sugars—glucose, galactose, and fructose—in equal concentrations are placed in a loop of bowel immersed in saline, glucose passes through the intestinal wall more rapidly than does galactose, and galactose more rapidly than fructose, provided that the intestinal mucosa remains viable and uninjured. After injury or death of the mucosa, the three sugars pass through it at equal rates—i.e., simply by diffusion.

After passing through the epithelium of the villi, the glucose and amino acids are absorbed into the network of blood vessels mentioned above; fat, synthesized in the epithelial cells from fatty acids and glycerin, passes, for the most part, into the lymph channels.

The reaction of the intestinal contents. The formation of feces. The reaction of the duodenal fluids is usually slightly acid but it depends upon the stage of gastric digestion at which a sample is taken; upon the degree of acidity of the gastric contents; upon the quantity of alkaline juices, pancreatic juice, and mucus secreted into the duodenum; and upon the nature of the food. It may be definitely alkaline. The contents of the remainder of the small intestine and of the entire large intestine have a reaction a little on the acid side (a pH of about 6.8). The feces are very faintly acid as a rule.

The feces are not simply and solely unabsorbed residues of the food, but are made up largely of materials excreted from the blood. During starvation, for example, the bulk of the feces may not be greatly less than at ordinary times, and a loop of intestine isolated from the rest of the intestine becomes filled after a few days with a pasty mass indistinguishable in consistency and composition from ordinary feces except that, of course, being isolated from the rest of the intestinal tract, it contains no bile and is, therefore, pale. Bacteria make up about 9 percent of the feces; the other main solid constituents are food residues, which vary considerably in amount with the proportion of indigestible material (chiefly cellulose) in the diet. They contain some fat, nitrogenous substances, and minerals eliminated from the blood, together with epithelial cells and leucocytes, shed from the intestinal mucosa. A very small proportion of digestible food appears in the feces. In other words, practically all the protein, fat, and carbohydrate which is eaten is absorbed, the food residues of the feces consisting al-

most entirely of indigestible substances. Vegetable material, since its framework is composed of cellulose, contributes more to the feces than do other foods. This indigestible material or "roughage," as it is commonly termed, serves a useful purpose in that it acts as a mechanical stimulus, increasing the motility as well as the secretions of the intestinal wall.

Substances that cause evacuation of the bowels: laxatives, cathartics, and purgatives. The manifold drugs and other substances em-ployed to stimulate the intestinal movements and cause evacuation of the bowels do not all bring about their effect after the same fashion. Some, such as Epsom salts (magnesium sulfate), sodium sulfate, and other saline cathartics, being not absorbable, cause through their osmotic effect a flow of water into the small intestine which, by increasing the bulk of the intestinal contents, causes distention, and consequent stretching of the bowel wall; this, we have seen (p. 337), acts as a powerful stimulus to peristalsis. Other cathartics, such as castor oil, contain an irritant principle which serves to excite intestinal motility. Croton oil is an especially powerful irritant which in very small dosage causes profuse purgation and, in larger amounts, may induce acute inflammation of the intestinal mucosa. Figs, raisins, etc., act by virtue of their indigestible residue, chiefly seeds, which serves mechanically to stimulate the intestinal movements, local reflexes being set up in the nerve plexuses of the intestinal wall Undigested material of any sort within the bowel—e.g., the skins of fresh fruit, the cellulose of raw vegetables, whole wheat, and bran—all serve as intestinal stimulants by increasing the bulk and consequently the degree of stretch of the bowel wall Paraffin oil, which is not absorbed from the intestine except perhaps in minute amounts, acts in a similar manner. There are a host of other laxatives and cathartics, many of them used from time immemorial, of which little is known definitely concerning their mode of action. Vitamin B, and other factors of the B complex tend to increase the tone of the intestinal musculature, and thus favor natural movements of evacuation.

Reliance upon medicines to ensure regular bowel movements should be avoided, for the intestinal musculature tends to become tolerant to their use and requires a gradual increase in dosage. Hygienic measures, diet, exercise, and habit should be depended upon whenever possible.

The sensibility of the alimentary canal. The mucosa of the gastrointestinal tract for the greater part of its length is insensitive to the

<sup>&</sup>lt;sup>7</sup> The term laxative implies an agent with a milder stimulating action than either a cathartic or a purgative.

several forms of stimulation that arouse sensations from the skin, such as touch, cold and warmth, and pressure. This is not surprising, for this part of the body is never exposed to these types of stimulus, and this part of the body is never exposure to these specific has therefore not developed end organs that can respond to them. Pain is aroused by stretching the intestinal wall, but not by cutting, burning, or crushing. Thus distention of the stomach or intestine may cause intense pain as a result of stretching of nerve fibers in their walls. In disease, pain is often felt, not in the diseased organ itself, but in some part more or less remote Thus in angina pectoris (Chap. 4) pain is felt in the left shoulder and arm, and even in the fingers or jaw of the same side, and occasionally in the right arm. This is called referred pain and has been explained upon the fact that the organ and the part in which the pain is felt have developed from closely associated embryonic structures, and, in the adult, nerves supplying them carry impulses to the same segment of the spinal cord from where they are relayed to consciousness The pain is therefore projected to the part that receives its sensory nerve supply from this segment of the spinal cord. In other words, it is interpreted as being located in the part from which impulses of pain would normally arrive

The only parts of the alimentary canal from which sensations of touch, warmth, and cold can be aroused are the mouth, pharynx, and anal canal, though extremes of heat and cold can be appreciated in the esophagus and, according to some, even in the stomach. Common experience may seem to contradict the statement that warmth is not felt in the stomach, but the sensation aroused by warm food or drink is attributed to the conduction of heat through the abdominal wall to stimulate end organs in the overlying skin.

Hunger and appetite. Though we have often experienced both appetite and hunger, it is possible to describe these sensations only in the vaguest terms. Appetite, though closely associated with hunger, differs from the latter in that it depends upon pleasurable memories of the taste and smell of food Extraneous factors such as the sight of food and pleasant appurtenances of the meal also influence appetite (p. 311). Conditioned responses enter largely into its make-up. It is psychological rather than, like hunger, physiological. A newborn babe no doubt feels hunger but probably does not know appetite. Yet appetite is whetted by hunger and has, therefore, a physiological element; it is influenced by the tone and general state of the stomach. This is evidenced by the stimulating effect of an alcoholic drink taken before a meal and the increased appetite which follows the first few mouthfuls of a meal.



Fig 8 37 A tracing of the contractions of the stomach in hunger

Obviously, the sensation of hunger serves a protective function, for it controls the food intake. Its origin has a local (gastrointestinal) as well as a general (metabolic or blood composition) basis. Emptiness or fullness of the stomach and the state of its wall are potent factors influencing hunger, as is also the presence or absence of food products in the duodenum. We have all experienced that "gone" or empty feeling in the pit of the stomach when a meal has been long postponed. It is hard to say whether this arises in the stomach or the duodenum Spasms of pain—the pangs of hunger—may occur. These are caused by contractions of the stomach and have been graphically recorded in man by Carlson by means of a balloon passed into the stomach and connected by tubing with a recording apparatus (Figs. 8.37 and 8.38).

But gastric factors are incidental rather than primary, the fundamental cause of hunger is general or metabolic rather than local A




Fig 8 38 Showing the method of recording hunger contractions

low level of blood sugar (glucose) has been thought to be an important factor in its causation. This theory, however, does not accord with the excessive hunger characteristic of diabetes in which the blood sugar is above normal. It is more probable that the general or meta-bolic factor is a slow rate of glucose utilization by the tissues rather than the blood-sugar level itself. Thus during fasting, as the blood sugar falls and the rate of glucose utilization by the tissues is slowed, and in diabetes, though the blood sugar is above normal, glucose utilization is also reduced. This, which is known as the glucostatic theory of hunger, reconciles two otherwise irreconcilable factors. The actual control of food intake devolves upon a center in the brain (certain hypothalamic nuclei) whose cells, like those of other tissues of the body, will be subjected to changes in the rate at which glucose is used. It is postulated that a fall in glucose utilization by the cells of this "feeding center" arouses the urge to increase the food intake The responsive hypothalamic cells have been named glucoreceptors. There is experimental evidence to support this conception of the hunger mechanism. A lesion suitably placed in the hypothalamus of animals is followed by an increase in the food intake The opposite effect reduction in the food intake—is caused by an injury placed in another but near-by part of the hypothalamus Furthermore, it has been recognized for many years that in man disease involving this or neighboring regions of the brain may be associated with excessive hunger or with a pathological absence of hunger.

Thirst. Like hunger, thirst serves a protective function. It is nature's signal to replenish the body's water supplies. This sensation is referred to the pharynx, and is caused by the stimulation of sensory nerves in this situation. There has been much discussion as to whether the underlying cause of thirst is local—drying of the mucous membranes of the mouth and pharynx—or of a more general nature. Drying of the oral and pharyngeal mucous membranes is certainly conducive to thirst. But, if one is really thirsty, washing out the mouth or even gargling has little effect in allaying the sensation. Blood composition, probably the osmotic pressure of the plasma, seems to be the essential factor concerned. It seems likely that cells (osmoreceptors) in the hypothalamus are sensitive to changes in the osmotic pressure of the blood. We have seen that such receptors are part of the mechanism controlling the water output (see Chap. 7). It is a reasonable supposition that they also govern the water intake.

# Metabolism and Nutrition

Metabolism is a general term applied to the various chemical processes, whatever their nature, taking place in living tissues—e.g., the oxidation of food materials with the liberation of energy; the decomposition of compounds into more elementary principles; the chemical transformation of one material into another; and the synthesis of complex compounds from others of simpler constitution, as in the processes of tissue repair and growth, or in the manufacture of internal secretions and enzymes. Reactions involving decompositions are embraced by the term catabolism; those of a synthetic nature are referred to as anabolism.

Though some of the reactions taking place in the body are accompanied by the absorption of heat (endothermic reactions), in the great majority (e.g., oxidations) heat is evolved (exothermic reactions). The sum total of all the chemical reactions occurring in the body is referred to as general metabolism, and is expressed in terms of heat given out by the body in a given time. Those chemical changes, whether of a catabolic or an anabolic nature, which a particular substance (e.g., carbohydrate, fat, protein, purine, calcium, etc.) undergoes in the body are referred to as special metabolism.

## General Metabolism

The sun is the source of all energy on the earth. Plant life through its possession of the green coloring matter, chlorophyll, is capable of utilizing the energy of sunlight to form carbohydrate material from the carbon dioxide of the atmosphere and water drawn from the soil. This process is called photosynthesis. The animal body, of course, cannot make direct use of solar energy; it must depend upon the energy stored by the plant or upon that provided by the tissues of other animals. The food—fat, carbohydrate, and protein—derived from either of these

sources therefore represents stored or potential energy. The carbon and hydrogen of the food, after absorption from the intestinal tract, are oxidized in the tissues (Chap. 6). Energy is liberated thereby for muscular activity and for maintaining the vital functions—e.g., the action of the heart, the movements of the gastromtestinal tract, the excitability of nervous tissues, etc. Thus, the various foodstuffs are to the body as fuel is to an engine; they represent a certain amount of potential energy which the body can convert to other forms of energy—mechanical, electrical, chemical, and thermal.

When the body is at rest—that is, when no external work is being performed—the energy liberated from food materials appears ultimately as heat. A heat unit, the large Calorie, is therefore employed as a measure of the energy liberated in the animal body. A Calorie is defined as the quantity of heat required to raise a kilogram of water from 15° to 16° C. Of the energy expended during muscular exercise, as in lifting a weight, walking, etc., about 30 percent appears as work, the remaining 70 percent is converted to heat.

Antoine Lavoisier (1743–1794), the celebrated French scientist of the eighteenth century, pioneered in the field of metabolism. He established the crucial fact that the heat produced by a burning candle and that generated by the body were fundamentally similar processes. He showed that the body burned (oxidized) the food, the carbon being converted to carbon dioxide (CO<sub>2</sub>) and the hydrogen to water (H<sub>2</sub>O). He devised an ice calorimeter calculating the heat produced by the candle flame and that given out by the body of a small animal (e.g., a mouse) from the weight of ice which had melted during the period of the experiment. He came to but one false conclusion—namely, that the oxidation of the food with the production of carbon dioxide and water occurred in the lungs. We know now, of course, that it occurs in the cells of all tissues, a process which is known as internal respiration.

The law of the conservation of energy states that, though one form of energy is convertible into any other form, the sum total of the energy in the universe remains constant—energy cannot be created or destroyed. This law holds true for the animal body. That is to say, a given quantity of food when completely oxidized in the tissues yields its entire store of potential energy to the body—the energy intake balances the energy output, as measured by the heat produced and the work performed.

 $<sup>^1</sup>$  The large Calorie used in physiology is written with a capital C to distinguish it from the small calorie used in the physical laboratory, which is the quantity of heat required to raise, I gram of water from 15° C to 16° C.

#### METABOLISM AND NUTRITION

352

The food may not be completely oxidized in the body; a part may be stored, or, as with protein (pp. 353 and 363), a part resists oxidation and is excreted. Nevertheless, an energy balance can be struck, if any gain in weight is noted (stored energy) and the quantity of unoxidized residue of protein food in the urine is determined.

Energy is stored in the adult body as carbohydrate (glycogen) and fat, mainly the latter, whenever the food intake exceeds the energy




Fig 9.1 The bomb calorimeter A, a plannum dish upon which is placed the sample of food, B, bomb filled with oxygen, C, vessel holding water in which bomb is immersed, D, double-walled in-sulating jacket, E, fuse which can be ignited by an electric spark; F, water stirrer turned by a motor, M, G, thermometer, H, electric wires for ignition of fuse (Courtesy of Emerson Apparatus Company, Boston, Mass.)

needs of the body—that is, when the energy of the food is not all expended in maintaining the vital processes, or in performing muscular work. This is by far the commonest cause of overweight (obesity). This store of energy, in the form of fat and carbohydrate, is readily available and will be drawn upon should the energy value of the diet at any time fall below the body's energy requirements. During a prolonged period of fasting, after the stores of glycogen and fat have been drawn upon and exhausted, the protein of the tissues is utilized to furnish energy.

In the growing body, a part of the food not used to furnish energy is converted to new tissue with a resulting increase in weight.

The bomb calorimeter. The potential energy of a foodstuff is determined by measuring the heat evolved when it is burned in an instrument known as a bomb calorimeter (see Fig 9.1). The heat generated by the animal body may be determined in a manner identical in principle. Now, the quantity of heat liberated by 1 gram of carbohydrate or fat is the same whether the material is metabolized in the body or burned in the bomb calorimeter; 4.1 Calories are generated in the oxidation of 1 gram of carbohydrate, 9.3 Calories in the oxidation of 1 gram of fat. Protein, on the other hand, does not undergo complete combustion in the body. The nitrogenous part of the molecule resists oxidation and is excreted in the urine, mainly as urea (p. 363). A gram of protein, therefore, evolves less heat when catabolized than when burned outside the body. In the latter instance, 5.3 Calories are generated, in the former only 4.1.

Animal calorimetry. The heat generated by the body may be measured directly—direct calorimetry. The animal is placed in a closed chamber with double insulated walls; coils of copper tubing, through which water is circulated, are situated on the inner walls of the chamber; thus the heat given out by the animal is absorbed. The temperature of the water as it enters and leaves the chamber is recorded by thermometers.

The rise in temperature of the water in degrees centigrade, during the period of observation, multiplied by the total volume of water in kilograms which has passed through the chamber in that time, gives in Calories the heat given off by the animal. A few calorimeters of this type have been constructed for metabolic experiments on man, but their great expense precludes their general use. Indirect calorimetric methods are therefore employed almost entirely in metabolism studies upon the human subject.

Indirect calorimetry. We have seen that when carbohydrate undergoes combustion in the body, the carbon is oxidized to carbon dioxide (CO<sub>2</sub>) and the hydrogen to water (H<sub>2</sub>O). Thus:

$$C_6H_{12}O_6 + 6O_2 = 6CO_2 + 6H_2O$$
  
glucose carbon water

The body, therefore, reverses the reaction carried out by the green plant which, as already stated, synthesizes carbohydrate from carbon dioxide and water. In the complete oxidation of a given weight of carbohydrate, whether in air or in the tissues of the body, the quantity of oxygen used and of carbon dioxide produced have definite and con-

stant values. It is also known that for every 1000 cc of oxygen consumed when carbohydrate is the material oxidized, 5.047 Calories heat are evolved. That is to say, there is a constant relationship between the volume of oxygen consumed and the quantity of heat evolved. It is evident, therefore, that the heat production of an animal subsisting upon carbohydrate food could be determined, provided that the quantity of oxygen consumed by the animal were known. The relationship between the oxygen consumetion and the heat production varies, however, with the type of food undergoing combustion. Carbohydrate material as compared with fat is relatively rich in oxygen. Thus, C,H,(C,1H,Q2), is the formula for triolein, a common fat. More oxygen must therefore be supplied from a source outside the food itself for the complete oxidation of the carbon and hydrogen in fat than for the oxidation of these elements in an equal quantity of carbohydrate. The consumption of 1000 cc of oxygen, when fat is oxidized, is accompanied, therefore, by a smaller heat production—namely, 4.686 Calories as compared with the figure (5.047) given above for carbohydrate. The corresponding value for protein is 4.485 Calories. These values are termed the heat equivalents of the foodstuffs.

In order, therefore, to calculate the heat production of the body from the oxygen consumption, one must know the nature of the food mixture (the proportions of carbohydrate, fat, and protein) which is being metabolized. This information is obtained from the respiratory quotient, which will now be explained.

The respiratory quotient. The volume of carbon dioxide eliminated by the body during a given time, over the volume of oxygen absorbed, is called the respiratory quotient (R.O.). Thus:

From the equation given on page 353 it will be seen that, when the food undergoing combustion is carbohydrate, the volume of carbon divide climinated and of oxygen absorbed is equal. When, for example, 100 grams of glucose are oxidized, 75 liters of oxygen are absorbed and 75 liters of carbon dioxide produced. The R.Q. is, therefore,  $\frac{1}{12}$  — 1. In the complete combustion of fat, which is relatively poor in oxygen, the volume of oxygen used (200 liters per 100 grams) is greater than the volume of carbon dioxide produced (142 liters per 100 grams). The R.Q. is, therefore, ( $\frac{12}{12}$  — 0.71. The respiratory quotient for protein is 0.80 and for alcohol, 0.67.

On an ordinary mixed diet, the R.Q. of the human subject is around

Table 9-1

(After Zuntz and Schumberg, modified by Lusk)


Nonprotein	Calories per	Calories deri	Calories derived from	
respiratory quotient	liter O,	Carbohydrate	Fat	
		percent	percent	
0.707	4.686	0	100	
0.71	4.690	1.10	98.9	
0.72	4.702	4.76	95 2	
0.73	4.714	8 40	91.6	
0.74	4.727	12.0	0.88	
0.75	4 739	15.6	84.4	
0.76	4.751	19.2	80.8	
0.77	4.764	22.8	77.2	
0.78	4.776	26.3	73.7	
0.79	4 788	29.9	70.1	
0.80	4.801	33.4	66.6	
0.81	4.813	36.9	63.1	
0,82	4.825	40.3	59.7	
0.83	4.838	43.8	56.2	
0.84	4.850	47.2	52.8	
0.85	4.862	50.7	49.3	
0.86	4.875	54.1	- 45.9	
0 87	4.887	57.5	42.5	
0.88	4.899	60.8	39.2	
0.89	4.911	64.2	35.8	
0 90	4.924	67.5	32.5	
0 91	4.936	708	29.2	
0.92	4.948	74.1	25.9	
0.93	4.961	77.4	22.6	
0.94	4.973	80.7	19.3	
0.95	4 985	84.0	16.0	
0.96	4.998	87.2	12.8	
0.97	5.010	90.4	9.58	
0.98	5 022	93 6	6.37	
0 99	5.035	96.8	3.18	
1.00	5.047	100.0	0	

0.85. After fasting for 12 hours, it has a value of 0 82. When, as in the fattening of farm animals, fat is being formed from carbohydrate -that is, an oxygen-rich material is being transformed into one poor in oxygen-the R.Q. rises above 1. The heat equivalents of a liter of oxygen at different respiratory quotients are given in Table 9-1.

The respiratory quotients given in this table are for mixtures of fat and carbohydrate only (nonprotein respiratory quotients). For very precise work, the Calories derived from the metabolism of protein are calculated separately from the quantity of nitrogen excreted in the urine. But, as a matter of fact, only a negligible error is introduced when the figures in the table are used for calculating the heat production-that is, without any attention being paid to the protein metabolism.

The basal metabolic rate (B.M.R.). The heat production of the human body at its lowest possible level in the waking state is called the basal metabolism or the basal metabolic rate (B M.R.). In its determination, all conditions which have a stimulating effect upon heat production must be eliminated. It is measured, therefore, 12 hours after the last meal (usually in the morning without breakfast) with the subject lying down at complete physical and mental rest, and at a room temperature of about 20° C. No strenuous muscular exertion should be undertaken for 24 hours prior to the determination. The B.M R. is usually expressed in Calories per square meter of body surface per hour all may be determined from either the quantity of carbon dioxide eliminated, or from the quantity of oxygen consumed over a known period of time; but the figure for the latter is now most commonly employed (see Fig. 9.2) The heat value of a liter of oxygen for various

<sup>2</sup> In the early investigations of metabolic problems, it was the practice to express the heat production of the body in relation to its weight-that is, in Calories per kilogram of body weight per hour. But it was soon found that the surface area of the body rather than its weight was the determining factor in heat production. Heat is lost from the surface of the body (p. 377) and in order to maintain a constant body temperature, must be balanced by heat production. The greater the surface area of the body the greater is the amount of heat which is dissipated. Now small animals. such as mice, rats, etc., which have a much greater surface area in proportion to their weight than have larger animals, such as dogs, cattle horses, etc. (just as a marble has in relation to its mass a much greater surface than a baseball), have a correspondingly greater heat production on a body weight basis than have the larger animals. When, however, the heat generated is calculated and expressed in relation animas. Ween, nowever, the near generated is calculated and expressed in freatment to the surface area of the body, it is found to be approximately the same for mam-mals of all sizes under comparable conditions. Thus the heat production of a mour is around 27 Calories per kilogram of body weight per hour, while that of a man is only about 1.1 Calone per kilogram per hour. But both have a heat production of around 40 Calories per square meter of body surface per hour (Of course, the srea of the moure's body amounts to only a very small fraction of a square meter and the absolute quantity of heat tigwen out is relatively minute.



Showing method of determining the basal metabolic rate by means of the Benedict-Roth apis calculated from the oxygen consumption of Warren E Collins Inc., Boston) Oxygen is inhaled from the spirometer through c quantity paratus (

alone The right hand drawing is a diagram of the interior of the Benedict Roth apparatus,

metabolic mixtures (carbohydrate and fat), as indicated by the respiratory quotients, is given in Table 9-1. As a rule, the R.Q. is not actually determined, but is taken to be 0.82, for this is its value after fasting for 12 hours, under which conditions, as just stated, the B.M.R is determined.




Fig. 9.3 Du Bois' chart for obtaining the surface area of the body in square meters (as indicated by the slanting lines) from a subject's height and weight. For example, a person 170 centimeters (approximately 5 feet 8 inches tall and weighing 70 kilograms (154 pounds) has a surface area of 18 square meters.

Sample calculation of the B.M.R. from the oxygen consumption. If the oxygen consumption over a period of 10 minutes is 2.500 liters, then the hourly heat production (under the condutions defined and at an R.O. of 0 82) is  $2.500 \times 4.825 \times \frac{a_0}{10} - 72.36$  Calories. If the subject has a surface area of say, 1.8 square meters, which is the figure for the average male adult, then the hourly heat production per square meter of body surface is:

$$\frac{72.36}{1.8}$$
 = 40.2 Calories

The surface area of the body bears a relationship to the height and weight and may be obtained for a given person from the chart shown in Figure 9.3.

In young healthy adult males, the basal metabolic rate is around 40 Calories per square meter of body surface per hour. It is lower in women and higher in children, being around 52 Calories during the early years, and diminishing gradually throughout life. It reaches the adult value at about 20 years of age; from then it declines more slowly,

reaching a value of around 36 Calories in old age. The heat production per square meter of body surface per hour for males and females of different ages is given in Table 9.2

Table 9-2

Oxygen consumption (liters) and heat production (Calories) per square

meter of body surface per hour

Ages	Male		Female	
	O <sub>z</sub>	Calories	0	Calones
14-15	9 53	45 9	8 91	42 9
16-17	8 91	429	8 29	39 9
18-19	8 50	409	7 88	37 9
20-29	8 19	39 4	7 67	369
30-39	8 19	39 4	7 57	36 4
40-49	7 98	38 4	7 46	35 9
50-59	7 77	37 4	7 25	34 9
60~69	7 57	364	7 05	33 9
70-79	7 36	35 4	6 84	32 9

Note the gradual diminution in heat production with advancing years

The physiological conditions which raise the metabolism are muscular exercise, a low environmental temperature, and the ingestion of food, especially protein. The term specific dynamic action is applied to this stimulating effect of food upon metabolism. During sleep the metabolism is lowered by about 15 percent below the basal level.

Among pathological conditions which increase the metabolism are hyperthyroidism (p 411) and fever (p 381), it is lowered in hypothyroidism (p 408), undernutrition, deficiency of the adrenal cortex (p 426), and in certain pituitary disorders

# The Metabolism of Protein

Protein material enters largely into the composition of all types of protoplasm, both of animal and vegetable origin. Among foods, meat (muscle), cheese, eggs, beans, and peas (legumes) are the richest sources. Its basic elements are carbon, hydrogen oxygen, nitrogen, and sulfur, and usually phosphorus. Its content of the last three elements distinguishes it from either fat or carbohydrate, which contain only carbon, hydrogen, and oxygen. The body is dependent almost entirely upon protein for its supplies of nitrogen and sulfur, and mainly for its phosphorus.

#### METAROLISM AND MILEPITION

The amino acids. The protein molecule is made up of a number of amino acids linked together Some twenty three different amino acids have been discovered. The simplest of these is called glycine or glycocol, which consists of acetic acid in which a hydrogen atom is re placed by an NH2 group Thus

> CH. CH. - NH. glycine

The majority of the other amino acids are constructed upon the same general plan They contain a fatty acid—e g, propionic, valetianic, caproic, succinic etc—and one or two NH<sub>2</sub> (amino) groups An in complete list of the amino acids with their empirical formulas is given in Table 9.3

#### Table 9 3

## A partial list of the amino acids (see also appendix)

Glycine (or glycocoli) C.H.NO. or amino acetic acid

- Alanine C<sub>2</sub>H<sub>2</sub>NO<sub>2</sub> or α amino propionic acid \*Threonine C<sub>1</sub>H<sub>2</sub>NO<sub>2</sub> or α amino-β hydroxy n butyric acid
- \*Valine, C.H., NO, or a amino isovaleric acid
- \*Leucine, C.H., NO, or a amino isocaproic acid
- \*Isoleucine C.H., NO, or a amino B ethyl B methyl propionic acid Aspartic acid C.H.NO. or amino succinic acid
- Glutamic acid C.H.NO, or a amino glutaric acid
- \*Arginine, C<sub>t</sub>H<sub>14</sub>N<sub>4</sub>O<sub>2</sub> or δ guanidin α amino valerianic acid
- \*Lysine C.H.N O1 or \alpha e-diamino caproic acid Cystine C<sub>6</sub>H <sub>1</sub>N<sub>2</sub>S-O<sub>4</sub> or di-cysteine or di-(β thio-α amino propionic acid)
- \*Methionine C.H., SNO, or a amino y methyl thiol n butyric acid
- \*Phenylalanine, C2H11NO2 or β phenyl α amino propionic acid
- Tyrosine, C<sub>2</sub>H<sub>11</sub>NO<sub>2</sub> or β parahydroxy phenyl α amino propionic acid
- \*Tryptophane, C11H 1N2O2 or β indole α amino propionic acid
- \*Histidine, C.H.N.O., or α amino-β imidazol propionic acid

The protein molecule, being built of amino acids linked together, possesses free COOH (acid) and NH<sub>2</sub> (basic) groups It is capable, therefore, of acting as a weak acid or a weak base depending upon the reaction of the medium in which it is dissolved. In alkaline solution, it behaves as a weak acid, in acid solution as a weak base. In the blood plasma and the other body fluids (which are slightly alkaline in reaction), protein, acting as an acid, forms with bases compounds called

proteinates. The chief base is sodium. The proteinate undergoes ionization (p. 12), into proteinate anions (P-) and cations (Na<sup>+</sup>, K<sup>+</sup>, etc.). In acid solution, protein hydrochloride, protein sulfate, etc., are formed, the protein then acting as a weak base. At a certain reaction

## Table 9-4

### Classification of proteins

Simple proteins

Albumins—e.g., ovalbumin of egg white, serum albumin, lactalbumin, and certain vegetable albumins

Globulins-e.g., serum globulin, fibrinogen, vitellin of egg yolk, and legumin of peas

Glutelins-e.g., glutenin of wheat

Gliadins—e.g., gliadin of wheat, hordein of barley, and zein of maize Scleroproteins—e.g., keratin of hair, elastin, collagen (which yields gelatin) of connective tissues

Histones-e.g., the globin in hemoglobin

Protamines, in spermatozoa—e g., salmine in spermatozoa of salmon Conjugated proteins—i e., proteins containing a nonprotein group

Nucleoproteins, protein combined with nucleic acid (p. 366)

Chromoproteins, protein containing a pigment group, e.g., hemoglobin

Glycoproteins, protein containing a sugar group—e g., mucin Lipoproteins, proteins whose molecule contains a fatty acid group; pres-

ent in protoplasm of cells of various tissues including blood

Phosphoproteins, proteins containing a phosphorus group (other than
nucleoproteins)—e.e., casein of milk

Derived proteins, products of the action of acids, heat, or enzymes upon proteins

Metaproteins

Coagulated proteins

Proteoses

Peptones

Peptids, dipeptids, tripeptids, and polypeptids

nearer the neutral point, ionization is reduced to zero; the protein particles, being electrically neutral, collect together into clumps. A precipitate of the protein material forms which is visible to the naked eye; this is called isoelectric precipitation, and the point or range of pH at which it occurs is known as the isoelectric point or range. The latter varies considerably for different proteins but is characteristic for a given protein. Proteins (or the constituent amino acids), on account of their ability to form acids or bases depending upon the H-ion concentration, are termed amphoteric electrolytes (Gk. ampho. both). There are several proteins which contain more than 15 varieties of amino acids, but the assortment varies between different types of protein. The total number of amino acids in a protein molecule may be 200 or more. Such proteins are composed, therefore, of molecules of relatively enormous size. The molecules of certain other proteins, such as the protamines, contain much fewer amino acids and are correspondingly smaller. There is thus a very wide variation in the size of the molecules of the different types of protein. Egg albumin, for example, as molecular weight of around 40,000, whereas the molecular weights of some other proteins are in the neighborhood of 200,000 or more. A few have a molecular weight of over a million. The shape of the protein molecule also varies. The molecules of some proteins such as the myosin of muscle and of that forming tendons and ligaments (collagen) are long and fiberlike in shape, while others, such as those of serum albumin, are globular. A short classification of proteins is given in Table 9.4.

# The History of Protein in the Body

The growth of body tissue in the young animal, and the repair of protein structure in the adult body are dependent upon the protein (amino acid) content of the diet. But since the tissue protein of a given animal is different in constitution from the protein of its food, the latter as a rule must be completely broken down into its separate amino acids before it can be utilized for building body tissue.

The utilization of protein in the construction of body tissue may be compared to the building of a number of houses of different types from materials obtained from the wrecking of other structures. Each brick and stone in the old buildings is separated, and then sorted and carted to the new sites. Some of this building material will be more suitable for one type of house, some more suitable for other types. Other materials again cannot be used at all, and are therefore discarded as refuse. The new buildings, though constructed from materials taken from the old, will therefore be quite different in structure and general plan.

From this analogy, the amino acids are sometimes called the "building stones" of the protein molecule. Under the action of the peptidases of the intestinal juice (p. 327), the protein molecule is completely demolished, in the sense that it is broken up into its constituent amino acids. The separate amino acids are absorbed into the blood stream and carried therein to the tissues. Each tissue chooses those which it can make use of, rejecting others.

Those amino acids which cannot be utilized by the body are broken up; the carbon part of the molecule (i.e., the fatty acid group) is oxidized to carbon dioxide and water, thus furnishing energy to the body or is converted to body fat; the nitrogen goes to form urea which is excreted in the urine. The removal of the nitrogen (amino) group from the amino acid molecule is called deamination; this process occurs chiefly in the liver-Certain amino acids not used for the construction of body protein may, after undergoing deamination, be converted to glucose which is then either oxidized or stored in the liver as glycogen (p. 368). In diabetes the glucose formed from amino acids is largely excreted in the urine. The diabetic subject, therefore, continues to excrete sugar though he receives no sugar or other carbohydrate; even during fasting he continues to excrete glucose, which is then derived from the protein of his own tissues.

The urea in the urine is formed mainly through the deamination of amino acids composing the protein of the food. Raising the protein content of the diet, therefore, increases the output of urea in the urine and vice versa. But the urea is not entirely derived from the diet; a small part is formed from the nitrogen released by the breakdown of body protein, for the tissues are constantly undergoing disintegration and repair. The urea of the urine is thus in part exogenous (from food) and partly endogenous (from body protein). Creatinine, another nitrogenous constituent of the urine, is also derived from tissue protein, but unlike urea it is derived almost entirely from this source. Varying the protein of the diet, therefore, exerts little effect upon the excretion of creatinine. A small quantity of nitrogen, combined in various ways, is also eliminated from the body by passing through the wall of the intestine into the feces.

During starvation or upon a protein-poor diet, the breakdown of body protein, of course, continues. The body loses nitrogen through the so-called "wear and tear" of tissue protein, but receives none to make good the loss. That is to say, the output of nitrogen in the urine and feces exceeds that of the food; the body is then said to be in negative nitrogen balance. Similarly, the nitrogen output will exceed the intake, and the body will be in negative nitrogen balance, if the food protein is inadequate in amount or of poor quality (ie., if its assortment of amino acids is unsuitable for repairing body protein, the nonutilizable amino acids being excreted in the urine). In the healthy adult receiving an adequate diet, the nitrogen excreted just balances the nitrogen taken in the food—that is, the nitrogen lost as a result of the breakdown of tissue protein is replaced from the food; the remainder

of the food nitrogen is excreted. The body is then said to be in nitrogen equilibrium.

During growth, after a period of starvation, in pregnancy, in muscular training, or in convalescence from some disease which has caused the excessive destruction of body protein, the quantity of nitrogen excerted is less than that taken in the food, provided the protein intake is adequate. The body is then in positive nitrogen balance. In other words, nitrogen is retained for the construction of body tissue.

The 'wear and tear quota of protein metabolism is reduced by car bohydrate food For example the nitrogen excretion is considerably less on a diet containing carbohydrate but no protein, than in starvation The carbohydrate diminishes the breakdown of tissue protein This is spoken of as the sparing effect of carbohydrate upon protein metabolism.

We have seen that only part of the protein molecule can be oxidized in the body and thus provide energy. Civilized man cannot subsist on protein food alone because he cannot eat and digest enough to supply the required energy. One gram of protein furnishes 4.1 Calories. A man of average size has a daily energy expenditure under basal conditions of about 1400 Calories, and between 2500 and 3000 Calories when engaged in light occupation. Now, lean meat is about 20 percent pure protein. Therefore about

$$\frac{1400}{41} \times \frac{100}{20} = 1700$$
 grams

or nearly 4 pounds of meat would need to be eaten daily, in order to obtain from protein the energy required by the body, even at rest From three to four times this amount of meat would be required to furnish the energy for heavy work. On the other hand, a carmivorous animal such as the dog which has an energy expenditure of some 600 Calories, can consume 2 or 3 pounds of meat in a few minutes, and can therefore derive its total calorie requirement from such food. The Eski mos are also accustomed to the consumption of relatively enormous quantities of meat and fat

The essential ammo acids and the relative nutritional values of different proteins. An essential or indispensable ammo acid may be dofined as one which must be present in the diet in order that the growth of young animals may proceed normally, and that the health of both young and old animals be maintained. In a sense, probably all the ammo acids are essential in one way or another for nutrition, but some (e.g., g(youe)) can be synthesized in the body or formed from others fur nished in the diet. It is not necessary, therefore, that such amino acids shall themselves be present in the food. There are ten essential amino acids; they are marked with an asterisk in Table 9-3 on page 360.

Proteins are not all of equal value in nutrition, for the reason that the various types differ widely in their amino-acid constitution. Those which possess an amino-acid assortment most closely resembling that of the body's proteins have the highest nutritive value. Generally speak-

Table 9-5
Character of proteins in some common foods

Food	Chief proteins present	Amino-acid constitution
Milk and cheese	Casein Lactalbumin	Complete, but low in cystine Complete
Corn (maize)	Zein	Lacks lysine and tryptophane and is low in cystine
Eggs	Ovalbumin Ovovitelline	Complete Complete
Meat	Albumin Myosin	Complete Complete
Peas	Legumin	Incomplete, low in cystine
Wheat	Gliadin Glutenin	Incomplete lacks lysine Complete
Gelatin	Gelatin	Incomplete, lacking tryptophane and tyrosine; very low in cys- tine

In part from M. S Rose, Foundations of Nutrition.

ing, and as might be expected, these are of animal origin—for instance, the proteins of milk, eggs, and meat. Some proteins, such as gelatin and zein (in maize) are incomplete—that is, they lack certain amino acids which are essential for growth and for the maintenance of nitrogen equilibrium in the adult. Others, such as gliadin (in wheat), hordein (in barley), and legumin (in peas), will serve to repair body protein in the adult but will not support growth, since they lack certain essential amino acids or contain them in insufficient amounts. Young animals which receive one or other of these as their sole protein fail to grow. Wheat, peas, and barley, however, contain other proteins which make good the amino-acid deficiencies of those which are incomplete (see Table 9-5). Lactalbumin of milk, and ovalbumin of egg white, the proteins of meat, and glutenin (of wheat) contain all the essential amino acids.

## Purine Metabolism


366

Uric acid (C<sub>1</sub>H<sub>1</sub>N<sub>1</sub>O<sub>2</sub>), a nitrogenous compound present in normal urine and blood, is a purine derivative Purines are constitutents of nu cleic acid and this, in turn, is found in the body combined with protein the compound being called nucleoprotein Nucleoprotein is present in the nuclei of cells generally, and such tissues as those of thymus liver kidney pancreas and other glandular stuctures are particularly rich in this material Viruses are now thought to be composed of nucleopro-

Nucleic acid is constituted of four compounds called nucleotides. A nucleotide contains a molecule each of phosphoric acid the pentose ribose (a sugar with 5 carbon atoms and 5 H,O molecules), and a purine (adenine or guanine) or a pyrimidine group. The nucleoprotein of food is broken down by intestinal enzymes into protein and nucleic acid. The latter is split into its constituent nucleotides by an enzyme in the intestinal juice called nuclease. The nucleotides are ab sorbed and through the actions of specific tissue enzymes are broken into their components. Prosphoric acid is first removed leaving a pentose purine (or pyrimidine) compound called a nucleoside. The sugar is then split off. The fate of the purine derivatives adenine and guanine only need concern us here. These as a result of deamina tion (splitting off of NH<sub>3</sub>) and oxidation by tissue enzymes give rise to urice acid.

Uric acid is of especial interest because in gout, a painful arthritic condition its excretion is reduced and its concentration in the blood increased Uric acid in the form of crystals of sodium urate is de posited in the tissues surrounding the affected joint Meats such as liver sweetbreads kidney etc. which are rich in nuclear material and con sequently in nucleic acid tend therefore to aggravate the symptoms of rout.

The origin of uric acid is briefly summarized in the following scheme



# Carbohydrate Metabolism

Classification of the carbohydrates. Carbohydrates (sugars, starches, etc.) are composed of carbon, hydrogen, and oxygen. The last two elements are present in the same proportions as in water (H<sub>2</sub>O). Thus, the three sugars, glucose (or dextrose), fructose (or levulose), and galactose, contain six molecules of water and six atoms of carbon, as shown by their formula, C<sub>4</sub>H<sub>11</sub>O<sub>4</sub>. These sugars are therefore called hexoses (Gk. hex, six). Though they have a common empirical formula—i e., the same number of each type of atom—the positions of the atoms in the molecule differ. The structural formulas of glucose, fructose, and galactose are given below for comparison.

Glucose is found in fruits and in the blood and tissues of animals Fructose is the chief sugar of honey. Galactose is present, as a component of the disaccharide lactose, in milk, being combined in the molecule with glucose. It is also found as a constituent of certain fatty compounds in brain tissue. Other much less common sugars have molecules containing seven, five, four, three, or two carbon atoms and a corresponding number of water molecules. Employing Greek prefixes, they are called heptoses, pentoses, tetroses, trioses, and dioses, respectively. Thus a heptose is represented by the formula C<sub>7</sub>H<sub>11</sub>O<sub>1</sub>, a pentose by C<sub>4</sub>H<sub>10</sub>O<sub>1</sub>, a tetrose by C<sub>4</sub>H<sub>10</sub>O<sub>1</sub>, and so on. Octoses, nonoses, and decoses, though not known in nature, have been prepared in the laboratory.

Sugars belonging to the class just outlined are called *monosaccha-rides*, and may be represented by the general formula  $C_n(H_2O)_n$ , in which n has the value 2, 3, 4, 5, 6, 7, and so forth.

Disaccharides are sugars composed of two monosaccharide molecules less a molecule of water. Sucrose (cane, beet, and maple sugars), maliose (sugar of malt), and lactose (sugar of milk) belong to this group Though a disaccharide may be made up of other monosaccharides, these three, which are important food elements and, therefore, of physiological interest, consist of hexoses either of the same or of different kinds Their formula is C<sub>11</sub>H<sub>22</sub>O<sub>11</sub>. They are split (hydrolyzed) into their constituent monosaccharides by the actions of specific enzymes in the intestine Thus—

#### $C_{10}H_{10}O_{11} + H_{10}O = 2C_{1}H_{10}O_{1}$

Sucrose is hydrolyzed into a molecule each of glucose and fructose, amaltose into two molecules of glucose, and lactose into glucose and galactose

galactose

Polysacchardes are made up of a large number of monosaccharde
molecules which may be either pentoses or hexoses, less water The
polysacchardes of physiological importance—namely, the vegetable
starches glycogen ('animal starch''), cellulose, and dextrins—are
constituted of glucose molecules, and are therefore given the general
formula of (C<sub>s</sub>H<sub>s</sub>O<sub>s</sub>). Upon hydrolysis they yield glucose

The history of carbohydrate in the body. Glycogen was discovered
by the great French physiologist, Claude Bernard, in 1857 It is found
in traces in most tissues of the body and in fairly large amounts in liver
and muscle Muscle may contain any amount of glycogen between
0 10 and 1 0 percent of fresh tissue Since approximately one half the
total weight of our bodies is muscle. It will be approximately one half the total weight of our bodies is muscle, it will be appreciated that the total quantity of muscle glycogen is very considerable. The liver often con tains as much as 10 or 15 percent of its wet weight of glycogen and,

though it makes up only about 3 percent of the body weight, its total glycogen content is comparable with that of the muscular tissues. Glucose is found in all body tissues, blood contains approximately 0 1 percent but there is considerable variation in the amount in health or percent out enter is considerable variation in the amount in reasonable as well as in disease. It is the most important single carbohydrate with which we have to deal in physiology. As such it can be used as food, passing unchanged from the small intestine into the blood while complex carbohydrates, such as the disaccharides and polysaccharides (starches and glycogen), must first be hydrolyzed into glucose or other hexoses before they can be absorbed in significant amounts


Glucose (as well as fructose and galactose) passes from the intestine into the blood of the portal vein, and is deposited in the liver as glycogen. The conversion of glucose to glycogen is a specific function of the liver cells, and is termed glycogeness (literally, glycogen formation)

The mixture is called invert sugar

The glucose in the blood is maintained at a fairly constant level by the reconversion of glycogen to glucose, which is then discharged into the general circulation. This process is termed glycogenolysis (literally, glycogen breakdown).

Muscle glycogen is derived from the glucose of the blood. During muscular contraction, energy is furnished mainly by carbohydrates; the muscle glycogen breaks down into its constituent glucose molecules. Lactic acid, which is ultimately produced, is in part resynthesized to glycogen and in part oxidized to carbon dioxide and water (see Chap. 11). Lactic acid, as well as the three hexoses-glucose, fructose, and galactose-is, therefore, a glycogen former. A part of the lactic acid produced diffuses into the blood and is deposited as glycogen in the liver, as well as in the heart muscle and certain other tissues. Glucose is also formed from the amino acids of food protein; in starvation or upon a protein-free diet, it is produced from body protein. This process is carried out in the liver, and is termed gluconeogenesis (literally, the new formation of sugar). There has been some question in the past whether sugar could be formed from fat, but it is now accepted by most authorities on carbohydrates that the conversion does occur.

The carbohydrate cycle just outlined may be represented in the following scheme.



Oxidation of carbohydrate. While it appears probable that glucose is the only sugar oxidized directly in the tissue cells, there may be some utilization of fructose in this manner (see p. 367).

Diabetes and the action of insulin. Insulin, the antidiabetic hormone, is produced in the pancreas; when this gland is completely removed from animals, a condition known as pancreatic diabetes rapidly develops. This is characterized by a high blood sugar (0.2-0.5 percent, as compared with the normal of 0.1 percent) and the excretion of sugar in the urine. The extent of the increase of blood glucose (hyper-glycema), and of the loss of glucose in the urine (glucosuria or glycosuria) is proportional to the carbohydrate content of the diet, but even on a diet composed entirely of protein, or during fasting, the depancreatized animal excretes a considerable amount of sugar. The glucose reserves of the body (e.g., liver glycogen) are rapidly lost Sugar is then made from tissue protein, the animal losing weight as a result. The fat depots are also called upon; the fat may be oxidized directly or first converted to glucose. At any rate, the increased rate of breakdown of fat results in the accumulation of ketone bodies in the blood (ketosis) and unne (ketonuria).

The ketone bodies are  $\beta$ -hydroxybutyric and acetoacetic acids and acetone. The two organic acids are normal intermediate products in the metabolism of fat,  $\beta$ -hydroxybutyric acid being derived, through oxidation, from acetoacetic. Both acids are present in the body in health though only in small amounts, acetoacetic acid undergoing almost complete oxidation to carbon dioxide and water. When, on the other hand, the metabolism of carbohydrate is defective, as in diabetes, for some reason not altogether clear, excessive quantities of fat are metabolized; acetoacetic acid is formed in such large amounts that it cannot be completely oxidized. It and  $\beta$ -hydroxybutyric acid accumulate in the blood and tissues, and are responsible for the acidosis and coma of diabetes. Acetoacetic acid is believed to be an especially toxic product. In diabetes, it is converted, through the loss of a molecule of carbon dioxide, to acetone, a volatile substance with a character-stic odor. Acetone is exhaled in the breath and excreted in the urine.

All the signs and symptoms seen in animals following removal of the pancreas are seen in human diabetes mellitus which, in most instances, is the result of disease of the islets of Langerhans in the pancreas. These structures are solely responsible for the body's supply of insulin They can be seen in a section of the pancreas as small groups of cells lying here and there between the pancreatic alveoli which secrete the pancreatic juice (Fig. 8.24, p. 323). In diabetes there is also an accumulation in the blood of neutral fat in the form of microscopic partucles called chylomicrons. This phenomenon is known as lipemia.

The disturbances of carbohydrate metabolism characteristic of the

diabetic state may be explained on the basis of (1) failure of utilization (oxidation) of carbohydrate or (2) abnormal rate of production of sugar from noncarbohydrate sources. The results of modern research favor the second alternative.

The discovery of insulin has not solved the problem of the cause of diabetes, but has provided an effective means of treating the disease. It has also furnished a most valuable "tool" for studying carbohydrate metabolism. Insulin administered subcutaneously or intravenously eliminates all the signs of diabetes in experimental animals or in human patients. Its action is to promote the storage of glycogen in muscles and liver, to depress the wasteful new production of sugar in the liver, and to increase the oxidation of carbohydrate. The hyperglycemia and glucosuria disappear. Ketosis is eliminated. The rapid loss of body tissues is checked.

Insulin has now been prepared in crystalline form; it is a protein containing eight or more amino acids. The highly purified insulin is absorbed quickly, when administered subcutaneously, and efforts have been made to lengthen the period of absorption—i.e., to prolong its action. The most satisfactory preparation thus far developed for this purpose is protamine zinc insulin, which is made by adding protamine and zinc to insulin Protamine is a simple protein obtained from fish sperm (see Table 9-4).

While a deficiency of insulin causes diabetes, overproduction is responsible for clinical condition called hyperinsulinism. The outstanding features of this disease are a low blood sugar (hypoglycenia), and the symptoms incident thereto. A similar condition may result from an overdose of insulin. The signs of hypoglycemia are neuromuscular hyperexcitability and hunger. If the blood sugar continues to fall, the excitability increases, leading to involuntary twitching of muscles, and later to generalized convulsions. Glucose is the best antidote for an excess of insulin. Other carbohydrates are effective in proportion to their ability to form glucose.

Other hormones and carbohydrate metabolism. The influence upon carbohydrate metabolism of certain hormones of the anterior lobe of the pituitary is dealt with on page 435, and of thyroxine and the hormones of the adrenal gland on pages 411 and 427, respectively.

These several hormones constitute a complex, but delicately balanced mechanism whereby the metabolism of carbohydrate is regulated. The various phases of, and the ceaseless changes in carbohydrate metabolism—the utilization of glucose by the tissues, the formation and storage of glycogen and its breakdown to glucose, and the new

372 METABO

formation of glucose from protein and fat—all come under the in fluence of one or more of the hormones mentioned. Due to their reciprocal actions the sugar of the blood and other body fluids is main tained at an approximately constant concentration.

The nervous system and carbohydrate metabolism In 1855 Claude Bernard showed that injury to the brain which involved the pons and the cerebellum as well as the floor of the fourth ventricle produced hyperglycemia and glucosuria. It appears probable that these lesions set up nerve impulses which caused the breakdown of liver glycogen. This may be due to the conduction of the impulses directly to the liver or may be an indirect result of adrenaline liberation. It will be appreciated that interference with the nerve supply of any of the glands whose secretions affect carbohydrates may exert a profound influence on the metabolism of these substances.

## Fat Metabolism

The neutral fats. The common fats of vegetable and animal tissues are compounds of the higher fatty acids palmitic  $(C_1H_1O_2)$  stearic  $(C_1H_2O_2)$  and oleic  $(C_1H_2O_1)$  in combination with the triatomic scombined with three molecules of one or other of these fatty acids. The resulting compound (or ester) is called a neutral fat or triglycer ide. Depending upon the particular fatty acids in the triglyceride mole cule the three chief fats are named tripalmitin tristearin and triolein respectively

The fatty tissues of animals consist of connective tissue in which is deposited a mixture of neutral fats triolein being in the greatest proportion. Tripalmitin is present in smaller and tristearin in least amount.

The triglycerides are hydrolyzed by the intestinal enzymes (lipases) into their constituents—fatty acids and glycerol. In the presence of alkali fat is decomposed the fatty acid then reacting to form soap. Thus

Fatty or waxy substances-sterols, steroids, and phospholipids or phosphatides. The sterols are secondary alcohols which are present in animal and vegetable tissues in combination with fatty acids. The sterols are but one group of a class of fatty or waxy substances known as steroids They are all closely allied chemically and include the hormones of the sex glands and of the adrenal cortex. Short descriptions of the more important steroids follow

Cholesterol is widely distributed throughout animal tissues both as such and as cholesterol esters. It was first isolated from gallstones and is an important constituent of bile, hence its name (Gk. chole, bile), and of blood. It is closely related chemically to cholic acid, vitamin D, and the sex hormones. Cholesterol and its esters are especially abundant in the sheaths of nerves, in brain tissue, and in the skin.

Ergosterol is found in plant tissues, it acquires antirachitic properties upon irradiation with ultraviolet light

The phospholipids or phosphatides are essential constituents of ani mal and vegetable cells Brain, muscle, liver bile, milk, and eggs con tain these substances in especially large amounts. To this class of substance lecithin cephalin and sphingomyelin belong. Upon hydrolysis they yield fatty acids, phosphoric acid and a nitrogenous base. The nitrogenous element in lecithin is choline.

Animal fat is formed from the fat of the food as well as from carbo hydrate. It represents a store of energy which the body can draw upon as need arises (e.g., during a prolonged period of fasting). The fat of the food is, of course, not deposited in the tissues unchanged. The fatty acids derived from the ingested fat are recombined with glycerin in such proportions as to produce a fat characteristic of that of the animal sown body—that is, with the proper mixture of the three types of neutral fat.

It will be recalled that the caloric value of fat (93) is more than double that of either carbohydrate or protein, which makes it the ideal energy storing material Furthermore, fatty tissue is almost pure fat whereas carbohydrate and protein materials are laid down in the tis sues with a relatively large quantity of water

The phospholipids are believed to play an important part in fat absorption and fat metabolism Fatty acids, it is thought, must first be transformed to phospholipids before they can be transported across cell membranes

The liver and fat metabolism This organ occupies such a central position in any consideration of fat metabolism that particular attention must be directed to it. The liver is the principal if not the only site of production of the ketone bodies (see p. 370). Under a great variety of pathological conditions, neutral fats accumulate in the liver. In diabetes, in anemias, and after various types of poisoning, large de posits of fats are found in this organ. This finding suggests that, under

these conditions, either the fat which normally is transported from the fat stores of the body (such as in the subcutaneous tissues or surround ing the abdominal organs) to be dealt with by the liver is, brought there in excessive amounts, or the hepatic cells are in some way functionally depressed and unable to deal with even the normal amount of fat

Oxidation of fats. The fact that the long chain fatty acids are oxidized very readily in the body makes the detection of intermediate products a difficult matter. The observations of Knoop in the living animal suggest that the so-called \( \theta \) oxidation takes place—that is, that there is successive removal of groups of two carbon atoms.—The fact that the ketone bodies—i.e., those with four carbon atoms—are formed only from fatty acids with an even number of carbon atoms obviously supports the theory of \( \theta \)-oxidation. The results of in vitro experiments, however, suggest that rupture of fatty acid chains takes place at various points in the chain. It is thus apparent that our knowledge of fat oxida tion is very incomplete.

The digestion and absorption of fat are dealt with in Chapter 8

Obesity, Overweight of the body due to the excessive accumulation of fat is called obesity or adiposity. Though a very common state, it must, unless of very moderate degree, be considered an abnormality obesity is a definite hazard to the general health and in several ways puts the physiological processes at a disadvantage. It places a greater than necessary burden upon the heart at all times but especially in the performance of muscular work. Diabetes arterial hyportension, degen crative changes in the arteries and gallistones are among the conditions which are more common in the obese than in persons of normal or subnormal weight. Diabetes has been called the 'fat man's folly' Obesity tends to reduce the body's resistance to infection, and the extremely obese person is looked upon as a poor risk by the surgeon, as well as by the insurance examiner. Someone has stated the case of the overweight person of middle age in respect to life expectancy in the epigram "the longer the belt the shorter the life." This is well borne out by insurance statistics. In persons over 45 years of age, the age at death of those who are overweight is considerably below the average, while that of lean persons is above average.

while that of lean persons is above average

The cause of obesity Some cases of obesity are due to an endocrine
disorder, or other specific disease But such causes are rare. The com
mon type of obesity, which is usually referred to as simple obesity, is
due to an intake of energy (food) in excess of the energy output
(work). That is to say, the subject eats too much for the physical work
he does, or he prefers foods of higher caloric value than does the per-

son of normal weight. The excess food is deposited as fat. This conclusion has been frequently questioned, but attempts to prove otherwise, by experiments upon human subjects, have failed repeatedly to prove that it is false. The metabolic rate, for example, is not lower in the obese than in lean persons. In other words, the metabolic processes of obese persons are not more than usually economical in the conversion of food energy into work.

# The Regulation of Body Temperature

The temperature of the human body is determined by placing a thermometer in the mouth, axilla (armpit), or rectum. In health, the mouth temperature is around 98.6° F. The temperature in the axilla is about a degree lower than this, and the rectal temperature about a degree higher. Even in health, the body temperature does not remain at a constant level throughout the 24 hours, but is from 1° to 1.5° lower in the early morning than in the late afternoon. The cause of this diurnal fluctuation in body temperature is unknown.

The body temperature represents the balance struck between the heat generated by the active tissues, mainly the muscles and liver, and

that lost from the body to the environment. It is remarkable how steady the body temperature remains under widely varying conditions Little change in body temperature occurs though the air temperature rises to 100° F or falls below zero, nor does the extra heat produced during light work occasion a rise in temperature. Strenuous muscular exercise may, however, cause a temporary rise in temperature of from 1° to 4° F.

Animals such as mammals and birds (warm-blooded animals) which can maintain a constant body temperature against variations in the temperature of their environment are called homoio-thermic. Those species—e.g., fish, frogs, and restricts (cold blooded pairwise).

and reptiles (cold-blooded animals)—that are unable to regulate their body temperature are called *poikilothermic*. Their body temperature is that of the environment (see Fig. 9.4).

Heat balance. It is obvious that the quantity of heat produced in the body (pp. 378-9) must just balance the quantity lost to the environ-




Fig 94 Variation in body temperature of different types of animals by exposure for 2 hours in an environment rising from 5° to 35° C (After Martin)

ment. If the body produced more heat than it gave off, the retained heat would cause a rise in temperature; if it lost more than it generated, its temperature would fall. The average daily heat production—that is, the metabolism—of a man following a light occupation is around 2500 Calories. This quantity of heat is dissipated in the ways described in the following section.




Fig. 9.5 Showing an arteriovenous anastomosis in the skin of an exposed part (e.g., ear, check, finger) through which warmer blood from deeper parts is conducted through the skin, just beneath the more superficial vessels. In I, the arteriovenous anastomosis is closed, and, as at ordinary environmental temperatures, blood is directed through the subspapillary venous plexus; in 2, the blood, as at low temperatures, is shunted through the communicating vessel, a small proportion only, passing through the more superficial vessels. The continuations of the arteriole into the subspapillary venous plexus and of the small vein leading therefrom are represented merely by double rows of dotted lines.

Physical factors in temperature control. 1. Radiation, convection, and conduction. Through these three physical processes the body, like any other warm object, loses heat to the cooler air surrounding it, and to any cooler objects in contact with it or in its immediate neighborhood. The heat lost in this way can be increased or diminished through variations in the quantity of blood flowing through the skin. On a hot day, for example, the vessels of the skin dilate; more blood is therefore brought from the deeper parts of the body to the surface, and heat loss accordingly increased. In cool weather, the vessels of the skin constrict, a greater proportion of the total blood volume being distributed to the internal structures; heat loss is thereby reduced. When the cold is extreme, the cutaneous vessels of covered parts of the body, such as the trunk and limbs, are constricted; but in the skin of the ears, nose,

checks, etc., anastomosing vessels existing between the arterioles, on the one hand, and small veins on the other dilate to permit a free flow of warm blood from deeper parts to the skin of the exposed regions; thus their temperature is prevented from falling too low. The anastomosing vessels lie beneath the subpapillary venous plexus from which they divert a considerable proportion of the swiftly flowing blood (Fig. 9.5 and Pl. 6 A). The skin is kept warm more effectively in this way. Thus the skin may appear pale or even of a bluish tint though quite warm; for it is the amount of blood flowing through the venous plexus and not that in the deeper cutaneous vessels which gives the skin its color (see capillaries of skin, Chap. 4).

It should be mentioned that under ordinary circumstances the amount of heat lost through conduction is negligible. Air is a very poor conductor, but the body immersed in water or on cold ground, especially if wet, loses heat rapidly by conduction.

Another important factor in temperature control is the variation in blood volume that results from changes in environmental temperature. The blood is increased in volume by heat, and is thus capable of absorbing and conveying more heat to the surface of the body for elimination through radiation, convection, and conduction. Cold causes the reverse effect—namely, a reduction in blood volume and, in consequence, a diminished heat loss.

2. Evaporation of water from the lungs and skin. The latent heat of evaporation of water is about 0.6 Calories —that is, this quantity of heat is absorbed in the vaporization of 1 cc of water. Under ordinary atmospheric conditions, about 300 cc of water are vaporized from the lungs daily, and about 500 cc from the skin. This represents a heat loss of (800 × 0.6 —) 480 Calories

But it should be understood that the evaporation of water from the surface of the body occurs constantly in the absence of any visible sweating, and even though the sweat glands are quite inactive or, as in some rare instances, when they are entirely lacking. Like any other moist mass, the body is continually losing weight by evaporation. The water "seeps" (diffuses) into the skin from the underlying tissues and is vaporized from the surface. This water loss, together with that from the lungs, was first observed and studied by an Italian physician Sanctorio (1561–1636), upon himself and called by him insensible perspiration. The quantity of body water lost from the skin surface as visible sweat and as insensible perspiration varies, not only with the temperature of the air, but also with the degree to which the

<sup>4</sup> The latent heat of the evaporation of sweat is 0.59 Calorie,

atmosphere is saturated with moisture—i e, with the relative humidity If this is very high, a large part of the sweat may simply drip from the skin, and such unevaporated sweat exerts no cooling effect. The relative humidity of the atmosphere is, therefore, of the greatest importance with respect to our comfort on a hot day.

It is clear that when the environmental temperature is higher than that of the body, heat cannot be dissipated by radiation and convection the body would gain rather than lose heat were it entirely dependent upon these processes. At such temperatures, evaporation of water from the skin and lungs plays the leading role in the regulation of body temperature. The sweat glands are stimulated and visible sweating appears when a rise of from 0.5 to 1°F in body temperature occurs. The rise in body temperature may result from an increase in the tem perature of the environment or from increased heat production, as in muscular exercise.

The effect upon the sweat glands is brought about through a center in the brain from which they receive impulses through the sympathetic nerves. A rise in temperature of the blood supplying the center is the most potent factor in causing the secretion of sweat. This is shown by the fact that heating the carotid artery in the cat causes sweat secretion from the toe pads, though the paws themselves are kept cool. But sweating may also be induced reflexly—that is, from the stimulation of afferent nerves in the peripheral tissues skin muscles, and so forth

Table 9 6
Daily heat loss

•	
	Calories
Radiation, convection and conduction	1850
Evaporation from the skin	400
Vaporization of water from the lungs	175
Warming inspired air to body temperature	80
Urine and feces (i.e., heat of these excreta over that	
of the food)	50
Total daily heat loss	2555

In dogs and cats, functional sweat glands are absent from the body surface, being confined to the skin of the pads of the paws. In these animals, rapid breathing (panting), with increased vaporization of water from the lungs as a consequence, is relied upon to increase heat loss at high temperatures.

3 Of the total heat lost from the body, over 95 percent is elimi

nated through (1) and (2) discussed above. The remaining 5 percent is lost in the urine and feces, and in raising the inspired air to body temperature.

The quantities of heat lost daily in each of the several ways just described are given (in round numbers) in Table 9-6.

Variations in the quantity of heat lost through these physical factors are capable, under ordinary circumstances, of maintaining the body temperature at the normal level. However, at very high environmental temperatures or when, as in strenuous muscular effort, heat production is very greatly increased, the mechanisms of temperature control may be inadequate; the body temperature then rises.

The chemical factors in temperature control. At low environmental temperatures, the physical mechanisms alone are incapable of reducing heat loss sufficiently to prevent a fall in body temperature. Chemi-




Fig. 9.6 Showing the effect of external temperature upon metabolism. (From Martin after Gessler from observations upon himself.)

cal regulation then comes into play—i.e., heat production increases. The body's fires, so to speak, are fanned. The point in the temperature scale where physical factors are aided by chemical means in the control of body temperature is called the critical temperature. It varies, of course, with the nature and thickness of the clothing, but for the naked human body it is around 28° C (82° F). Cold, therefore, acts as a powerful stimulus to metabolism (heat production). The greater heat production in response to cold is brought about mainly by increased tone of the skeletal muscles and, in some instances, by invol-

untary contractions of these muscles (shivering) and of the smooth muscle of the skin (goose flesh) When cold, a person usually indulges, also, in some form of muscular activity—walking, swinging his arms, or stamping his feet—which very materially increases his heat production (see Fig 9 6)

At temperatures between 28° and 30° C, the body (naked) is able to establish a balance between heat production and heat elimination without calling accessory mechanisms into play—sweating on the one hand or muscular contraction (shivering or voluntary muscular movements) on the other—and one, therefore, feels quite comfortable This temperature range, or correspondingly lower temperatures for the clothed body, is for this reason called the comfort zone

In cold climates protein food, owing to its specific stimulating effect upon metabolism (specific dynamic action, p 359), gives valuable assistance to the chemical mechanism of temperature control. On the other hand, a high protein diet is unsuitable in hot weather, since the greater heat production which it induces throws an additional bur den upon the mechanisms responsible for heat dissipation.

Heat-controlling centers. The main center for the control of body temperature is situated in the fore part of the hypothalamus—that region at the base of the brain near the origin of the pituitary stalk (p 541) Isolation of this region by section of the brain stem or upper spinal cord renders an animal poikilothermic—that is, incapable of maintaining the height of its temperature independently of the temperature of the environment. This operation also paralyzes the skeletal muscles, the power to increase heat production is therefore lost Stimulation of the hypothalamic center causes a rise in temperature. The center exerts its influence upon body temperature through the autonomic nervous system causing vasoconstriction or vasodilatation, sweating, increased muscular tone, contraction of smooth muscle in the skin (shivering), and the liberation of adrenaline from the adrenal medulla which, as mentioned below, stimulates heat production

These effects upon the sweat glands, vessels, and the adrenal gland are largely reflex in nature, the center being influenced by afferth impulses initiated in the temperature receptors of the skin. But such effects are also induced by a rise in the temperature of the blood supplying this region of the brain

The endocrines and heat regulation. The secretions of both the thyroid and adrenal glands stimulate heat production. This effect, known as the calorigenic action of these hormones, enables an animal to with stand a greater degree of cold than is possible for one that has been deprived of either or both of these endocrines; the temperature of such animals is usually subnormal. Prolonged exposure to cold (over a period of weeks) causes an overproduction of the thyroid hormone. At any rate, the thyroid glands of rats who have been so treated show pronounced indications of increased activity, and a rise in heat production. The latter effect does not appear in thyroidectomized animals similarly treated.

The hypersensitivity of hypothyroid patients to cold and of hyperthyroid to a warm environment is well known.

Fever or pyrexia. A rise in body temperature above the normal level, unless of a temporary nature as in strenuous exercise or as a result of exposure to a high air temperature, is called fever or pyrexia. The highest body temperature compatible with life is between 110° and 112° F. The rise in temperature is due primarily to an impairment of the mechanisms for heat elimination (e.g., increased blood volume, vasodilatation, and sweat secretion) and not to increased heat production At the commencement of a fever, for example, the skin is pale and dry; the metabolism is not increased. Heat elimination is reduced; heat is therefore retained in the body and the temperature

rises. Having reached a certain height, the temperature remains fairly steady, for now heat elimination keeps pace with heat production; a balance is struck, but at a higher level than in health. Though, as just mentioned, increased heat production is not responsible for the onset of the fever, the higher temperature induced through heat retention




Fig 9.7 Chart from Du Bois showing the effect upon metabolism of an increase in body temperature (fever).

causes, in turn, greater heat production; for the heat generated in the body is the result of chemical reactions, mainly oxidative in nature and, like chemical reactions in general, they are accelerated by a rise in temperature (see Fig. 9.7). A rise of 1° F in body temperature increases the metabolism by about 7 percent.

In acute infectious diseases and in bacterial infections generally, the accompanying fever is due to an action of the bacterial toxin upon the heat-regulating centers; in animals whose brain has been cut across below these centers, the injection of a toxin, which in a normal animal would cause a sharp rise in temperature, does not cause fever. In

some instances, the bacterial toxin may act directly upon the heat regulating centers. As a rule however, its action is probably an indurect one, the first effect being to produce chemical changes in the tissues which alter the osmotic pressure of the tissue fluids, causing the withdrawal of water from the vascular system and, as a consethe withdrawal of water from the vascular system and, as a consequence, a reduction in the dissipation of heat by radiation Furthermore the reduced blood volume causes the drainage of blood from the cutaneous vessels, the skin thus becoming cold, stimulates cold receptors, and brings about a reflex response from the centers controlling heat production Cutaneous vasoconstruction occurs, which further reduces heat dissipation accompanied by shivering. One of the first symptoms at the onset of an acute infectious fever is a feeling of in tense cold, often with goose flesh, shivering chattering of the teeth and shaking of the limbs for, as just mentioned, the withdrawal of warm blood from the skin has stimulated receptors sensitive to cold. The temperature taken in the mouth at this time of the "chill is found to be high though the skin may feel cold even to another person, and the patient's lips and nose appear pinched pale and even of a blush tint. The mechanism of a chill at the commencement of a fever may be compared (though not exactly) to that controlling the tempera be compared (though not exactly) to that controlling the tempera ture of a room isolated from a temperature sensitive device which has been exposed to a cold draft The cold sensitive device corresponds to the cold receptors in the skin the thermostat to which it is connected by wires and which controls the furnace, is comparable to the heat

by wires and which controls the turnace, is comparative to the acceptability and the isolated room to the deeper tissues. The effect of certain drugs upon the body temperature. There are many drugs which have a fever reducing or antipyrine, and quinne. Some of these drugs appear to bring about their antipyrietic effect by increasing the blood volume, and thus increasing heat dissipation, for the greater volume of fluid available for circulation through the vessels near the body surface, the greater the amount of heat which will be radiated to the surroundings. The increase in blood volume is thought to be due to the rise in blood sugar which these drugs induce, water is drawn into the vessels by the raised osmotic pressure of the blood plasma caused by the increased sugar concentration. On the other hand, reduction in circulating fluid, as in dehydration, may cause fever. Other drugs such as morphine alcohol in large amounts and general aneithetics (ether, chloroform etc.) are capable of reducing even a normal body temperature through their depressant action upon the heat-controlling centers.

There are other agents which raise the body temperature above the normal. Dinitrophenol, a drug which has been used—not without danger—as a cure for obesity, raises the body temperature by increasing the rate of oxidation in the tissues. The action of the hormones, adrenaline and thyroxine, in this regard, has been referred to on pages 413 and 422.

Heat stroke, heat exhaustion, and sunstroke. After a person has been exposed to a high temperature for a time, his heat-regulating mechanism may fail, and very serious symptoms, even death, may result. The effects of extreme heat fall into two categories known, respectively, as heat stroke and heat exhaustion. In the former, a person exposed to great heat, often while undergoing some physical exertion, suddenly becomes unconscious. The body temperature is found to be very high, and in a severe case may reach 108° or 110° F. Sweat secretion is suppressed; both blood volume and urine volume are increased. In heat exhaustion, the symptoms develop more gradually and consist of abdominal cramps, rapid pulse, low systolic blood pressure, and sometimes dizziness and dyspnea. The blood and urine volumes are reduced, and the chloride of the blood and urine are much below normal (due to loss of chloride in the sweat). Sunstroke is merely a form of heat stroke or heat exhaustion, the body absorbing heat from the direct rays of the sun. There is no special effect of the sun's rays, apart from their heat, which causes the symptoms of so-called sunstroke.

Heat cramps. Men working in a very hot environment for a long period may suffer severe cramplike contractions of the muscles. The muscles are rendered hyperexcitable by the reduction in the salt concentration of the tissue fluids; and this is due, in turn, to the excessive loss of sodium chloride in the sweat.

## The Vitamins

It was discovered nearly 50 years ago that diets composed of purified protein, fat, and carbohydrate, and containing the essential minerals would not support life. It was concluded, therefore, that foods in their natural state contained substances which, though present only in minute amounts, were nevertheless essential to life. Sir F. Gowland Hopkins, of Cambridge University, was among the first to suspect the presence of such materials in natural foods. In 1906, he expressed his views in the following words, "No animal can live upon a mixture of pure protein, fat, and carbohydrate; and even when the necessary in384

organic material is carefully supplied, the animal still cannot flourish." The chemical nature of these materials was unknown, so they were called simply accessory food factors. Later these vital elements were thought to belong to a group of nitrogenous substances known as amines They were therefore named vitamines. It was soon found, however, that this conclusion was false. The name was retained, however, but it was suggested that the terminal "e" be dropped. The generic term vitamin was then generally adopted. There are a number of different vitamins. Each is designated by a letter of the alphabet. The vitamins A. D. E. and K are soluble in fats and fat solvents (the fat-soluble vitamins); the vitamins of the B complex and vitamin C are soluble in water (the water-soluble vitamins).

The growth of our knowledge of the vitamins within the past three decades ranks as one of the most brilliant achievements of physiological and biochemical research, the practical application of this knowledge has conferred immense benefits upon mankind.

#### Table 9.7

### Summary of the vitamins

Vitamin A (antixerophthalmic)

thiamin or vitamin B, (antineuritic)

riboflavin, B<sub>2</sub> or G
nicotinic acid or niacin
pyridoxin, B<sub>6</sub>
pantothenic acid
folic acid
folic acid
antipernicious anemia factor, B<sub>12</sub>
biotin

inositol para-aminobenzoic acid

Vitamin C (antiscorbutic)

Vitamin D (antirachitic)

Vitamin E (antisterility)

Vitamin K (antihemorrhagic)

Vitamin A (C20H10O), the antixerophthalmic vitamin. The chief sources of vitamin As are fish liver oils (e.g., cod and halibut), dairy

There are actually two vitamins A Each is derived from a separate carotene (carotene a and carotene b), which differ slightly chemically. The vitamin A first discovered and present in mammalian livers, and in the commoner types of food is now known as vitamin A, it is derived from caroteners. The second vitamin A, products, and many vegetable foods. The following is a list of materials especially rich in this vitamin.

Fish liver oils
Butter and cream
Egg yolk
Carrots, yellow maize (corn), alfalfa, spinach, peas, beans, and other
vellow and green vegetables

It is not strictly correct to say that plants contain vitamin A. They contain rather a yellow pigment called *carotene* ( $C_{10}H_{10}$ ) which, after being taken into the animal body, is converted by the liver into vitamin A according to the following equation:

#### $C_{40}H_{44} + 2H_4O = 2C_{40}HO_{20}$

Carotene, being a precursor of the vitamin, is called provitamin A. The green and yellow parts of plants contain more of the provitamin than do the paler portions. The contents of vitamin A in milk and butter varies with the carotene content of the fodder of the cow. After ingestion the carotene is transformed in part to vitamin A and secreted as such in the milk. A part of the carotene is also secreted unchanged. Vitamin A itself is colorless, so that a pale milk may be just as rich a source of the vitamin as one more deeply colored. The vitamin A of fish liver oils is also derived ultimately from plant life. Small invertebrates of the sea, known as copepods, feed upon marine plants, and serve in turn as food for small fish. The latter convert the carotene, originally formed by the plant, into the vitamin. Larger fish devour the smaller fish and store the vitamin so obtained in their livers.

Vitamin A deficiency. When the diet is deficient in vitamin A (and in the provitamin), the epithelium lining the respiratory and alimentary tracts as well as the ducts of certain glands, becomes abnormal; it undergoes transformation to the stratified squamous type. Such nuccous surfaces are especially susceptible to infection. Inhibition of lacrimal secretion, followed by drying of the cornea (xerophthalmia), is an outstanding effect of vitamin A deficiency. In some instances softening of the cornea (keratomalacia), which may progress to ulceration and consequent blindness, results. Vitamin A is necessary for the regeneration of the visual purple of the retina which, in health, readily takes place after the eyes have been exposed to light (Chap. 13).

designated  $A_3$ , is absent from mammalian liver but is the chief vitamin A found in the livers of fresh water fish; its provitamin is carotene- $\beta$ .

Night blindness (nyctalopia) is, therefore a common accompaniment of vitamin A deficiency A sufferer from night blindness sees well in bright light, but is quite or almost blind in dim light Vitamin A deficiency may also lead to degenerative changes in the long fiber tracts of the spinal cord and in certain of the peripheral nerves

of the spinal cord and in certain of the peripheral nerves
An adequate supply of vitamin A is also required for the normal
growth of bone In experiments upon growing animals, pronounced
localized overgrowths of bone results, causing deformities. These ab
normalities were particularly noticeable in the bones of the skull and
spinal column Pressure upon nerves issuing from the crainal cavity or
vertebral column by the redundant bone are largely responsible for
the abnormalities of the nerves mentioned above

The diets of persons on this continent are rarely deficient in vita min A to the point where serious abnormalities supervene, but in In dia China, and other Eastern countries the diet is frequently deficient in this and other vitamins. In localities far removed from larger centers of population owing to difficulties of tránsportation such as in



Fig 9 8 Xerophthalmia in a child caused by vitamin A deficiency (Drawn from a photograph after Block)

lumber camps of the north country, especially in winter, night blindness and other signs of vitamin A deficiency occasionally make their appearance Infants reared upon a badly planned artificial diet may also suffer from deficiency of vitamin A (Fig. 98)


The viamin B complex The first of this large group of vitamins to be discovered was thiamin or vitamin B; the antineuritic vitamin Severe deficiency of this vitamin is the cause of a disease known as beriberi Beriberi is a disease of rice eating countries. Its chief features are paralysis of the limbs (due to de

generative changes in the peripheral nerves—polyneurits), dilation of the heart and edema. It was shown by Eikiman a physician work ing in the Dutch West Indies to be due to a dietary deficiency. He demonstrated that the essential factor missing from the diet was contained in the coverings of the rice kernel. Persons who live almost exclusively upon a diet of polished rice—that is rice from which the pericarp (bran) and germ have been removed in the milling process—

develop the disease, but are readily cured when they are given unpolished rice or an extract prepared from the rice "polishings." This factor in the polishings of rice which protected against beriberi was designated vitamin B or the antineuritic vitamin. From researches of



Fig. 9.9 Upper figure (after Harris); left, distended stomach in thiamine (B,) deficiency due to loss of tone of gastric muscle; right, normal stomach Lower figure: curves showing effect of graded quantities of vitamin B, upon growth of nine groups of rats; P, point where polyneuritis developed, broken lines are drawn from the points where some individuals of the group died. Figures from 0.0 to I.0 indicate grams of whole wheat (B,) received by the respective groups. (After Chase.)



recent years it has been shown that so-called vitamin B is actually a complex substance consisting of several vitamins, only one of which is protective agains beriberi. This latter, the antineuritic principle, is now known as vitamin B<sub>1</sub>, or thiamin.

Other effects of vitamin B, deficiency are (a) gastrointestinal disorders (e.g., reduced motility of the stomach and intestines resulting in constipation, inhibition of the digestive secretions, and loss of appetite); and (b) retarded growth (see Fig. 9.9).

In birds (e.g., pigeons) a condition analogous to human beriberi







Fig. 9.10 Left, pigeon suffering from polyneuritis; right, the same bird an hour after treatment with vitamin B., (After Funk, redrawn)

results from vitamin B, deficiency. Its chief features are paralysis due to involvement of the peripheral nerves (polyneuritis), retraction of the head, and convulsions. The condition is cured in a remarkably short time by feeding a concentrated preparation of vitamin B, (see Fig. 9.10). Vitamin B, deficiency in a dog and its rapid cure by administration of the vitamin is shown in Figure 9.11.

Thiamin has been synthesized. It exists in the tissues as an integral




Upper drawing, dog with polyneuritis showing paralysis of the hind limbs due to a deficiency of vitamin B. Lower drawing, the same animal cured by the vitamin given in the form of tomato juice (Drawn from a photograph after Cowgill and Mendell )

part of an enzyme system essential for the metabolism of carbohydrates. In its absence, carbohydrate metabolism does not proceed normally. Lactic acid accumulates in the blood, heart, and brain. The accumulation of lactic acid in the brain substance appears to be responsible for the head retraction seen in birds.

There are several other vitamins of the B complex (see Table 9-8, p. 396), the most important of which in respect to human nutrition are riboflavin and nicotinic acid (niacin).

Riboflavin. This is one of a group of yellow fluorescent pigments found in animal tissues. Liver, kidney, and milk are especially rich sources of this factor of the B complex When it is lacking from the diet, reddening of the eyes, due to the growth of fine blood vessels

Fig. 9.12 Vascularization of the cornea due to severe riboflavin deficiency. Note the growth of vascular loops into the cornea, which normally is bloodless, from the vessels at the periphery (lumbic plexus).



into the cornea, results, together with opacities of the cornea (Fig. 9.12). A sensation of burning and itching of the eyeballs is experienced. There is intolerance to light which may be so severe that the eyes cannot be opened in an illumination of even ordinary intensity (photophobia). Another effect of deficiency of this vitamin is the development of fissures in the skin radiating from the angles of the mouth.

Animals with riboflavin deficiency grow slowly. Chicks on a diet in which it is lacking develop a degenerative disorder of the peripheral nerves, a type of paralysis resulting known as "curled toe."

Nicotinic acid (niacin). A disease occurring among the poor of the southern United States and less commonly in some European countries has within recent years been recognized as being due, mainly at least, to a deficiency of this vitamin. The chief features of this disease, which is known as pellagra, are reddening and drying of the skin together with, in the severer cases, nervous and gastrointestinal disturbances It occurs in those who subsist mainly upon maize

Pyridoxin Rats on a diet deficient in pyridoxin develop a cutaneous

disorder consisting of reddening and scaliness of the skin and loss of hair There is some indication that this factor is concerned with the synthesis of hemoglobin and the manufacture of erythrocytes. The regeneration of blood following hemorrhage is hastened by feeding pyridoxin Deficiency of this vitamin has not been observed in man

Pantothenic acid Most animal and vegetable tissues contain this acid It stimulates the growth of yeast Its role in human nutrition is unknown But when lacking from the diet of chicks, it causes der matitis and degenerative changes in the nervous system. In rats, pantothenic acid deficiency causes graying of the fur and 'spectacle eyes" (an appearance due to an inflamed hairless ring around the eyes), as well as damage to the kidneys, heart, and adrenal cortex

Biotin This vitamin is present chiefly in yeast, liver, kidney, and the yolk of egg and in smaller amounts in other foods. Its deficiency does not occur in man so far as is known but it is important, nevertheless in human nutrition, for deficiency has been produced experimentally in man by giving a diet containing a high proportion of raw white of egg Unpleasant symptoms including drowsiness, greasiness of the skin and some cardiac distress, developed In rats fed upon a diet of raw egg white, a severe dermatitis accompanied by "spectacle eyes appears A protein in raw egg white known as avidin combines with biotin in the intestinal tract and renders the latter unavailable in nu trition, the avidin biotin compound cannot be utilized

Choline A constituent of the lipid known as lecithin, choline when lacking from the diet results in fatty livers in animals. Deficiency of choline apparently does not occur in man. Nor has inositol deficiency been observed in man This vitamin is found chiefly in cereals as a complex compound with calcium, phosphorus, and magnesium It is also present in animal tissues such as liver and ordinary meats. Mice fed upon a diet deficient in inositol show patches of baldness, but it does not appear to be related to loss of hair in man Para aminobenzoic acid deficiency has not been observed in man and, though deficiency in rats causes graying of the hair, there is no evidence that this vitamin is related in any way to the loss of color of human hair Para amino-benzoic acid's chief interest lies in the fact that it is necessary for the

growth of certain disease producing bacteria (see footnote, p 221)

The role played by folic acid in blood formation is described on

page 74, and of vitamin Biz on page 76

A number of vitamins of the B complex are essential for the growth of several types of microorganism and are dietary requirements for chicks, rats, mice, and other laboratory animals, but only thiamin, riboflavin, niacin, folic acid, and vitamin  $B_{12}$  have been shown to be indispensable constituents of human diets. This does not necessarily mean that those which are not required in the diet are not utilized by the human body. Several are synthesized by intestinal bacteria, and such synthesis probably occurs in the human intestine. Most of the factors of the vitamin B complex have been shown to exist as integral parts of tissue enzyme systems with which their functions are intimately associated.

Vitamin C, the antiscorbutic vitamin. Vitamin C is present in greatest amounts in citrus fruits (e.g., lemons limes oranges, and grapefruit). It is also contained in most other fruits and green vegetables. Tomatoes and red and green peppers are exceptionally rich sources, potatoes, turnips, meat, and milk contain moderate amounts. Vitamin C has been identified chemically as ascorbic acid (CeH2O4). It possesses high reducing powers, it has been synthesized in the laboratory, and by the intestinal bacteria of many animals. Vitamin C serves the important function of maintaining a healthy state of the walls of the capillaries. When it is lacking from the diet,

Vitamin C serves the important function of maintaining a healthy state of the walls of the capillaries. When it is lacking from the diet, the cement substance interposed between the endothelial cells, and holding them together, becomes deficient. The vessels, as a consequence, develop leaks which permit the escape of blood into the surrounding tissues.

Vitamin C is in relatively high concentration in the cortex of the adrenal gland (see p 428)

Scurvy (scorbutus) is caused by vitamin C deficiency. The chief manifestation of this disease is bleeding from the mucous membranes, beneath the skin and into joints. The skin, as a result of numerous capillary hemorrhages (petechiae), may show extensive mottling. The guins are swollen and inflamed and bleed easily. The subjects become anemic, weak, and emaciated. Unless fresh food containing the vitamin is supplied, death results. Scurvy, though unusual in the general population except in times of famine, was a common disease in sailors, soldiers, explorers, and others who, of necessity, subsisted upon diets lacking in fresh fruits, vegetables. meat, and milk. Scurvy is readily produced in guinea pigs by placing them upon a diet of hay and oats——i.e., one lacking in green stuffs. The disease is sometimes seen in persons living on the outskirts of civilization, such as lumbermen, especially during the winter months when fresh food is difficult to pro-

302 METAROLISM AND NUTRITION

cure, and provision for a supply of vitamin C in concentrated form has not been made. The disease may also appear in infants who are artificially fed; infantile scurvy (Barlow's disease) should, however, never occur, since the administration of tomato or orange juice is a certain preventive.

Vitamin D (C<sub>2</sub>,H<sub>4</sub>,O), the antirachitic vitamin. This vitamin is formed by the action of ultraviolet light upon certain waxy compounds known as sterols. Among such compounds are ergosterol, found in yeast, and cholesterol, present in the skin and other animal tissues. The ultraviolet rays cause a rearrangement of the atoms in the sterol molecule, thereby endowing it with the properties characteristic of vitamin D Vitamin D is formed naturally by the action of sunlight upon a sterol in the skin closely associated with cholesterol. This sterol is known chemically as 7-dehydrocholesterol. The antirachitic vitamin is prepared commercially by irradiating yeast (rich in ergosterol) by means of the mercury vapor lamp or the carbon arc. Such artificial preparations of vitamin D are referred to as viosterol, or simply as uradiated or activated ergosterol. The wavelengths in the ultraviolet part of the spectrum which are effective in the formation of this




Fig 9 13 Scheme of wavelengths of spectrum, showing the range of those possessing antirachitic power. Figures in millimicrons (mu). (Redrawn and modified from Blunt and Cowan.)

vitamin have a range of from 250 to 313 mu\* (see Fig. 9.13). Vitamin D has been synthesized in the laboratory. It is a yellow crystalline substance which has been named calciferol. The richest natural sources of the vitamin are fish liver oils, especially those from the cod and halibut. These large fish obtain the vitamin from smaller species upon which they feed (e.g., herring, whiting, etc.). The small fish in turn acquire the vitamin from invertebrate marine forms, such as copepods and mollusks, which probably synthesize it. Some authorities believe that the cod and certain other fish are also capable of synthesizing

<sup>\*</sup> mu = millunicron, the one millionth part of a millumeter

vitamin D. There are several vitamins D. The two of medical interest are the two just mentioned. They are designated  $D_2$  (irradiated 7-dehydrocholesterol of skin and in cod liver oil and other fish oils) and  $D_1$  (irradiated ergosterol, calciferol). There is no vitamin  $D_1$ .

The following is a list of the chief sources of vitamin D.

Fish liver oils (e.g., halbut and cod)
Irradiated ergosterol (of yeast)
Irradiated milk
Irradiation of the skin

The main action of vitamin D is upon the metabolism of calcium and phosphorus; it is essential for the mineralization of growing bone, and for maintaining the normal mineral composition of the adult skeleton. It also plays an important role in dental health; defective tooth structure, leading to decay of the teeth (dental caries), results when the diet is deficient in the antirachitic vitamin. When the vitamin in the form of viosterol is administered in excessively large doses, a marked rise in the calcium of the serum (hypercalcemia) results, calcium being drawn from the skeleton to the blood. The calcium of the urnne is increased. Vitamin D acts, in this respect, like the hormone of the parathyroid glands. Toxic symptoms resembling those caused by overdosage with parathyroid extract also follow the administration of large amounts of viosterol. The manner in which the vitamin, in physiological dosage, exerts its effect upon calcium metabolism is not altogether clear. But it is generally believed to increase the absorption of calcium from the intestinal tract; its chief action, however, appears to be upon the skeleton, influencing, in some manner not understood, the deposition of calcium and phosphorus in the developing bone.

Rickets (rachitis) results from vitamin D deficiency. It is a disease of young children, chiefly, between the ages of 6 months and 3 years. Its essential feature is defective development of the bones. The deposition of phosphorus and calcium is interfered with; the bones are, therefore, soft and yielding. Deformities of the limb bones and spine and defective ossification of the skull bones result. Small round swellings are found at the junctions of the ribs with the sternum; these are referred to as the "rachitic rosary." The head is somewhat larger than normal with a prominent brow; soft areas may be felt in the skull. The abdomen is large and round, which has given rise to the term "potbelly" (see Fig. 9.14). The disease occurs most often in bottle-fed infants, especially in those who are growing rapidly and receiving a diet containing a large proportion of carbohydrate. Rickets is readily cured




Fig 9 14 Diagram showing the chief features of infantile rickets (After Harris redrawn)

or prevented by the administration of vitamin D, either in the form of cod liver or halibut liver oil, viosterol or irradiated milk, or by exposing the child to ultraviolet radiations (sunlight, the mercury vapor lamp, or carbon arc) The value of sunshine in the treatment of rickets was pointed out over 70 years ago (1880) by Palm, an English physician. but it is little more than 30 years since the curative effects of ultraviolet irradiation and of cod liver oil have come to be generally recognized

Rickets is seen most commonly in the large cities of northern latitudes, since infants in these localities are less exposed to sunshine than are those in southern climates. The disease is almost unknown in the tropics. The smoke and dust of cities also act as a screen through which the shorter wavelengths of the sun's rays cannot penetrate, or do so to a minimal extent. It should be remembered, furthermore, that window glass is opaque to radiations shorter than 320 mm. Sunshine that has passed through ordinary glass therefore possesses no antirachitic action On the other hand, in order to gain benefit from ultraviolet radiation a person need not be in direct sunshine. The sun's rays reflected from the sky (skyshine), or from water, snow, or light colored objects are effective

Osteomalacia (soft bones) is a disease-essentially the same as rickets—which occurs in adults in eastern countries especially India and China Women are mainly affected with the disease since they, as well as receiving in common with the general population littly via min D in their diet, are often confined within doors and thus deprived of the benefit of sunshine

Vitamin (or vitamins) E, the antisterility vitamin, Vitamin E is necessary for normal reproduction. Female rats upon diets lacking in vitamin E though they become pregnant do not bear living young The embryos after developing for a short time die In male rats, The embryos after developing for a short time die in male rats, vitamin E deficiency results in degenerative changes in the testes, spermatozoa are not produced and the sex instinct is lost. The addition of lettuce or other food containing vitamin E corrects the defects of either the male or female reproductive functions.

Vitamin E is fat soluble and has been obtained in crystalline form. Its chief sources are green vegetable foods—e g lettuce peas, alfalfa and the germ of various cereals. Wheat germ oil has a very high vita.

min E content

Besides the vitamin E first discovered two other compounds closely related to it chemically, have been shown to have similar physiological actions We may now speak of vitamins E. All three belong to a class of compounds known as the tocopherols. The three vitamins E are, therefore frequently referred to by their chemical designations Pre

therefore requently referred to by their chemical designations. Fre fixing the Greek letters to distinguish them, they are known as alpha, beta and gamma tocopherol, respectively.

Viamin E exerts an influence upon the metabolism of muscular tissue (nutritional disorders of muscle occur in animals when it is lacking from the diet) and upon the biochemical processes in the nuclei of cells generally, its action is exhibited especially upon rapidly growing

Table 9 B
Summarizing the principal facts concerning certain vitamins

λame	Ch of functions	Efects exused by deficiency	Chief sources
l tam n A ant zerophthalm e	Mainta as the integrity of epithel al tusues Essential for the regen eration of visual purple.	Aerophthalm's Keratomalacia Convers on of columnar epithelium to equamous type N ght blundness	Carrots Letture Alfalfa Yellow corp. Butter cream and mill Egg yolk
Thism s (B and rin) antineut tic	Essential in the metabo- ium of carbohydrate serving as a coensyme in the ties on Main ta as the tone of the gastrointest as! tract. Promotes appetite	Beriberi Polyncuritis in animals (taperially hards) Loss of tone of gas- trointest nal tract Loss of appet te	Yeast, germ of when and other cereals.
Riboflavin (B <sub>1</sub> vitamin G)	Essent al component of certain tusue ensymes	Vascularization of cor nea with opacities Fize area around 1 ps Retarded growth Loss of hair (n rata)	Liver Ridney Mak.
Niac a)	Coenzyme in t said res- pirat on	Pollagra	Liver meats Brower s yeast, Wheat germ
Pyridox a (B <sub>4</sub> )	Coenryme in the me- sabolism of tyros ne tryptophane and cer tain other amino ac da	Retarded growth (in rate) Anomis in pigs dogs and rabbits The effects of sta lack in human nutrit on unknown	Mik. Wheat germ Yeast Aleast kidney and veg table fats.
Pantothenie acid	St mulates the growth of yeast.	Spectacle eyes and graying of fur in rats anemia in dozs de- fic ency in man not observed.	Wdely distributed animal and vegetable insures
E otm	Tiesus enzyme concerned in carbohydrate metab- elism	Spectacla eyes and nerrous (spast c) symp- toms in rats Deficiency (escept ex perimental) not ob- served in man Deficiency produced by feeding raw egg white	Yeart, kidney liver at egg yolk
Para-am nobemso o acid	Essential for some en syme system in becta- rial reli (see footnote p 2-1)	Graying of fur in black rate Deficiency not observed in man	Various animal tissue cultures of certain be- teria (e.g. Staphyl- cocci)
Vitamin C antiscorbutic	Maintains the integrity of the capillary walls	Seurvy	C true fruite. Tomatoes Turnips
Vitamin D antirachitin	Regulates calcium and phosphorus metabo- lism Essential for the normal development of bones and teeth	Rickets, Osteomalscis, Dental carles	Cod fver and hal be liver mis Exposure of the skin f ultraviolet irradiation
Vitamin E antistorility	Essential for reproduc- tion.	Death of young in uterus. Steril ty of male de- generative changes in testes failure of sper- matagenesis	Latture Water crees. Wheat germ
Vitamia K Antibemerrhagie	Necessary for the pro- duction of prothrom his by the liver	Hemorrhagia tendency	Alfalfa and other gree foods


structures. It has been employed, but on the whole with disappointing results, in the treatment of certain wasting diseases of the muscles; and, in consideration of its influence upon the reproductive functions, it has been given to pregnant women who have had repeated miscarriages or abortions (habitual abortion) with the hope that they will be able to give birth to a living child.

Vitamin K (the antihemorrhagic vitamin). This vitamin is found in largest amounts in alfalfa, spinach, cauliflower, cabbage, and other green foods and less abundantly in cereals, carrots, and yeast. Chicks and other farm birds develop a fatal hemorrhagic disease when this vitamin is lacking from the diet. The hemorrhages are due to a low concentration of prothrombin in the blood. This, as mentioned on page 103, is an essential factor in the coagulation of the blood. Hemorrhages in the human subject are rarely if ever due to the lack of vitamin K in the diet because vitamin K is synthesized by bacteria in the intestine of man. For the same reason it has not been found possible to induce a hemorrhagic tendency in other mammals by feeding them a diet lacking in this vitamin. But vitamin K is not absorbed from the intestine in the absence of bile. Therefore, in obstructive jaundice in which bile does not enter the intestine, a tendency to bleed is a common feature. It is now the custom to give vitamin K by injection or by mouth accompanied by bile or bile salts before operating upon a patient with obstructive jaundice. The prothrombin concentration of the blood is raised thereby and the defect in the clotting mechanism corrected.

A grave hemorrhagic disease occurs in newborn infants due to an insufficient amount of vitamin K received by the fetus from the mother. The prothrombin concentration of the infant's blood, when this occurs, is less than 50 percent of the normal. Death may occur within the first few hours after birth. If the child is untreated but survives longer than this, the condition usually corrects itself, for bacteria swallowed in the milk synthesize the vitamin in the intestine. The disease can now be treated at the first sign of its appearance, and quickly corrected by the intravenous administration of vitamin K.

## The Principles of Dietetics

The importance of a well-balanced diet for the normal growth and development of the young, and for the health of both young and old, is recognized today as never before. There seems little doubt that the finer physique and better health of the present generation is largely the result of the more varied and ample character of modern diets (see



Upper photograph class of elementary school children at Southwark England in 1894 Lower photograph corresponding class 30 years later The improvement of this group over that in the upper photograph is attributed largely to the more adequate diet received by the children in the lower group (After Harris)

Fig 9 15) Deficiency diseases which have been so prevalent in the past, especially among the poor, are now comparatively rare Rickets and scurvy, for example, have almost vanished from civilized com munities and, were the knowledge which has been gained from modern research sedulously applied, the eradication of these and other diseases due to defective diets-such as beriberi and pellagra-would soon be an accomplished fact. In planning a diet, the following factors must be taken into account.

399

- 1. The caloric (energy) requirement of the subject.
- The proportions of the three main types of food carbohydrate, protein, and fat.
  - 3. Essential fatty acids.
- 4. The vitamin requirement.
- 5. The mineral constituents.

Determination of the caloric requirement. A man of average size and following a light occupation has an average energy output of from 2500 to 3000 Calories daily. In order that the body's tissues shall not be called upon to furnish any of the required energy, the caloric value of the diet must not be less than the energy expenditure; otherwise the body would lose weight Nor should the caloric value of the diet be greater than the energy expended. The excess energy would then be simply stored as fat (see p 374). Excess of energy intake over energy output is the usual cause of obesity. In order, therefore, to calculate the caloric requirement in any given case, one must first know the energy expenditure during the resting state, that is, the basal metabolism (p. 356). To this is added the energy expenditure of the particular occupation in which the person is engaged.

The energy expended daily by different persons, or by the same person at different times, varies widely, depending upon the extent of the exercise undertaken. The energy allowances for work of several grades

are given in the following table.

Dail	y energy expenditure
	(above basal)
	Calories
Sedentary life	500~ 700
Light work-e.g, professional and business-	
men	700-1200
Moderate work—e.g., mechanics Heavy work—e g, laborers, lumbermen,	1200-1500
athletes, etc.	1500-4000 and upward

The metabolism during sleep is lower than the basal metabolism by from 15 to 20 percent. The lower metabolism of the 8 hours of sleep must therefore be taken into account in determining the caloric requirements. The calculations just outlined are exemplified in the following table in which the figures are for a man following a light occupation.

#### METABOLISM AND NUTRITION

400

Basal metabolism (16 hours)	1200 Calories 1	
Metabolism during sleep (8 hours)	500 "	
Allowance for light work	800 "	
Total	2500 Calories	

Since the food itself has a stimulating effect upon metabolism (see specific dynamic action, p. 359), an allowance of from 6 to 10 percent must be made in order to establish the energy balance. In the example given above, this would amount to between 150 and 250 Calories. If this allowance is not made, the subject would lose weight, for his own tissues would be drawn upon to furnish the extra heat production associated with the specific dynamic action.

The basal metabolism of a normal subject can be derived from his height and weight (see chart, Fig. 9.3, p. 358). The energy values of the various articles of diet are obtained from standard tables, and the quantities apportioned accordingly. For example, beef (with a medium amount of fat) has an energy value of about 1300 Calories per pound; butter (almost pure fat) yields 3600 Calories per pound; milk, 325; eggs, 765; cheese (largely protein), 2200; sugar, 1860; and white bread, 1200. Children, owing to their greater activity and the requirements of growth, should receive a more ample diet in proportion to their size than that of the average adult A boy of 16 years, for example, especially if he is growing rapidly and taking part in athletics, requires at least as much food as a full-grown man engaged in moderately heavy work

The proportions of carbohydrate, fat, and protein. A little over 50 percent of the total caloric requirement should be furnished by carbohydrate food, about 35 percent by fat, and 12 percent by protein. The protein in the diet of the average adult should be from 70 to 90 grams daily, or around 1 gram per kilogram of body weight. Children, especially in the early years of life, require a higher protein intake (2 to 3.5 grams per kilogram) than do adults, and a larger proportion (60 to 90 percent) of the total protein should be of the highest biological value, such as is furnished by milk, eggs, and meat. In adults receiving 80 to 90 grams of protein daily, at least 50 percent should be first class, as in meat and eggs. If the protein is mainly of low biological value (e.g., vegetable proteins), the total allowance must be larger.

Fat is an indispensable element of the diet. Fats not only contain fatsoluble vitamins, but also certain unsaturated fatty acids (linoleic and

<sup>7</sup> This is about 40 Calones per hour for a man having a surface area of 1.8 square meters.

linolenic acids) which cannot be synthesized in the body, yet are essential for normal nutrition

The mineral requirements. Sodium, potassium, magnesium, and phosphorus are present in adequate amounts in a diet which is ample in other respects and, as a rule, no special attention need be paid to them Sodium chlqride, for example, besides that which is present nat urally in many foodstuffs is added in cooking, and as table salt in quan tities determined by individual taste. Protein foods constitute the chief source of the phosphorus of the diet and, when the allowance of protein is adequate, the phosphorus intake takes care of itself. Potassium and magnesium are derived from cereals and vegetables, and are also present in sufficient amounts in an ordinary diet.

The minerals in which the diet is most likely to be deficient are cal cium, iron and iodine. The intake of calcium, which enters so largely into the composition of bone, is especially likely to be inadequate in the diets of children. According to Professor Sherman, children require at least a gram of calcium per day. The adult requirement is about 0.8 gram daily. As a result of the deposition of mineral in the bones of the fetus, the demand for calcium increases during pregnancy when the allowance should be from 1.5 to 2 grams daily. The mother should also receive a high allowance of calcium after childbirth if she nurses her baby at the breast, for the secretion of milk is a severe drain upon the body's calcium stores. By the same token milk is especially rich in calcium, containing about a gram per quart. Cereals and certain vegetables (such as beans, peas, and turnips) are also good sources of this mineral, but the calcium of milk, as compared with that of vegetables, is more readily absorbed. Only a small part of the calcium in cereals and vegetables is utilized by the body. Meat contains minimal amounts Milk is therefore, especially for children, the best source of calcium.

mineral, but the calcium of milk, as compared with that of vegetables, is more readily absorbed. Only a small part of the calcium in cereals and vegetables is utilized by the body. Meat contains minimal amounts. Milk is therefore, especially for children, the best source of calcium. Iron is an indispensable constituent of the diet, since it is necessary for the synthesis of hemoglobin. The daily requirement is from 15 to 20 mg. The chief sources of food iron are meats (especially liver), eggs, and such vegetables and cereals as spinach, beans and peas, whole wheat, and oatmeal Milk is very poor in iron. Iodine is an essential constituent of the thyroid hormone, goiter (p. 412) results when the diet is deficient in this element. Sea foods are the chief natural sources of judine, though many brands of table salt.

Iodine is an essential constituent of the thyroid hormone, goiter (p 412) results when the diet is deficient in this element. Sea foods are the chief natural sources of iodine, though many brands of table salt (rodized salt) contain small quantities (1 part in 100 000) which have been added by the manufacturer. The daily requirement of iodine is placed at about 100 micrograms.

<sup>•</sup> A microgram or gamma (γ) = 1000 milligram.

#### 402 METABOLISM AND NUTRITION

water is the most usual source of this mineral.

to be taken in minute quantities for special purposes—e.g., the elaboration of certain hormones and enzymes or for the incorporation into particular tissues. Among these so-called "trace elements" are: cobalt (in vitamin B<sub>13</sub>, p. 74), zinc (in insulin, p. 371), manganese (in tissue enzymes), and copper (required in hemopolesis). No attention need be paid to these elements in planning a diet since the small amounts required are present in a diet adequate in other respects. Traces of fluorine, which is a constituent of osseous and dental tissues, are indis-

pensable for the formation of teeth of normal structure. The drinking

There are a number of other mineral elements which are required

# The Endocrine Glands

Glands such as those of the mouth, stomach, and intestines, which deliver their secretions into the alimentary tract, or those such as the lachrymal, mammary, and sweat glands, which discharge upon the surface of the body, are called glands of external secretion or exocrine glands. The glands which we are about to consider do not possess ducts or any openings to the exterior, their secretions pass into the blood stream, and are thus conveyed to the various tissues of the body upon which they exert their action. These glands are therefore called endocrine glands or organs, ductless glands, or glands of internal secretion.

The secretions of the endocrine glands are for the most part excitatory in their actions, they stimulate the growth and development, or the functional activity of certain tissues; internal secretions acting in this manner are therefore called hormones (Gk. hormaō, I excite). The term hormone is not, however, restricted to the secretions of the ductless glands. Any substance formed by a tissue of the body and carried in the blood stream to act as an excitant to some other tissue or organ may be called a hormone. Thus, secretin (p. 325), pancreozymin, and even carbon dioxide (p. 254), since it is carried by the blood and acts upon the respiratory and vasomotor centers, come into this category. Chalone is an allied term, sometimes used to designate an internal secretion having an inhibitory action, such as enterogastrone (p. 314).

Methods of investigation. Our knowledge of the endocrines has been advanced enormously in the past thirty years, and especially within the last twenty. The chief methods used in the investigation of endocrine function are three in number. First, an extract of the gland tissue may be prepared, which is then injected into animals and a study made of its effects. Second, a given gland may be excised, and the subsequent life history of the animal followed, careful notice being taken of its development and growth, or of any unusual feature. Third, studies may

be made upon human subjects in whom one or other gland is known to be deficient or overactive. The first and second methods are often employed together. For example, after extirpation of the gland, the animal is treated with an extract prepared from the same type of gland, observations being made of the power of the preparation to correct the defects resulting from the operation. This is called the substitution or replacement method.

The general chemical nature of the hormones. The active principles of the endocrine glands have a complex chemical structure. Those of the male and female sex glands and of the adrenal cortex belong to the large class of fatty and waxlike substances known as steroids (p. 373). Their chemical formulae are well known and many have been synthesized in the laboratory. The hormones of the pituitary gland, the parathyroid hormone, and insulin, on the contrary, are proteins, little is known more precisely of their chemical constitution. The thyroid hormone, thyroxine (see p. 412), is an amino acid.

Potent extracts of a number of ductless glands have now been prepared These preparations, with the exception of thyroid extract and some forms of the male and female sex hormones, are relatively or quite inactive when given by mouth, their full effect is exerted only when given by injection

Most of the ductless glands are present in all orders of vertebrates, and an extract obtained from the gland of one order exerts, as a rule, its specific effect when administered to a member of another order. Thus the hormone of the sheep's thyroid influences the growth and development of frog larvae (tadpoles)

## The Thyroid Gland

The thyroid gland is composed of two lobes which he one on eithe side of the larynx (see Figs 10 1 and 10 2) Under the microscope, the glandular tissue is seen to be composed of a mass of alveoli (also calle acun or follucles) lined by a single layer of cuboidal epithelial cell. The alveolar cavities are filled with a homogeneous viscous matericalled colloid this is secreted by the lining cells and contains the thy roid hormone.

The thyroid is supplied with nerves, but whether these have an secretory function is questionable. It is certain, at any rate, that a hoi mone liberated by the pituitary gland (thyrotrophic hormone, p. 435 is of much greater importance than are nerve fibers in the control  $\epsilon$  thyroid activity

Excision of the thyroid (thyroidectomy) in young mammals (e.g., pigs, goats, calves, rabbits, etc.) results in retardation of skeletal growth, and failure of sexual development (Fig. 10.3) Tissue oxida tions are depressed, the basal metabolic rate falling after the operation to 60 or even 50 percent of the normal. The skin becomes thick and leathery in texture, the hair sparse, dry, lusterless, and brittle. The




Fig 10 1 Showing the relation of the thyroid gland to neighboring structures




Fig 10.2 Microscopic section of normal thyroid tissue

long bones continue to grow in thickness but longitudinal growth is very slow. The administration of thyroid extract after thyroidectomy prevents the onset of these effects or, if they have appeared abolishes them or at least arrests their progress.

The effects following thyroidectomy in lower orders are even more striking Thyroidectomized tadpoles for example do not develop into frogs. Metamorphosis is at once resumed however upon the addition of thyroid extract to the water in which the thyroidess animals live (Fig. 10.4). Furthermore, the administration of thyroid extract to nor mal tadpoles causes them to metamorphose in about a third of the usual time. The effect of thyroid extract upon the axoloil—an aquatic form allied to the frog—is still more extraordinary. In the adult form, this animal is purely aquatic in its habits possessing gills a finned tail, and four short limbs, suggesting some fabulous gigantic tadpole whose metamorphosis, after proceeding so far, has become arrested (Fig.

40Å

10.5). Thyroid feeding causes this creature to lose its gills and fin, to develop air-breathing organs, and forsake the aquatic life of its kind.

Goiler. This term is applied to chronic enlargement of the thyroid. When the enlargement is not associated with any general effects referable to disturbed function of the thyroid, it is called simple gotter. Goiter may, however, be accompanied by underfunctioning or by over-




Fig 10.3 The effects of thyrordectomy on growth and development. The kid on the right was thyroidectomized soon after birth, both kids are the same age. (Drawn from a photograph after Sutherland Simpson)

functioning of the thyroid—hypothyroidism and hyperthyroidism, respectively.

The histological appearance of the goitrous thyroid varies In one type, the alveoli are enlarged, irregular in size and shape, and filled with colloid material. This is called simple colloid goiter (Fig. 10 6). Colloid goiter may progress to atrophy of the secretory cells, and their replacement by fibrous tissue; the signs of thyroid deficiency (hypothyroidism) then make their appearance. In a second type, the cells lining the alveoli enlarge and multiply; the wall of the alveolis is thrown into folds which project into, and almost obliterate, its cavity. This form, which is known as diffuse parenchymatous goiter, may be associated with signs of increased functional activity of the gland (hyperthyroidism). In the third type, the gland contains a tumor of thyroid tissue (adenoma); the surrounding glandular tissue may be normal. This form is called adenomatous goiter. It too may give rise to hyperthyroidsm.

All forms of simple goiter are due to a deficiency of iodine in the food and drinking water. Other forms of goiter, in many instances at any rate, are also believed to be due to some deficiency of this element. It is, therefore, in regions where iodine is low, such as those far




Fig 10.4 (Above) Showing the effect of thyroidectomy upon the metamorphosis of tadpoles A thyroidecto mized B normal frogs of the same hatching (After Allen redrawn)



Fig 10.5 (Right) Axolotls A has not been treated B has received thyroid extract Note the disappearance of the gills and fin in the latter

removed from the sea (which contains large stores of iodine), that gotter is prevalent The inhabitants of mountain villages—e.g., in the Alps, Pyrenees, and Himalayas—have been among the chief sufferers Through the ages iodine has been leached from the soils of these re gions and, as a consequence, the water and food have a very low iodine content. In certain parts of North America, especially in areas near the Great Lakes, and in some inland and hilly counties of England, there

**∡**∩8

is a poverty of iodine and goiter is occasionally seen, even today. Since the importance of iodine for normal thyroid function has been recog nized, small amounts of iodine are taken by the population of goitrous districts, as a result of this preventive measure, goiter is much less




Fig 10.6 Types of gotter as defined by microscopical appearance A colloid gotter B parenchymatous gotter C adenomatous gotter

prevalent today than in the past Iodine, as we shall see presently (p 412) is an essential constituent of the thyroid hormone

Hypothyroidsm (thyroid deficiency) In infants and young children thyroid deficiency produces effects essentially similar to those seen in animals following thyroidectomy. The defects of physical sexual and mental development form a characteristic picture to which the term cretinism is applied the child is called a cretin. These young victims of thyroid deficiency are dwarfed to a marked degree and unless the condition is corrected at an early age by the administration of thyroid extract or of thyroxine (p. 412), their height when they reach adult age




Fig 10.8 Cretinism On left at age of 3½ years (before treatment) right same subject (after treatment with thyroid hormone) at the age of 7 years (Drawn from a photograph after McCullagh)

**41**∩

may be no greater that that of a normal child of 7 or 8 years. The sexual organs do not develop, nor do the secondary characters of sex appear. The basal metabolic rate is lowered. Mental growth is, as a rule, very seriously retarded; many cretins are imbecile or idiotic, and a large proportion are deaf mutes (Figs 10.7 to 10.9). The facial features are coarse; the bridge of the nose is depressed; the tongue in many instances is enlarged and, as if too large for the mouth, protrudes



Fig 10.9 A group of cretins in an institution (After Falta)

between the lips. The skin of the entire body appears thick and puffy, due to the deposition of a gelatinous fluid in the subcutaneous tissues. The hair is dry, brittle, and sparse.

Cretinism is very commonly associated with goiter, and is therefore most prevalent in goitrous districts; for as already mentioned, goitrous enlargement may lead to atrophy of the secretory cells. Cretinism, if recognized early, can be corrected by continued treatment with thyroid extract, powdered thyroid gland, or thyroxine. Almost normal growth, and sexual and mental development can be induced (Fig. 10.8). The longer that treatment is postponed, the less benefit will be derived; if the condition has existed untreated for years, little improvement can be expected.

Myxedema. This condition is the result of thyroid deficiency commencing in adult life. Its general features—namely, the low metabolic rate, the thickness and puffiness of the skin (myxedema) from which the condition originally got its name, the scantiness and dryness of the hair—are similar to those just described for cretinism. But inasmuch as full development—skeletal, sexual and mental—has been attained before the onset of the disease, many of the characteristic features of cretinism are absent. Myxedematous subjects, though they are apathetic and lethargic and think slowly, are not mentally defective.





Fig 10 10 Myxedema Left before treatment with thyroid extract Right after treatment (Drawn from a photograph after Joll)

The myxedematous subject is very quickly restored to normal by thyroxine or thyroid extract.

Hyperthyroidism, thyrotoxicosis (overactivity of the thyroid gland) Hyperthyroidism occurs in two forms, considered on the basis of the microscopical appearance of the gland. In the first form, called exophthalmic goiter, the whole gland is enlarged. The chief features of this condition are (a) enlargement of the thyroid (b) elevation of the basal metabolic rate, (c) disturbances of carbohydrate metabolism (e.g., reduction in the glycogen stores hyperglycemia, and glucosuria), (d) nervousness and tremor, (e) rapid action of the heart and (f) protrusion of the eyeballs (exophthalmos, see Fig. 10.11).

<sup>2.</sup> Exophihalmos or protrus on of the eyes often to the extent that the whites of the eyes show above the trix, is due to the accumulation of fluid (edema) in the tissues behind the globes The eyeballs are literally pushed forward and may be parily dislocated from their sockets in extreme cases the lids cannot be closed over them The surfaces of the eyes then become dry and inflamed and may utcerate Exophihalmos is not due as might be thought to an excess of the thyroid hormone itself, but to the thyrotrophic hormone of the anterior lobe of the pitutary body (p. 435) it can be produced in thyroidectoms ed animals by the continued administra tion of this hormone.



Fig. 10.11 Exophthalm c go ter Note the staring eyes and the enlarge ment of the thyro d

The name given to this type is not always appropriate since there may not be exophthalmos and this may be present in the second form called toxic adenoma in which the gland contains one or more tumors of thyroid tissue (adenoma) surrounded by normal appearing thyroid tissue. In both types the affected thyroid tissue shows signs of great activity presenting those features described on page 406 as parenchymatous gotter.


The thyroid hormone It was shown over 60 years ago that the thyroid was a gland of internal secretion. An English physician George Murray in 1891 prepared a glycerin extract of sheeps thyroids. This preparation when ad

ministered by injection to a subject of myxedema restored him to normal within a very short time. It was discovered later that thyroid tissue dried and powdered and given by mouth was equally effective. At this time nothing was known regarding the chemical nature of the hormone but it was soon found that thyroid extracts contained iodine combined with protein. This iodine protein compound was called thy reoglobidin (or thyroglobidin). The active principle of the extract was later obtained in crystalline form. It was shown to contain a large proportion (60 percent) of iodine it was named thyroxine. Thyroxine—the thyroid hormone—has since been synthesized It is an amino acid derived from the amino acid tyrosine and has the following formula.

Thyroxine whether prepared from thyroid tissue or synthesized has an action identical with that of the dried gland but is much more petent A fraction of a milligram given daily is sufficient to maintain the basal metabolic rate of a subject of myxedema at the normal level

The thyroid hormone appears to act as a catalyst (p. 290), hastening very greatly oxidation in the tissues. When given to a normal animal, the metabolic rate is raised and the other signs of hyperthyroidism (except exophthalmos) make their appearance. The effects are not observed, however, until the lapse of several hours after the hormone has been administered, and are not fully developed until the lapse of from 8 to 10 days. But, even though a single dose is given, the effects persist for 5 or 6 weeks.

The production of the thyroid hormone and its discharge into the circulation are automatically controlled through the hypothalamus and the anterior lobe of the pituitary (thyrotrophic hormone, p. 435). This controlling mechanism is a very complicated one, several factors are



concerned, but probably the most important one is the concentration of the thyroid hormone in the blood. A fall in the concentration of the thyroxine in the circulation stimulates the hypothalamic-pituitary mechanism, and more of the thyroid-stimulating (thyrotrophic) hormone is released from the pituitary. Any tendency for the thyroid hormone concentration in the blood to increase beyond the body's requirements causes a depression of the production of the thyrotrophic hormone. Thus the thyroid hormone governs its own production (see

414

Fig 10 12) It is also probable that an excess of thyroid hormone in the circulation acts directly upon the thyroid, "backfires" as it were, to depress hormone production Furthermore, it is most likely that the nervous system, through the hypothalamus, exerts an influence upon the production of the thyroid stimulating hormone of the pituitary. Summary of thyroid function. The thyroid gland extracts iodine from

Summary of thyroid function. The thyroid gland extracts iodine from the blood together with the amino acid, tyrosine, and from these synthesizes its hormone thyroxine, whose molecule, as will be seen from the formula shown above, contains four atoms of iodine (1) Thyroxine presumably acts upon all itssues as a catalyst, hastening oxidative processes. It is tempting to attribute the developmental retardation caused by thyroid deficiency to the reduction in the rate of tissue oxidations, but the problem is not so simple as this, for dinitrophenol, a drug which increases oxygen consumption by the tissues and raises the metabolic rate, is not capable of replacing thyroxine in a subject of thyroid deficiency. The thyroid, apart from increasing the oxygen consumption, exerts some specific effect upon the tissue cells

# The Parathyroid Glands

The parathyroid glands are two pairs of small oval structures (about a quarter of an inch long) lying in close relation to the thyroid. The upper pair is situated, one on each side of the mid line, behind the up




Fig 10 13 The thyroid region of the neck from behind

per poles of the thyroid lobes, the lower pair hes behind, or a little below, the inferior poles of the thyroid (Fig 10 12) Sometimes the parathyroids, especially the upper pair, are embedded in the posterior part of the thyroid


Removal of the parathyroids (parathyroidectomy) causes increased excitability of the neuromuscular tissues, which culminates in a severe convulsive
disorder known as tetany. The
manifestations of tetany, as they
occur in animals after parathyroidectomy, are as follows (a)
witching and spasmodic contractions of the muscles, which

usually increase in violence until the whole body is thrown into convulsions; (b) rapid breathing; (c) acceleration of the heart; (d) rise in temperature; (e) death, usually occurring from exhaustion or from asphyxia resulting from spasm of the muscles which close the opening of the larynx (laryngeal spasm).

Parathyroid deficiency resulting in tetany is sometimes seen in the human subject following removal of the thyroid for goiter, the parathyroids having been inadvertently or unavoidably removed with the thyroid, or severely injured. The phenomena of parathyroid deficiency in man are usually much less pronounced than those which follow parathyroidectomy in animals Twitchings, nervousness, and an occasional spasm of the facial or limb muscles are often the only manifestations. In other instances, tetany is in a latent form and is revealed only by means of special tests or under special circumstances (p 417).

The function of the parathyroid glands. Normal blood contains from 5 to 6 mg of calcium per 100 cc; this is all contained in the plasma, so that the concentration of calcium in the plasma (or in the serum, for

Fig 10 14 The effect of parathyroid extract on the calcium of the blood in parathyroid tetany. A single dose of extract was given when the serum calcium had fallen to 6 mg percent and the symptoms of tetany had reached their height (After Macleod and Taylor.)



none is removed in the clot) is from 9 to 11 mg per 100 cc (i.e., 6 mg per 100 cc of blood). Parathyroidectomy causes a pronounced fall in the serum calcium (hypocalcema). When the serum calcium level falls to about 6 mg percent, tetany appears and increases in severity with further reduction in the serum calcium concentration. Injections of calcium relieve the condition within a few minutes; there is no doubt, then, that calcium deficiency and the development of the tetanic state are directly related. These observations led to the conclusion that the parathyroids are concerned in some way with the control of calcium metabolism. This view of parathyroid function was amply confirmed in 1925, by the important discovery made by Professor J. B. Collip.

He found that an extract prepared from the parathyroid glands of cattle, when administered by subcutaneous injection to parathyroidectomized animals, raised the serum calcium and abolished all the manifestations of tetany (Fig. 10.14).

When normal animals are given repeated doses of parathyroid extract, the serum calcium is raised above the normal level (hypercal-



Fig 10 15 Carpopedal spasm. The positions of the hands and feet in tetany

cemia); it may be increased to 20 mg or more per 100 cc of serum. The excretion of calcium and phosphorus in the urine and feces is increased. The excess calcium in the serum is derived from the bones. The skeleton, especially the ends of the long bones, contains stores of calcium and, under normal circumstances, these are released by the action of the parathyroid hormone. Thus the normal level of calcium in the blood is maintained

Other forms of tetany. Tetany occurs from other causes than parathyroid deficiency. It is not uncommon in young children (infantlle tetany), and is then usually an accompaniment of rickets, the parathyroids being very rarely at fault. The serum calcium, however, is low and, as in parathyroid tetany, it is the hypocalcemia which is directly responsible for the tetanic seizures. Tetany is also seen in osteomalacia (p. 395), an adult form of rickets occurring in China and India; in sprue, a rare condition associated with defective absorption of calcium from the intestine; and in certain other conditions but especially in alkaloxis, which may result from persistent vomiting (loss of HCl), forced breathing (excessive elimination of  $CO_2$ ), or the administration of large doses of an alkali, such as sodium bicarbonate.

In the tetany of children, the hands and feet assume characteristic attitudes (see Fig. 10.15). This phenomenon is referred to as *carpopedal spasm*. Spasm of the laryngeal muscles with asphyxial attacks are common; the child becomes blue (cyanosis), but, presently, the obstruction to breathing is overcome, the air being then drawn into the chest with a highpitched "crowing" sound. In other instances, generalized convulsions occur.

In either infants or adults, the serum calcium is sometimes reduced below the normal level, but not quite low enough to bring on a tetanic attack, under ordinary circumstances. This is called latent tetany. A typical tetanic seizure may be precipitated, however, by some upset in the general health. Latent tetany may be detected by the following tests: (a) Tapping over the facial nerve in front of the ear causes twitchings of the muscles of the face, if latent tetany exists. This is called Chvostek's sign. (b) Erb's sign. The muscles show increased excitability to stimulation by a galvanic current (applied to the skin overlying a muscle). (c) Trousseau's sign. Tying a tourniquet around the arm above the elbow, so as temporarily to occlude the blood supply, causes the hand to assume the characteristic attitude.

Hyperparathyroidism (excess of the parathyroid hormone). In animals, overdosage with parathyroid extract causes (besides the hypercalcemia and the increased excretion of calcium and phosphorus, already mentioned) very grave effects—e.g., depression of spirits, muscular weakness, and hemorrhages into the stomach and intestines. Death occurs within a few days. If given in smaller doses over a long period, softening of the bones, as a result of the removal of calcium, results. In man, there is little doubt that excessive amounts of the extract would produce similar effects.

A rare disease, known as osteins fibrosa cystica, is caused by overactivity of the parathyroid glands themselves. A tumor of one or other
of the four glands is usually present; this secretes excessive amounts of
the parathyroid hormone into the blood stream, producing effects
closely similar to those resulting from overdosage with parathyroid
extract—namely, hypercalcemia, increased excretion of calcium and
phosphorus in the urine, and softening of the bones, which in some
cases leads to marked deformity. The bony tissue, which normally is
dense and contains a high percentage of mineral, becomes demineral-




Fig. 10.16 Longitudinal section of bone (tibia) showing structural changes in osteits fibrosa cystica (Drawn from a photograph) Note deformity of shaft and cystic cavities in the head of the bone

ized and more fibrous in character. In places, the bone structure is re placed by cavities (cysts) of various sizes and shapes (see Fig. 10.16) Renal stones (calculi) are common in this disease and are due to the deposition of calcium excreted in excess by the kidney. They are formed from part of the excess calcium brought to the kidney for excretion.

# The Adrenal Glands (or Suprarenal Capsules)

These are two somewhat pyramidal structures lying, one on each side of the body, in contact with the upper pole of the kidney (Fig 10 17) Each gland measures from 1 to 2 inches in length. The adrenal




Fig 10 17 The adrenal

gland, however, is actually two glands in one, for each consists of two functionally distinct parts-a central portion which is called the medulla, and a surrounding zone of tissue called the cortex The cells of the medulla give a specific chemical reactionnamely, an affinity for the salts of chromic acid, by which they are stained brown They are therefore referred to as chromaffin cells In certain fish (the elasmobranchs) the two parts are quite separate (Fig. 10 18) The tissue corresponding to the mammalian adrenal medulla is present as several small masses of cells lying in close relation to the ganglia of the sympathetic chain In man, also, small collections of chromaffin cells are to be found in close association with the

sympathetic ganglia. They are known as paraganglia. The two parts of the adrenal gland have quite different origins, and there is but little evidence that their functions are closely related. The medulla develops from a group of cells, split off from the neural crest of the early em-




Fig 10.18 Adrenal tissue in elasmobranch fish (A), in mammals (B) medul lary (chromaffin) tissue black cortical tissue white The small masses of chromaffin tissue lying close to the sympathetic ganglia are the paraganglia

bryo, to which the ganglion cells of the sympathetic nervous system also trace their ancestry. It is important to remember this fact in order to understand the functions of this part of the gland. The cortex, on the other hand arises from the multiplication of cells in close relation to those from which the sex glands (testes and ovaries) are developed A microscopical section of the adrenal cortex is shown in Figure 10.19



Fig 10 19 Microscopical section of adrenal cortex and adjacent part of medulia

The cellular structure of this part of the gland shows three distinct zones—an outer (zona glomeru-losa), a middle (zona fasciculata), and an inner lying next the me dulla (zona reticulars)

The functions of the adrenal medulla. Over 60 years ago (1894), two physiologists in Edinburgh (Professor Schafer and Dr Oliver) showed that an extract of the adrenal medulla caused a pronounced rise in blood pressure when injected subcutaneously or intravenously The active principle of the extract was later isolated in pure crystal line form and finally synthesized It was named epinephrine, but adrenaline is the name more commonly used today Adrenaline is related to the amino acid, tyrosine, it has the following structural formula.

The extract of the adrenal medulla is prepared commercially, and is marketed under various names—

adrenalin, suprarenin, etc —depending upon the manufacturer It was discovered a few years ago that the adrenal medulia contained another hormone which was similar to but not identical with addrenaline. This was named noradrenaline or norepinephrine. The commercial extract is, therefore, a mixture of the two hormones. The description which follows applies to the actions of the commercial extract.

Actions of a commercial extract of the adrenal medulla. The rise in blood pressure following the injection of a commercial extract of the adrenal medulla (Fig. 10 20) is the result of constriction of the small

<sup>&</sup>lt;sup>2</sup> Adrenalm as a name for a commercial extract is spelled without the terminal e When referred to as a separate hormone it is spelled with the e

blood vessels (arterioles), chiefly of the skin and abdomen. The vessels of the heart (coronaries) and of the skeletal muscles are dilated by the extract. The heart is slowed; this action is the reflex result of the rise in blood pressure (Marey's law, p. 190); for if the vagus nerves are first cut, the heart is accelerated. The quantity of the extract required to cause cardiac acceleration is minute. The denervated heart—that is, the heart completely isolated from nervous control by interruption of the vagus and sympathetic pathways—is increased in the rate of its




Fig. 10.20 Showing effect of adrenaline upon intestinal contractions, A, and blood pressure, B 1 cc of 1 in 10,000 adrenaline solution injected at X. (Modified from Jackson)

beating by as little as 1 part of the material in 1,400,000,000 parts of blood. The denervated heart therefore provides the most sensitive test object for detecting the presence of adrenaline in blood or other fluid.

Other effects produced by an extract of the adrenal medulla are inhibition of the movements of the intestine, and contraction of the pyloric, iliocolic, and internal and sphinciers. It relaxes the smooth muscle of the bronchioles, thus causing dilatation of these parts of the bronchioles in attacks of great value in relieving spasm of the bronchioles in attacks of asthma. It inhibits the wall of the urinary bladder, but causes contraction of the ureter, and of the sphincier of the bladder. In most animals, the uterus contracts under its influence. It dilates the pupil of the eye and contracts the smooth muscle of the skin through which the hairs or feathers are elevated. It exerts a favorable effect upon the contraction of skeletal muscle; an isolated muscle subjected to repeated stimulation fatigues less readily if treated with the extract (Fig. 10 21).

### 422 THE ENDOCRINE GLANDS

An extract of the adrenal medulla also raises the metabolic rate, this is spoken of as its calorigenic action. It increases the sugar content of the blood by causing an increased breakdown in the liver of glycogen to sugar. The decomposition of muscle glycogen is also accelerated, lactic acid being produced which, escaping into the circula





Fig. 10.21. Showing the effect of adrenaline in postponing muscular fatigue. Note the increase in the height of the contractions following each injection of adrenaline.

tion, raises the level of the lactic acid in the blood. The lactic acid is carried to the liver and there reconverted to glycogen. These effects in their relation to the carbohydrate cycle are shown in the following scheme.



Adrenaline hastens the clotting of blood, but this effect is brought about only if the hormone is injected into the body. It exerts no effect upon the clotting process when added to blood after it has been shed—i.e., when added to blood in a test tube. This difference in the action of adrenaline upon the clotting process within and outside the body suggests that the hormone acts in the former case, not directly, but through some other tissue or organ acting as an intermediary. The nature of the intermediary is unknown

Most of the effects listed above as following the administration of an adrenal extract can be produced by stimulation of sympathetic

nerve fibers. Indeed, the actions of the hormones of the adrenal medulla may be summed up by saying that they imitate the effects of sympathetic excitation. This remarkable sympathomimetic action is perhaps not so surprising when we recall that the adrenal medulla and sympathetic nerve cells have a common origin (p. 419). See also p. 566.

The actions of adrenaline and noradrenaline compared. Informa-tion concerning the separate actions of adrenaline and noradrenaline is not yet complete, but the following comparisons can be drawn. Noradrenaline is a general vasoconstrictor, showing little or no selective action in respect to different sets of blood vessels, whereas adrenaline, while it constricts the cutaneous vessels, dilates the coronary vessels and the vessel of the muscles. Its effect upon the vascular system as a whole is dilatation, the vasodilator action overshadowing the vasoconstriction. The peripheral resistance (Chap. 4) is therefore reduced. Since the overall effect of noradrenaline is vasoconstriction. it increases the peripheral resistance. The action of a commercial extract upon the heart is a pure adrenaline effect, noradrenaline having no effect upon cardiac action. The effect upon plain (smooth) muscle-stomach, intestine, bronchioles, etc -is also due mainly to adrenaline; noradrenaline has but little effect upon smooth muscle, nor does the latter hormone act to postpone the fatigue of skeletal muscle. The effects of the extract upon carbohydrate metabolism and upon heat production (calorigenic action) are chiefly due to adrenaline.

The emergency theory of the function of the adrenal medulla. By a great number of ingenious experiments, Professor Cannon and his several associates at Harvard University have demonstrated that the hormone of the adrenal medulla serves an emergency function. It reinforces the action of the sympathetic nervous system to raise the defense mechanisms of the body against the dangers and rigors of the environment to a high level of efficiency. These workers experimented with cats in which the heart had been denervated by cutting the vagus nerves and excising the stellate ganglion of the sympathetic chain which gives rise to or transmits the sympathetic fibers to the heart. It was found that fright (caused by a barking dog), exposure to cold, asphyxia, the administration of an anesthetic, pain, or muscular contraction caused acceleration of the heart. Since the cardiac acceleration could not have been caused by nervous influences, and did not

<sup>&</sup>lt;sup>3</sup> The heart receives directly a few fibers from the sympathetic chain below the stellate ganglion which must also be severed to completely denervate the heart.

occur after removal of the adrenals, it must have been due to the

occur after removal of the adrenals, it must have been due to the action of adrenaline discharged from the animal's own adrenal glands
Under quiet, resting conditions the blood contains only a negligible amount of adrenaline, about one part in one or two billion, but during states of excitement or any of the conditions just mentioned about 004 mg per kilogram of body weight may be discharged from the glands per minute. It is believed, therefore, that the adrenal medulla functions only under circumstances which demand some special effort on the part of the body, as when an animal is defending itself or running down its prey, fleeing from danger, or exposed to cold. It will be recalled that the reactions of an animal under such circumstances. eg, the bristing of fur or the ruffling of feathers (caused by the contraction of the smooth muscle in the skin), the dilatation of the contraction of the smooth muscle in the sam), the dilatation of the pupil, the acceleration of the heart, and, in certain cold blooded species, the contraction of the melanophores (pigment cells) of the skin—are characteristic of sympathetic stimulation or of adrenaline administration Other physiological adjustments which increase the efficiency of the body to the maximum in times of stress—such as the mobilization of sugar from the liver (fuel for muscular work), the rise in blood pressure, and the diversion of blood from the skin and splanch nic region to the vessels of the brain, heart, and contracting muscles, the dilatation of the bronchioles, and the postponement of muscular fatigue—are brought about through the sympathetic nervous system, as well as by the hormone of the adrenal medulla The increased coagulability of the blood caused by adrenaline would also seem to be a part of the general defense reaction, for should a wound be sustained,

The veretion of the adrenal medulla is under the control of the The vecretion of the adrenal medulla is under the control of the sympathetic nervous system Stimulation of the splanchine nerve causes the liberation of adrenaline, and prolonged stimulation causes exhaustion of the adrenaline stores. It is a very interesting and significant fact, in view of the functional relationship between the adrenal medulla and the sympathetic nervous system, that the fibers to the adrenal, unlike any other sympathetic fibers, do not form connections with ganglion cells but pass directly to the cells of the medulla. The adrenal cell thus appears to take the place of the sympathetic ganglion cell. A center, controlling adrenal secretion, is situated in the medulla oblongata. A higher center is present in the breokalanus, and in the oblongata A higher center is present in the hypothalamus, and in the cerebral cortex

When the blood pressure raising (pressor) effect of adrenal extracts was first discovered, it was naturally suggested that the adrenal glands

were responsible for maintaining the normal arterial blood pressure. The fallacy of such an idea is proved by the observation that destruction of the medullary tissue of both adrenals is not followed by a fall in blood pressure. Animals subjected to this operation show no abnormality. The adrenal medulla then, unlike the adrenal cortex, is not indispensable. Nor is there any evidence that so-called high blood pressure (essential arterial hypertension) is due to the secretion of excessive amounts of adrenaline However, a rare type of arterial hypertension, which occurs usually in paroxysms, is caused by the secretion of excessive amounts of adrenaline and noradrenaline. Certain tumors composed of adrenal tissue, or hypertrophy of the adrenal medulla itself, are associated with such hypersecretion of the hormones.

Sympathin. It has been shown by Professor Cannon and his colleagues that the sympathetic nerves bring about their effects through the medium of a chemical substance (or substances) liberated from their terminals. This substance, though resembling adrenaline in its actions, was found to differ in certain respects. In view of its origin at sympathetic nerve endings, it was called sympathin. It was later concluded that there were two such substances, one excitatory, the other inhibitory. It now appears that the first of these, which was named sympathin E, is noradrenaline and the second or sympathin I is adrenaline.

The functions of the adrenal cortex. Animals die in from 10 to 15 days after complete removal of both adrenal glands. We have seen that excision of the adrenal medulla alone is not fatal; the cortex, therefore, is the part of the gland which is essential to life. If one sixth or so of the total amount of adrenal cortical tissue is left, the animal survives; this part of the gland must, therefore, produce a highly potent hormone.

During the time that the animal survives after double adrenalectomy, the following effects are observed: (1) loss of appetite, vomiting, fall in blood pressure, muscular weakness and, just before death, profound prostration; (2) subnormal body temperature and reduced basal metabolic rate; (3) concentration of the blood, due to a loss of plasma water; (4) reduction in the sodium chloride of the blood and a rise in the potassium, calcium, phosphate, and nonprotein nitrogen; (5) hypoglycemia; (6) signs of renal failure.

Foremost among the functions of the adrenal cortex is the influence that it exerts upon the metabolism of water and salts. In the absence of the adrenal cortex, sodium and water are excreted in excessive amounts, whereas potassium escapes into the body fluids from the

tissue cells, where normally it is in relatively high concentration. Excretion of this mineral by the kidneys is also impaired, its concentration, therefore, rises in the blood and other tissues.

The adrenal cortex also exerts an influence upon carbohydrate me tabolism which is due to a hormone other than that controlling the metabolism of water and salt

Destructive disease of the adrenal cortex, usually of a tuberculous nature, occurs in man. The manifestations of this condition, which is known as Addison's disease are similar to, though less acute than those just described as resulting from adrenalectomy in animals. The disease if untreated, is gradually progressive, death occurring within 2 or 3 years. A bronze or dirty gray discoloration of the skin, which is not seen as a rule, in animals after adrenalectomy, is a prominent feature in Addison's disease. The skin may be mottled, pale areas of depigmentation lying adjacent to overpigmented areas. Anemia and gastrointestinal disturbances are also outstanding features of adrenal insufficiency in the human subject. The pigmentation is due to an in crease in the normal pigment (melanin) of the skin. The disturbances in water and salt metabolism already described as occurring in adrenalectomized animals are prominent features.

Addison's disease is treated by the administration of a hormone (usually desoxycorticosterone cortisone or both see below) obtained from the adrenal cortex (or prepared synthetically) in combination with a diet high in sodium content and low in potassium

The active principles isolvited from the adrenal cortex. Several active principles have been extracted from the adrenal cortex and prepared in crystalline form Chemically they all belong to the class of fatty or wax like substances known as steroids (p 372) and are referred to collectively as corticoids. They are basically similar in molecular structure to the sex hormones A hormone was isolated from the adrenal cortex some years ago by Dr. E. C. Kendall of the Mayo Foundation. It has a long chemical name but is known more briefly as Kendall's compound E. More recently, owing to the possible confusion with vitamin E. it has been designated cortisone. It was discovered by Dr. P. S. Hench and his associates of the Mayo Clinic that this principle exerts a most favorable effect upon the course of chronic (rheumatod) arthus and acute rheumatic fever.

The physiological action of cortisone is upon the redundant connective tissues (collagenous fibers) surrounding the affected joints. This principle has also been employed in affections of the skin associated

<sup>4 17</sup> Hydroxy 11-dehydrocorticosterone.

with disorders of the collagenous tissues. Besides the diseases mentioned, it has been used in a wide variety of pathological states (e.g., leukemia, asthma, multiple enlargement of lymph nodes, etc.), sometimes with considerable benefit, though the improvement is usually only temporary. The mode of action of cortisone in many of these disorders is not clearly understood. The secretion of cortisone, like that of

Table 10-1
The corticoids

Name	Principal actions
Corticosterone	On carbohydrate, protein, and fat metabolism; antagonizes the action of insulin.
11-Desoxycorticosterone	On water and mineral metabo- lism, increases the retention of sodium and the excretion of potassium
Aldosterone	Similar to preceding but more pronounced
11-Dehydrocorticosterone	On carbohydrate metabolism
17-Hydroxycorticosterone	On carbohydrate metabolism; on connective tissues, antitheumatic.
17-Hydroxy-11-dehydrocorticosterone (cortisone)	On carbohydrate metabolism; on connective tissues; antirheumatic
17-Hydroxy-11-desoxycorticosterone	On connective tissues; antirheu- matic.

other hormones of the adrenal cortex, is under the control of the adrenocorticotrophic hormone (ACTH) of the pituitary body. So far as is known, the latter hormone is the sole means whereby the function of the adrenal cortex is controlled. Unlike the adrenal medulla, the liberation of its hormones is not regulated by nerves.

Among the other steroid principles obtained from the adrenal cortex are corticosterone and desoxycorticosterone. The former exerts its chief action on the metabolism of carbohydrate, causing the formation of sugar from protein, increase in the glycogen stores, and hyperglycemia. The action of desoxycorticosterone is directed mainly to the metabolism of sodium and potassium, increasing the retention in the



Fig. 10 22 Child aged 41 years suffering from adrenal tumor. Note the florid "beefy" face and the precocious development of the genital structures. (Drawn from a photograph after Guthrie.)

body of the former, and stimulating the renal excretion of the latter More recently, a hormone has been isolated from the adrenal cortex that has a much more powerful effect upon salt metabolism than has desoxycorticosterone. It is called electrocortin or aldosterone; like desoxycorticosterone, it causes the retention of sodium and the excretion of potassium, but, as just mentioned, to a much more pronounced degree. The principal corticoids and their actions are given in Table 10-1.


The adrenal cortex also contains the female sex hormones estrone and progestin, as well as one with male hormone activity known as adrenosterone.

The adrenal cortex is relatively rich in cholesterol and in vitamin C (ascorbic acid) The former is probably the chief substance from which the steroid hormones of the adrenal are synthesized in the body. The concentrations of both these substances are reduced when the adrenal cortex is stimulated, as by the adrenocorticotrophic hormone of the pituitary (p. 435), to liberate its hormones But the role played by vitamin C with respect to the function of the adrenal cortex is unknown. It is not essential for the production of the hor-

mones, for they continue to be formed in animals who have been rendered severely deficient in vitamin C

Other functions of the adrenal cortex. The adrenal cortex performs other functions besides those just described. Several observations suggest that it influences the development of the sex functions. Enlargement of the adrenal cortex, for example, or tumors of cortical tissue sometimes occur, accompanied by sexual abnormalities. In children with such growths, precocious development of the sexual organs and of the secondary sex characters (see p. 444) is seen; these subjects often also show unusual muscular development for their age, and

sometimes an abnormally "beefy" florid face. Such an appearance has prompted the comparison with "a burly brewer's drayman" (Fig. 10.22). Little girls, subjects of this disease, also show precocious sexual development, and in appearance resemble stout women in miniature. Menstruation may occur as early as 3 or 4 years of age. Women who are the victims of cortical enlargement or tumor become obese,



rig 10.25 vinism due to a tumor of the adrenal cortex. On test, at age of 24, syears; on right, at age of 32, about one year after the onset of the disease (Drawn from photographs after Hare, Ross, and Crooke)

and masculine in appearance; hair grows upon the face and chest, and their disposition tends toward "mannishness". This condition is referred to as virilism (Fig. 10 23).

The cortex of the adrenal, as well as the medulla, appears to play a role in states of stress, for when such a state as exposure to extreme cold, hemorrhage, bacterial infection, or severe mechanical injury is suffered by an animal, the output of the adrenal cortex is increased. See also Cushing's disease, page 444

# The Pituitary Body or Gland (Hypophysis Cerebri)

Development and structure. The pituitary gland is connected by a narrow stalk, called the neural stalk or infundibulum, to the base of the brain just behind the optic chiasma. In man it is an ovoid body and measures a little over half an inch in its longest diameter. It is lodged

#### THE ENDOCRINE GLANDS

430

within a small recess in the bone of the base of the skull, known as the sella turcica (L., Turkish saddle). The pituitary consists of an anterior and a posterior portion. These have separate origins and are different in structure and function. The anterior part arises from the embryonic mouth cavity as a hollow pouch (Rathle's pouch), the posterior part is formed by a downgrowth of nervous tissue from the floor of the third ventricle at the base of the developing brain. Rathle's pouch grows





Fig 10 24 Diagram of the pituitary body (hypophysis cerebri)

upward and, meeting the downgrowth from the brain fuses with it The cavity of Ratike's pouch becomes reduced to a narrow cleft by pressure against the posterior part, the anterior wall of the pouch—that is, the portion lying in front of the cleft becomes thickened and is now called the adenohypophysis, pars distalls, or pars anterior of the pituitary (Fig. 10 24). The posterior wall of Ratike's pouch is represented by a thin shell of tissue, closely applied to the anterior aspect of the posterior part of the gland, it is called the pars intermedia. That portion of the pituitary developed from the base of the brain and lying be hind the pars intermedia is known as the neurohypophysis, neural lobe, or the pars nervosa.

The part of the pituitary situated in front—that is, the adenohypophysis—is readily separated from the rest of the gland and is very commonly referred to as the anterior lobe, whereas the portion lying behind the natural line of cleavage offered by the cleft is called the posterior lobe. The latter term, therefore, includes both the pars intermedia and the neurohypophysis. A part of the tissue of Rathke's pouch also extends upward as far as the base of the brain, where it covers a mass of nerve cells in this situation known as the tuber cinereum. This part of the pituitary, which—having the same origin as the anterior lobe—is usually included with it, is called the pars tuberalis. As development proceeds, the connection of the pituitary with the mouth cavity disappears, but that with the base of the brain persists and, as mentioned above, is called the neural stalk or infundibulum.

The nomenclature of the parts of the pituitary is confusing owing to varied designations which different authors from time to time have adopted The different terms are given in the following table, with the names in most common use today given first and the older terms in small brackets.



The adenohypophysis consists of cords of cells separated by relatively wide blood channels (sinuses) The cells are of three main types. (a) those which are stained poorly by ordinary dyes—chromophobe cells; (b) those which stain with acid dyes—acidophil (alpha) cells; and (c) those which stain with basic dyes—basophil (beta) cells (see Fig. 10.25).

The posterior lobe of the pituitary receives numerous fibers from nerve cells situated in the region of the brain near the origin of the infundibulum (see hypothalamus, p. 541). A few nerve fibers have been traced into the anterior lobe, though it is believed that they are not secretory, but entirely vasomotor in function. The structure of the neural lobe is quite different from that of the anterior lobe. It contains nerve fibers, spindle-shaped cells called pituicytes, neuroglial cells, and hyaline bodies. These latter are cells filled with a clear homogeneous material; their function is doubtful, though they are believed by some to furnish the active principle of the posterior lobe secretion.

The functions of the anterior lobe. The functions of the anterior lobe of the pituitary are many. The growth of the skeleton, the development of the sex glands, and the activity of the thyroid, adrenal, and parathyroid glands are dependent upon this part of the pituitary. Within

the last 15 years or so the anterior lobe has been the object of intensive investigation. A number of separate active principles or hormones, each of which exerts its specific effect when injected into animals has been obtained by the extraction of anterior lobe tissue. It has been apily termed the master gland of the endocrine system (see Fig. 10.48, p. 455). Its hormones are protein in nature.




Fig. 10.25 Microscopical section of the adenohypophysis

The pituitary principles which upon injection give relatively pure effects, and are probably distinct and separate hormones, are as follows

- (1) The growth hormone
- (2) The thyrotrophic hormone (TTH)
- (3) The adrenocorticotrophic hormone (ACTH)
- (4) and (5) Two gonadotrophic hormones (FSH and LSH)
- (6) Prolactin
- Certain effects other than those produced by the foregoing more orless pure principles can be produced by relatively crude (unfractioned) extracts of the anterior lobe. These effects are
  - (1) Diabetogenic
    - (2) Glycotropic
  - (3) Ketogenic
  - (4) Parathyrotrophic
  - (5) Pancreotrophic
- Arrest of skeletal growth follows removal of the pituitary (hypophysectomy) from young animals, and the sex glands do not develop,




Fig 10 26 Curves showing the effect upon the growth of rats of daily injections of an extract of the anterior lobe of the pituitary Upper curve, treated animals, lower curve, untreated littler mate sisters (After Evans)




Fig 10 27 Drawing of an hypophysectomized white rat (left) and its normal litter mate

the thyroid and cortex of the adrenal atrophy. The daily transplantation of anterior lobe tissue into the hypophysectomized animals is followed by the resumption of growth, by the regeneration and normal development of the atrophic testes or ovaries, and restoration to normal of the adrenal cortex and thyroid gland. The effects of hypophysec tomy can be corrected also by anterior lobe extracts, and extracts have been prepared which will abolish one or other defect caused by hypophysectomy but leave the others almost or quite uncorrected. That is to say, extracts more or less pure in one or other of the pituitary hor mones as listed above are now available. These extracts are also capable of causing their respective effects in normal animals.




Fig. 10.28 The effect of an extract of the anterior lobe of the pituitary upon the growth of the dachshund. Normal dog above dog of same litter below, injected. (Drawn from a photograph after Van Dyke.)




Fig. 10.29. The thyrotrophic hormone. Section of thyroid from normal guinea pigo in the *left* that of litter mate treated with 8 mg of anterior lobe substance on *right*. Note the greatly increased cellularity of the latter and the infolding of the follicular walls (After Van Dyke)

Young rats treated daily with a purified growth hormone attain a size nearly double that of untreated animals of the same litter (Figs 10 26 and 10 27). The characteristic effect of the growth hormone has also been demonstrated on puppies (Fig 10 28).

The thyrotrophic's hormone (TTH) exerts its effect upon the thy roid gland. An animal receiving an extract containing this hormone shows marked hypertrophy and increased activity of the thyroid gland and in consequence, a rise in the basal metabolic rate (Fig. 10.29)




Fig 10.30 Showing effect of hypophysectomy upon the size of the adrenal Adrenal from an hypophysectomized animal on the right from normal an mal on the left. Note that the effect is exerted chefly upon the cortex (After P Smith)

The adrenocorticotrophic hormone (ACTH) maintains the functional activity of the adrenal cortex. It stimulates the elaboration and output of cortisone and the other corticoids (see Fig. 10.30)

The gonadotrophic hormones are essential for the normal development of the gonads and for the production of the respective sex hormones of the male and female animal. They will be considered in the section on the sex hormones (p. 449). Prolactin, which stimulates the secretion of milk, will also be dealt with in that section.

The diabetogenic and glycotropic effects appear to be due to differ ent principles in anterior lobe extracts. A crude extract given over a period of days produces a permanent diabetes—hyperglycemia gly cosuria increased production of urine (polyuria), thirst and wasting. The glycotropic effect is a temporary hyperglycemia and glycosuria.

• It has been recommended that the suffix tropic derived from the Greek trope (a turning) be changed to trophic from the Greek trophic (nourishment) on the grounds that the latter more correctly expresses the action of these p tu tary princ ples. The end ng tropic is used in such words as geotropic heliotropic glycotrop c etc.

This action is antagonistic to that of insulin. The effect upon carbohydrate metabolism exerted by the anterior lobe of the pituitary can also be demonstrated by the following experiments

(a) Hypophysectomy, by removing the source of the sugar raising principle, results in a fall in the blood sugar (b) Excision of the pancreas alone (removal of the source of insulin, p 370) causes hyperglycemia and the other signs of diabetes, but if the pituitary is removed as well, intile or no rise in blood sugar occurs, there may, indeed, be hypoglycemia when the animal is fasting The transplantation of anterior lobe tissue into such a pancreatectomized hypophysecto-




Fig 10 31 Showing the effect of pituitrin upon the arterial blood pressure B.P., carotid blood pressure white square indicates the injection of 0.5 cc pituitrin (After Herring)

mized animal causes the appearance of the usual effects of pancreatec tomy. A pituitary extract with an effect upon carbohydrate metabolism, but entirely free from growth effects, has not been prepared. It is probable that the diabetogenic effect is simply a manifestation of the growth hormone.


The ketogenic effect is evident when an anterior lobe extract is in jected into a fasting animal, or better, into one on a diet of fat when ketone bones—acetoacetic acid and  $\beta$  hydroxybutyric acid—appear in relatively high concentration in the blood and urine. Associated with the ketogenic effect are a reduction in body fat and increase in the fat of the liver

The relationship between the pituitary and parathyroid glands is rather indefinite but it does appear that the master gland everts an in fluence upon the activity of the parathyroids (parathyroine effect). The effect of a crude anterior lobe extract upon the pancreas is more definite, and following the administration of such an extract, the growth of the insulin producing cells (islands of Langerhams p 323) in the pancreas is stimulated. This is the so-called pancreotrophic effect.

The physiology of the posterior lobe. An extract of the posterior lobe of the pituitary has been used in medicine for a number of years. It is sold under various trade names—e.g., pituitrin, infundin, etc. It produces the following effects when injected subcutaneously:

- (a) A rise in arterial blood pressure occurs as a result of general vasoconstriction (Fig. 10.31). This action differs from that of an extract of the adrenal medulla, for all vessels—cutaneous, muscular, coronary, etc.—are constricted alike, there being no relationship between its effects upon various structures and those caused by sympathetic stimulation.
- (b) Contraction of the smooth muscle of the uterus, intestine, gall bladder, urinary bladder, and ureter, as well as of that in the walls of

Fig. 10.32 Melanophores of frog's skin A, fully expanded, B, partially contracted. C, fully contractedi e., collection of pigment at centers of the cells (Redrawn from Hogben ) Below, enlarged views of a melanophore, expanded on the left, contracted on the right The branching processes are invisible in the latter because they contain no granules



the follicles and ducts of the mammary gland The contraction of the uterine muscle is called its oxytocre action. The potency of a posterior pituitary extract is assayed by testing its oxytocic action upon the excised uterus of a virgin guinea pig. It is for this effect that the extract is most often employed in medicine, being administered in the later stages of labor to stimulate uterine contractions. It is also used to stimulate the muscle of the intestinal wall and thus to relieve trouble-some abdominal distention, especially after operation upon the abdomen. The effect of pituitary extract upon the mammary gland is called its galactogogue action (GK. gala [galact—]. milk + agōgos, drawing (orth), for it causes the expression of milk from the follicles and along the ducts to the surface of the nipple. This is not a true secretory effect—that is, a production of milk by the gland cells—but merely an expression of preformed milk from the alveoli and ducts by the con-

traction of smooth muscle fibers surrounding these structures. The commencement of the milk flow in cows at milking time, and what farmers speak of as the 'letting down' of the milk, is thought to be due to the galactorogue principle.

(c) Reduced production of urine-antiduretic effect (p 281)

(d) Hyperglycemia glucosuria and reduction of liver glycogen

(e) Expansion of the melanophores in the skin of various kinds of cold blooded vertebrates. The skin of the frog is abundantly supplied with melanophores. These are irregularly shaped cells possessing several branching processes and granules of a dark pigment (Fig. 10.32). The processes of adjacent cells interlace with one another to form a rich network. The pigment at one time may be concentrated near the center of the body of the cell, the next moment the granules may migrate and fill the processes. In the former event the skin is pale,




Fig 10.33 Upper drawing A the color of a normal frog B the same frog after the injection of an extract of the posterior lobe of the patientry (Re drawn from Hogben) Below the effect of hypophysictomy upon the color of tupoles A hypophysectomized B nor

in the latter it is dark. The movements of the pigment granules, and the consequent darkening or lightening of the skin depend upon the illumination of the en vironment thus by changing its color to match its surroundings the animal is rendered less con spicuous to its enemies. An injec tion of pituitrin causes the gran ules to move into the branching processes the animal then be coming a deep brown or green (Fig 10 33) Hypophysectomy, on the contrary, by removing the supply of posterior lobe princi ple results in permanent pallor of the skin Blinding the animal as by sealing its eyes with wax, also abolishes the skin responses so long as the eyes are closed The skin color remains unaltered in light or shade. The effects are evidently initiated by light re flexes and are brought about in the following way The pituitary secretes the melanophore-expanding principle continuously, so long as no light, or light of low intensity, falls upon the retina, the skin is then dark. Retinal illumination, on the other hand, sets up nerve impulses which suppress temporarily the pituitary secretion, the skin, therefore, becomes pale Many species of fish and certain lizards and toads possess a similar mechanism for altering the depth of color of their skins in conformity




Fig 10.34 An example of giantism Father on right and boy on left are of aver age height.

with the illumination of their immediate environment. The pigment granules in some forms are red (erythrophores), in others yellow (xanthophores)

The several effects of posterior lobe extract are due, not to one, but to at least three and possibly four, distinct principles. A short time ago two substances in the form of white powders were separated from the relatively crude posterior lobe extract (pituitin). These are known as pitressin and pitocin (or oxylocin). The former is vasoconstrictor (pressor), intestine stimulating, and antiduretic in action, the latter contains the uterine stimulating (oxylocic) principle. Both raise the sugar of the blood. The melanophore expanding effect is caused only by pitressin, though it appears that this effect is due to a separate prin

ciple which is merely associated as a contaminent with pitressin; there is convincing evidence that it is a separate hormone elaborated by the pars intermedia, and has been named intermedia. Indeed pitressin and pitocin are far from being pure substances, and it is possible that the antidiuretic effect of the former is also due to a distinct hormone.

Disorders of the pituitary in man. Giantism and acromegaly-hyperpituitarism. The great stature of those persons whom one sees in




Fig. 10.35 Acromegaly. (Drawn from a photograph after Lisser and Goldberg.)

circuses or occasionally reads about in the daily papers is caused by overactivity of the anterior lobe of the pituitary. Some of these giants are between 8 and 9 feet tall (Fig. 10.34). Of course, general overgrowth of the skeleton can occur only if the hyperpituitarism commences during the normal period of growth. Hypersecretion of the growth hormone of the pituitary after adolescence does not cause an increase in stature. Nevertheless, the stimulating effect of the hormone upon growth is seen in the bones of the face, hands, and feet. This condition, resulting from hyperpituitarism in adult life, is called acromegaly (Gk. acron, extremity + megas (megal-), large); the appearance of a person suffering from it is characteristic. The nose and lower jaw, especially the latter, are abnormally large and prominent, and the forehead unusually massive (beetling) where it ordinarily forms a slight eminence above the eye sockets. The skin of the face is thick and

coarse. There may be overgrowth of hair upon the chest; in women hair may appear upon the face. There are often signs of overproduction of other hormones of the anterior lobe of the pituitary—e.g., hyperthyroidism (thyrotrophic hormone), signs of overstimulation of

the sex functions (gonadotrophic hormones), or hyperglycemia, glucosuria, and other diabetic manifestations (diabetogenic principle). The disease is gradually progressive (see Fig. 10.35). In most instances a pituitary tumor composed of acidophil cells is found after death.

Dwarfs—hypoputuitarism. Deficiency of the anterior lobe secretion in childhood produces dwarfism. There are two main types of pituitary dwarf. In one type, the diminutive body is of normal proportions, or rather, shows the proportions of the normal young child, the head being large, relatively, to the rest of the body (see Fig. 10 36). These subjects may or may not show failure of sexual development. This variety of dwarf is usually intelligent, and not unattractive in appearance; it is referred to as the Loran type. The midget of the circus belongs to this group.

In the second type of pituitary dwarf, the chief features, other than the stunting itself, are obesity and arrested sexual development. Children affected by this pituitary disorder are often subnormal mentally, usually lethargic or somnolent, and have large appetites, especially for sweets. The fat boy of Pickwick Papers was undoubtedly an example of this disorder. The condition is called



Fig 10 36 Lorain type of pituitary dwarf, age 21 years. Man on the right 5 feet 7 inches. (Drawn from a photograph after Lisser.)

dystrophia adiposogenitalis or Frohlich's type of dwarfism. Diabetes insipidus often accompanies this condition. It is thought to be due to destructive disease involving the neurohypophysis or the base of the brain in the region of the origin of the pituitary stalk (hypothalamic region), together with deficiency of the growth hormone of the anterior lobe (see Fig. 10.37).

Essentially the same disorder occurs in adolescents or adults but,




Fig 10 37 Frolich's type of dwarf ism (juvenile form of dystroph a adsposogenital's (From a photograph after Cushing)

Fig 10 38 Dystrophia ad posogeni tal's in a woman (Drawn from a photograph)

of course, dwarfing is absent. The obesity is often extreme the sex functions are suppressed. When it occurs in the male, the fat has a feminine distribution—that is, over the hips, thighs and chest, the skin is smooth and soft, the hair is scarce or absent from the face and body, but fine and plentiful on the head, such persons tend toward ef feminacy in temperament and disposition (see Figs. 10.38 and 10.39).




Fig 10.39 Aduposity in a female rat (on left) following purtial destruction of the pituitary by the injection into it of chromic acid hypothalamius also probably injured Litter mate normal female rat on the right (Drawn from a photograph after P Smith)

Ptuutary cachexia or Summond's disease. This is a rare but very grave disease of the pituitary more commonly affecting women. It is due to atrophy of the anterior lobe occurring in early or middle adult life. The manifestations of the condition are those of a premature se nility. For example, a young woman giving every appearance of healthy youth may within 5 years or so show all those features which one ordinarily associates with old age. The skin becomes dry, sallow, and wrinkled, the hair gray and sparse the body emaciated, and the sexual organs atrophic, menstruation ceases. The bones are frail and appear to diminish gradually in size. Hypoglycemia (due to deficiency of glycotropic principle) often occurs. Mental changes ultimately su pervene. In a large proportion of cases, the changes in the pituitary which lead to this grave condition follow pregnancy.



Fig. 10 40 Cushing's disease.

Cushing's disease or syndrome. Hypersecretion of the adrencocriteotrophic hormone of the anterior lobe of the pituitary causes a state first described by Dr. Harvey Cushing of Boston. It is marked by obesity involving the face and trunk, but not, as a rule, the limbs The face is round and the features almost obscured by fat, the so-called "moon face." The

back of the neck is heaped up with fat, an appearance referred to as "buffalo neck" There may be hypertension. This disease may not only result indirectly from a pituitary disorder affecting the adrenal cortex, but may be caused by hypertrophy or a tumor of the adrenal cortex occurring as the primary disease.

Diabetes insipidus. This is the name given to a disorder characterized by excessive thirst and the passage of large quantities of very ditute urine. The urine has a specific gravity of around 1001 to 1002, and may have a daily volume of over 20 liters. The disease is due to a lesson involving any part of the hypothalamico-pituitary mechanism—namely, the hypothalamus, the posterior lobe of the pituitary, or the nerve fibers descending from the hypothalamus to the posterior lobe. Failure of the secretion of the antiduretic principle of the neural lobe is quite evidently the cause of the excessive production of urine (polyuria) Administration of posterior lobe extract (pituitrin) temporarily relieves the condition For a period of several hours after a single dose has been given by hypodermic injection or sprayed into the nostrils, the urine is reduced in volume and much less didute (see also Chap. 7).

## The Sex Glands or Gonads

The sex glands or gonads of the female are called the ovaries, those of the male, the testes (see Chap. 14) The gonads are also referred to as the primary organs of sex. The uterus, Fallopian tubes, and vagina of the female and the epididymis, seminal vesicles, prostate, and penis of the male are called the accessory organs of sex. Those changes which make their appearance at the time of sexual maturity (puberty), and which in many species distinguish at a glance the male from the female, are known as the secondary sex characters. In the human subject these

include the growth of hair upon the pubis and in the axillae of both sexes, the bass voice and beard of men, and the development of the breats of women. The large comb, wattles, spurs, and tail feathers of the domestic cock (rooster) and the antlers of the stag are other examples of secondary sex characters.

Excision of the sex glands, whether of the male or female, is called castration. Excision of the female gonads is also termed ovariectomy, or spaying. The effects of castration upon the male have been known from ancient times. When this operation is performed upon the male chicken (cockerel), the secondary sex characters do not appear; sex instinct is suppressed. The body weight of the castrated birds, which are known as capons, is greater, however, than that of uncastrated males of the same age. Castration of young stags prevents the development of the antiers. In boys who have been castrated before the age of puberty (a practice even of modern times, in certain countries), the larynx does not develop, and the voice remains high pitched, the accessory organs of sex do not mature, and the secondary sex characters fail to appear. The operation exerts an effect upon fat deposition, the subjects tending toward obesity. The proportions of the skeleton are also altered, the bones of the lower limbs tending to be abnormally long in relation to the trunk. That the gonads are endocrine organs is shown by the fact that the effects of castration in animals can be prevented by grafting tissue of the excessed gland into some other part of the body. Any influence which the transplanted tissue can then exert upon other tissues must, obviously, be solely through the blood stream.

upon other tissues must, obviously, be solely through the blood stream.

The sexual phenomena of the mature female. The organs of sex in both the male and the female animal mature at a definite age which varies in different species. At this time, which is called puberty, the reproductive functions commence. Sex desire is aroused and mating occurs. The reproductive period extends throughout the greater part of the animal's life; it ends with atrophic changes in the sex organs and the gradual suppression of sexual activity. In the human species, puberty occurs at from 13 to 15 years of age, usually being a little earlier in girls than in boys The secondary sex characters already described now make their appearance.

now make their appearance.

In the majority of animal species, the female will receive the male only at a certain period or periods of the year. These so-called mating seasons are characterized by certain sexual phenomena which are generally referred to by animal breeders as "heat" and by physiologists as estrus (also spelled oestrus). Estrus, and the sex phases immediately preceding and following it, are together referred to as the estrus cycle.

The phase preceding estrus is called proestrus that following it postes trus. The changes during proestrus consist of swelling and increased vascularity of the vulva and vagina. The uterus becomes enlarged and its glands hypertrophy. In some animals, at this time, bleeding occurs from the uterus and appears externally. The changes of proestrus are preparatory in nature, the female organs being brought into a condition suitable for the reception of the male and the fertilization of the ovum. The Graafian follicles in the ovary (p. 648) are undergoing maturation. Estrus itself is the period during which ovulation occurs and the female will mate. The changes occurring during postestrus are anticipatory to the implantation of the fertilized ovum in the uterus.




Fig 10 41 Diagram of the estrus cycle (Modified after Corner)

The uterine mucosa hypertrophies and its glands show increased secretory activity. During postestrus the corpus luteum is developing. The uterine changes in postestrus resemble those taking place during pregnancy which indeed, are an extension or continuation of the former.


In a number of animals the postestrus preparation of the uterus for the implantation of the fertilized ovum—te, for pregnancy—is a very prominent feature of the estrus cycle. Postestrus in such species is therefore called pseudopregnancy. If fertilization of the ovum by a sperm cell occurs, the uterine changes of postestrus merge into those of pregnancy. If fertilization does not result, the uterus returns to its resting state and all sexual activity subsides until the commencement (proestrus) of the next estrus cycle.

resting state and all sexual activity subsides until the commencement (proestrus) of the next estrus cycle

Some animals, such as the dog are called monestrous since a single estrus cycle occurs in each mating season. The term anestrus is applied to the quiescent periods intervening between the mating seasons. In other species such as the cow, mouse, and rat, two or more estrus cycles occur during each mating season, such animals are termed polyestrous. The interval clapsing between any two estrus cycles is then

termed diestrus and the period between mating seasons, as in the case of monestrous animals is called anestrus

The phases of the estrus cycle in a monestrous animal are shown in the following scheme and in Figure 10 41.



The menstrual cycle The rhythmical series of changes in the sex organs, which occurs about every 28 days throughout the reproductive life of women, is analogous to the estrus cycle of lower animals. It is called the menstrual cycle (L mens month). In one phase of the men strual cycle uterine bleeding occurs. This phase is referred to as men struation or the menstrual period, it has a duration of from 3 to 5 days. Ovulation occurs about midway between the menstrual periods—that is, somewhere between the thirteenth and seventeenth days following the commencement of the bleeding (see Fig. 10.42).

Though, as just mentioned the menstrual cycle corresponds to the estrus cycle, the two differ in this respect bleeding which is a prominent feature of the former, is absent or inconspicuous in the latter It is not possible to say to what phase of the estrus cycle (whether to proestrus or postestrus) menstruation itself corresponds Since bleeding in certain species occurs in proestrus, it has been assumed by some that such bleeding is analogous to menstruation others have thought that menstruation corresponds to the end of postestrus—that is, to the breakdown of the uterine mucosa built up during postestrus. There are objections to both these views

Conspicuous histological changes occur in the uterine mucous membrane (endometrium) during the menstrual cycle Though these changes have been very minutely described and divided into stages, it will suffice to give a brief outline here. The endometrium becomes thicker and more vascular, the small arteries become tortuous and coiled in a corkscrewlike fashion, the surface epithelium hypertrophies. The glands of the mucous membrane elongate and show secretory activity. The hypertrophy of the epithelium and the vascular changes are due to the action of estradiol (follicular phase of the cycle), the

AAR

somewhat later glandular changes are caused by progestin (luteal phase) Should fertilization and implantation of the ovum (p 669) occur, the stimulating effects of the ovarian hormones become more pronounced, and merge with those of the pregnant state II, on the other




Fig 10 42 Diagram of the menstrual cycle Note the thickening of the mucosa the increasing tortuosity of the vessels and glands as progestin production increases Breakdown of the mucosa with menstrual bleeding occurs as the corpus luteum degenerates or retrogresses

hand, the ovum is not impregnated by a male sex cell (spermatozoon, Chap 14), it degenerates as does the corpus luteum. The hypertrophed endometrum cannot be sustained in the absence of the latter, its epithelium is shed, and bleeding (menstruation) then occurs. Menstruation, therefore, is the sign that a fertilized ovum has not become implanted in the uterine mucous membrane, in other words, the pregnancy has not commenced. After from 3 to 5 days of menstrual bleed-

ing, repair processes restore the surface epithelium, the vascular and glandular activities of the endometrium subside, and the uterus returns to the resting condition, the menstrual cycle has been completed

The first menstrual cycle commences at puberty Toward the end of the reproductive period, the cycles become irregular and finally cease, usually between 45 to 50 years of age, the cessation of the menstrual cycles at this time is called the menopause It is accom-




Fig 10 43 Section of a portion of an ovary showing Graafian follicles in various stages of development

panied by atrophic changes in the ovaries and the accessory organs of reproduction Psychic phenomena, depression or irritability are common during this period Instability of the vasomotor system, causing sudden flushing of the face, and a sensation of warmth—hot flushes—are of frequent occurrence

Ovarian hormones. The phenomena of the estrus and menstrual cycles, described in the preceding section, are dependent upon hormones produced in the ovary The Graafian follicles of the ovary increase in size during proestrus (or, in the human subject, in the phase of the menstrual cycle immediately following menstruation) and become filled with fluid, the ova enlarge and approach maturity A hormone called estradiod or the follicular hormone s is produced at this

6 Other substances closely similar chemically and possessing the physiological activity of the follicular hormone have been obtained from the ovaries as well as from female urine and other body fluids and tissues Estrone (also known as thee

time by the cells lining the follicles. When a follicle reaches maturity, it ruptures and discharges the ovum. This event, called ovulation, occurs in animals during estrus, and in women midway between the menstrual periods—that is, about 18 days after menstruation. The cells lining the ruptured follicle then multiply and form a mass or body called the corpus luteum (yellow body), for the cells composing it contain a yellow, fatty material known as lutein. The corpus luteum replaces the small blood clot which filled the follicle immediately after its rupture and the discharge of the ovum. The corpus luteum is virtually a small temporary ductless gland (see Figs 10.42 and 10.43), for it produces an internal secretion—the hormone of the corpus luteum. This is more commonly referred to as progestin or progesterone. The latter is a chemical term and is used more especially when referring to the hormone in pure crystalline form.

mone in pure crystatine form.

Estradiol is responsible for the changes in the accessory organs of reproduction during proestrus and estrus, for the psychic phenomena of estrus, and for the development of the secondary sex characters (p. 444) of the mature female. It is, therefore, sometimes called the female sex hormone, and corresponds to the male hormone elaborated by the testes (p. 451). As already mentioned, removal of the ovaries of a young female animal prevents it from becoming sexually mature. The accessory organs fail to develop, estrus does not occur, the secondary sex characters do not appear, and the sex instinct is never manifested. Injections of the female sex hormone into such an animal correct all these effects of ovariectomy. On the other hand, young immature animals treated with estradiol show precocious sexual development. The role played by estradiol in pregnancy and parturition is described in Chapter 14.

The hormone of the corpus luteum—progestin or progesterone prepares the uterus for the reception of the fertilized ovum. It is responsible for the uterine changes characteristic of postestrus or pseudopregnancy, and for the development of the placenta—the organ which enables the embryo to receive nourishment from the mother during pregnancy (p. 670). For example, if a pregnant rabbit is castrated

lin), which differs from estradiol in chemical structure to a minor extent, was formerely thought to be the naturally occurring hormone produced by the maturing Graafian follicle Estradiol has been synthesized and has also been isolated in crystalline form from the urine of pregnant women

Estrin is an earlier term given to the follicular hormone before its chemical structure was fully known. It is still in use as a physiological term but carrier no precise chemical meaning <u>Estrogen</u> is another name with a purely physiological commutation, it is applied to any agent, whether of biological or other origin, which is capable of inducing estrus and often other effects of the follicular hormone

(ovariectomized) early in pregnancy or if the corpora lutea are excised, the embryo dies and is expelled from the uterus (abortion). The hormone of the corpus luteum can be prepared most conveniently from hog's ovaries. This preparation, when injected into ovariectomized animals after they have become pregnant, permits pregnancy to continue to full term. Normally, if fertilization of the ovum occurs, the corpus luteum continues to increase in size until the later months




Fig 10 44 Cells of seminal vesicles of rat. Left, of normal animal, middle, 3 weeks after castration; nght, of castrated animal after daily injections of male hormone.

of pregnancy, its hormone exerting a constant influence upon the growth and functional integrity of the placenta. On the other hand, if the ovum remains unfertilized, the life of the corpus luteum is short. After exerting its hormonal influence for a time and causing the uterine changes characteristic of pseudopregnancy, it undergoes degeneration. In the human subject, changes in the uterine mucosa occur after ovulation analogous to those of pseudopregnancy but, again, failure of the ovum to become fertilized is followed by degeneration of the corpus luteum. The uterine mucosa then reverts to its resting state; it is at this time that menstruation occurs (see Figs. 10 41 and 10.42).

Progestin also brings the mammary glands to full maturity during pregnancy, induces multiplication of the uterine muscle fibers, and inhibits contractions of the uterus (see Chap. 14).

The male hormone—testosterone. In 1929 McGec obtained the male hormone in crude form by the extraction of bull's testes with acctone.

457

This preparation was shown to cause the growth of the comb and wat tles of the castrated cockere! (capon) and to bring about the normal development of the accessory organs (penis, seminal vesicles, and prostate) of castrated male rats (Figs. 10.44 and 10.45). In immature rats treated with the extract, the accessory sex organs develop and sexual



Fig 10 45 Castrated cockerel (capon) on *left* note the undeveloped comb and wattles Capon after receiving daily in jections of male hormone on *right* 

maturity is reached prematurely The extract also stimulates the comb growth in hens and inhibits ovulation—masculinizing effect


More recently, the normone has been obtained in crystalline form from testicular tussue This purified material is called testos terone, and has the empirical formula C<sub>13</sub>H<sub>28</sub>O<sub>2</sub>. The hormone has also been obtained in crystalline form from male urine Chemically, this latter material differs slightly from testosterone

and has been designated androsterone its activity is also much less (one seventh to one tenth) than that of testosterone. The male hormone does not appear in the urine of boys until after the tenth year but oddly enough it is present in the urine of normal women. The male hormone belongs with the ovarian hormones, to the class of substances known as steroids (p 372). Both testosterone and androster one have been synthesized in the laboratory from cholesterol. This suggests the possibility that the male hormone is formed in the body from this substance, which is a constituent of nearly all animal tissues.

Cells in the interstitual tissue of the testes—the so-called cells of Leydig—are generally believed to manufacture the male hormone That these cells and not the spermatogenic cells (p. 656) are respon sible seems evident from the fact that castration effects do not follow irradiation of the testes with x rays which destroys the spermatogenic cells but leaves the cells of Leydig unaffected

The relation of the anterior lobe of the pituitary to the sexual functions—gonadotrophic hormones. The anterior lobe of either the male or female pituitary contains two hormones which act upon the gonads In the female one of these gonadotrophic hormones stimulates the growth and maturation of the Graafian follicles in the ovary Thus in directly this hormone causes the production of the follicular hormone It is called the follicle stimulating hormone (FSH) of the anterior

pituitary The other gonadotrophic hormone of the pituitary acts upon the second phase of the ovarian cycle—i e, it stimulates the growth of the corpus luteum This gonadotrophic hormone is therefore known as the luteinizing hormone (LH), or as the lutein stimulating hor-




mone (LSH) <sup>7</sup> It therefore, through its stimulating effect upon progestin formation indirectly causes the changes in the uterine mucosa characteristic of pseudopregnancy and of the corresponding phase of the menstrual cycle, or, if pregnancy ensues, of the development of the placenta (cf p 670) Both gonadotrophic hormones, whether derived from female or male pituitaries, produce analogous effects upon the testes. The follicle stimulating hormone acts upon the elements of the testes corresponding to the lining cells of the Graafian follicles—

<sup>&</sup>lt;sup>7</sup> The growth of the corpus luteum after it has been formed and its maintenance in a functioning state depends not upon the lutein stimulating hormone but upon another pituitary hormone. This has been given the name of the luteotrophic hor mone. It now appears, however that it is simply prolactin (p. 454)

namely, the spermatogenic cells. The luteinizing hormone stimulates the interstitual cells of Leydig (Chap 65) On this account it is also known as the interstitial cell stimulating hormone, abbreviated ICSH

Removal of the pituitary is followed by atrophy of the gonads and secondarily, of the accessory sex organs (Figs 10 45 10 46 and





ysectomy on the testis of hypoph ysectomized rat on left of normal rat on right (Drawn from photographs after Van Dyke )

10 47) Transplantation of tissue of the anterior lobe or injections of anterior lobe extracts prevent these otherwise inevitable results of hypophysectomy The anterior lobe of the pituitary is therefore ultimately responsible for the sex ual development of the male or female animal The phenomena of puberty-namely, the development and maturation of the gon ads and, through the intermediary of the hormones liberated by the latter the growth of the accessory organs of sex and the develop

ment of the secondary sex characters-are dependent upon the gonado trophic hormones

Prolactin the lactogenic hormone The anterior lobe of the pituitary also liberates a principle which stimulates milk secretion in the mother after the birth of the young This hormone is called the lactogenic principle or prolactin. It is obtained from the pituitary by extraction with an acid solution. If the pituitary is removed late in premancy only a small quantity of milk is secreted after the young are born and the secretion soon dries up. This hormone evidently also exerts an in fluence upon the maternal instinct for hypophysectomized animals do not care for their young like normal mothers Furthermore, injections of prolactin arouse the maternal instinct in young virgin rats they commence to build a nest from straw wool or other soft material which they can gather

Prolactin is also responsible for the growth of the corpus luteum during pregnancy (see footnote p 453)

The liberation of prolactin from the hypophysis and its stimulating action on the secretion of milk coincide so closely with the termination of pregnancy that curiosity must naturally be aroused concerning the mechanism underlying such accurate timing \* Many details of the

. Some secretion of milk into the foll cles of the mammary gland occurs before

highly complex mechanism are obscure, but it appears from the great volume of experimental work carried out within recent years that the immediate or direct cause the flow of milk from the breasts is hormonal rather than neural in nature. There is abundant evidence that prolactin is not concerned alone, but that the galactogogue principle (p. 437), estradiol, and progestin, a hormone of the adrenal cortex,




Fig. 10 48 Diagram showing the influence of the hormones of the hypophysis upon endocrine and other structures; anterior lobe, black; posterior lobe, white.

and even the thyroid hormone also play their parts. However, the liberation of prolactin from the hypophysis is initiated by nervous impulses. The latter set up in the nipple by the act of suckling and conveyed to the pituitary body (through autonomic nerves) constitute probably the most important single nervous factor in inducing the production and liberation of the lactogenic hormone Suckling also causes the passage of the galactogogue principle into the circulation. But that there are other nervous influences affecting milk secretion can be attested to by any dairyman. Contentment of milking cows, and even music in the stable, facilitates the "letting down" of the milk, an effect which is be-

the end of pregnancy, but as a rule, it is not until the uterus has been emptied that milk appears externally. This milk secreted before the birth of the offspring and retained by the gland is yellow in color and creamy in consistency, it is known as colortrum

lieved to be due, at least in part, to the release of the galactogogue principle and the resulting stimulation of the smooth muscle in the walls of the milk follicles and ducts. During pregnancy progestin is in high concentration in the blood and, according to some investigators, this depresses prolactin production but, when toward the end of pregnancy




Fig 10.49 Showing the relation of the thymus to the thoracic viscera and great vessels C common carotid artery, H, heart, L, lung S subclavian artery, T thymus

a rather abrupt reduction in the blood content of this hormone occurs prolactin liberation follows automatically.

The actions of the various hormones of the anterior lobe of the pituitary body are summarized in Figure 10 48

The hormones of the placenta The placenta contains two hormones One of these is estradiol, the other, called by Collip the anterior pitutarylike (APL) principle, acts upon the ovary, stimulating the growth of the corpus luteum It also stimulates, though to a slight extent, the maturation of the Graafian follicles The estradiol found in the placenta is apparently manufactured by this organ, and does not simply represent a store of the hor

mone produced by the ovary This conclusion is justified by the observation that in pregnancy large quantities of the follicular hormone continue to appear in the urine, though the ovaries have been removed APL also appears in the urine of pregnant women. This fact is the basis for a well known test for pregnancy Two German workers, Aschheim and Zondek, found that the injection of a small quantity of urine of a pregnant woman into a young (sexually immature) mouse caused within 100 hour the following effects

- (a) The onset of estrus
- (b) Small hemorrhages into some of the Graafian follicles
- (c) The formation of corpora lutea

a In certain abnormal states of the hypophysis associated with overactivity of the anterior lobe such as acromegaly finit may be secreted in the absence of prepancy or even in males This is due presumably to the production and liberation of prolation. Also the mammary glands of newborn infants of either sex sometimes secrete milk. This was long known as "witch's milk" and as now attributed to pro-

lactin in the infant's circulation derived from the blood of the mother before birth action of prolatin upon the crop glands of doves and pigeons is of especial interest. These glands secrete a creamy material with which the birds feed their young. The crop glands of virgin pigeons enlarge and secrete an abundance of crop milk when injected with a potent preparation of the Jactogenic hormone. The Aschheim-Zondek test for pregnancy is almost infallible; its error is only about 1 percent. The close resemblance between the effects of the urine of pregnancy and those of the gonadotrophic principles of the pituitary, led Aschheim and Zondek to the conclusion that




Fig 10 50 Microscopical section of a follicle of the thymus

during pregnancy the pituitary principles were produced in excess and excreted in the urine. It is now generally agreed, however, that the characteristic effects of the urine of pregnant women are due to a single substance, and that it is not derived from the pituitary, but is the anterior pituitarylike principle (APL) of the placenta

### The Thymus

The thymus lies behind the upper part of the breast bone and extends upward for a short distance into the neck. Below, it lies in relation to the base of the heart and the great vessels (Fig. 10.49). It is of relatively large size in the infant, but commences to shrink at about the age of puberty and, by adult life, it is reduced to little more than a vestigial structure. In the infant, it is soft and pink. When sectioned, it is seen to be made up of a large number of lymph follicles, in each of which an outer zone or cortex and an inner zone or medulla can be distinguished (Fig. 10.50). The cortex is composed of masses of lymphocytes supported by a meshwork of slender branching cells. The

45R medulla also contains lymphocytes but in much smaller members, and

the meshwork is coarser. It also contains peculiar bodies called Hassall's corpuscles, formed of epitheliallike cells arranged concentrically around a mass of granular cells

Many functions have been attributed to the thymus. Though proof is lacking, it is concerned in all probability with the production of lym



Fig 10.51 Diagram to show position of the pineal gland 1, cere bral hemisphere, 2 corpus cal losum 3, pineal, 4 third ventricle of the brain 5, optic chiasma, 6, pituitary 7 pons 8 midbrain

phocytes The structure of the cortical portion of its follicles suggests such a function. Several endocrine functions at one time or another have been attributed to it, but most of the theories which have been advanced are based more upon conjecture than upon fact.

## The Pineal Body

The pineal is a small body lying deep in the brain under the shelter of the posterior extremity of the corpus callosum (p 530 see also Fig 10 51) Though its structure suggests the possibility that it serves some endocrine function there is no reliable evidence that such is the case.

# The Physiology of Nerve and Muscle

The structure of the nerve fiber. A nerve fiber is the elongated extension of a nerve cell whose body is situated in the central nervous system (brain or spinal cord), or in an outlying ganglion (of the sympathetic gangliated cord or of a posterior spinal nerve root). (See also Chap 12). Some nerve fibers transmit messages (e.g., of touch, pain, sound, light, etc.) from the periphery to conscious centers and give rise to the respective sensations. They are, therefore, sensory in function Others carry messages to the muscles, causing them to contract: these are therefore called motor nerve fibers. Certain nerves are composed entirely of sensory fibers; some contain only motor fibers; others. again, contain both types of fiber, and are called mixed nerves (see also Fig. 12.2, p 508). The peripheral nerves-that is, the nerve trunks distributed to the various parts of the body-are constituted of nerve fibers, often in great numbers, bound together into a variable number of bundles. Each bundle is surrounded by a connective tissue sheath called the perineurum. The individual fibers are enveloped in turn and separated, each from its neighbor, by a thin covering of connective tissue known as the endoneurium. Finally, the entire nerve trunk is enclosed and separated from other structures by a thick connective tissue sheath called the epineurium (see Figs. 11.1 and 11.2).

The message transmitted by the nerve fiber is called the nerve impulse; its nature will engage our attention later.

The nerve fiber, then, whether in a peripheral nerve or in the central nervous system itself, is the conducting unit of the nervous system, and may be compared to a single wire in an electric cable or in the central

exchange of a telephone system Nerves which transmit impulses from the periphery to the nerve centers are called afferent those which transmit impulses in the opposite direction are called efferent. Now, of course, all sensory nerve fibers are afferent, but not all afferent fibers are sensory, for some transmit impulses which make no impression upon consciousness—they arouse no sensation Similarly, all motor fibers are efferent but not all efferent fibers are motor, since certain




Fig 11 l Cross section of a portion of a peripheral nerve The small circles represent the nerve fibers

fibers of the autonomic nervous system terminating in glands are se cretory in function

Each fiber entering into the composition of the motor and sensory nerves consists of a delicate filament of protoplasm—the axis cylinder—ensheathed by a layer of fatty material called myelin

Nerve fibers which possess a sheath of this fatlike substance, and they constitute the great majority comprising the tracts in the central nervous system and in the peripheral nerves, are called myelinated or medullated fibers. Fibers which have no myelin sheath such as the postganglionic fibers of the autonomic nervous system, are called amyelinated or nonmedulated fibers.

The layer of myelin enveloping the axis cylinder—the myelin sheath—is enclosed in turn by a very thin, transparent nucleated membrane known as the neurilemma or sheath of Schwam The myelin sheath is interrupted at regular intervals, at these points the neu-




Fig 11 2 Drawing to show the structure of a nerve fiber in longitudinal section and in cross section at two levels

rilemma dips sharply toward, and comes almost into contact with, the axis cylinder These interruptions give the impression of a series of equally spaced constrictions along the nerve fiber, they are called the nodes of Ranwer The axis cylinder contains a number of fine thread like strands known as neurofibrils which run throughout its length (see also p 506) The endoneurum, mentioned on page 459 encloses the nerve fiber lying immediately outside the neurilemma

The myelin sheath is believed by some physiologists to serve as in sulation and thus to prevent the pas sage crosswise of the impulses from fiber to fiber. Others consider that it is chiefly nutritive in function and that its metabolic processes are closely related to those of the axis cylinder. There is no conclusive evidence to support either of these ideas.


The myelinization of nerve fibers is a progressive process and does not occur simultaneously in the various types of nerve It is not complete at birth its completion appears to coin

cide with the establishment of function. The tracts of the spinal cord supplying voluntary nerve fibers to the lower limbs for example are not fully myclinated in infants until about the second year or about the time that the child has learned to walk. The association tracts of the brain are myclinated still later, whereas myclination of the sensory tracts of the cord occurs early (fourth month) in fetal life.

Nuclei are to be seen in the neurilemma at fairly regular intervals, one to each section of nerve between the nodes of Ranvier The sheath of Schwann is absent from the fibers of the central nervous system and from the fibers of the optic nerve Its presence is absolutely essential for the regeneration of an interrupted nerve fiber (p 464), fibers in

the situations just mentioned, therefore, once their continuity has been destroyed as a result of disease or mechanical injury, are never repenerated

Degeneration of nerve. Since the nerve fiber is simply the elongated process of a nerve cell, its nutrition depends upon its remaining a part of the cell When, therefore, a nerve fiber is divided or crushed, de generative changes begin almost at once in the section distal to (1e. on the side away from the body of the cell) the point of injury These changes, as they occur in a peripheral nerve, were first described by



Degeneration of nerve Nerve injured at X

Waller, a physiologist of the last century, and are usually referred to as Wallerian degeneration. The first alteration in the fibers of the nerve appears within 24 hours after the injury. The neurofibrils in the axis cylinder become wavy or tortuous, and then break up into sections Next, the myelin sheath swells and droplets of myelin appear (Fig 11 3) The myelin itself undergoes decomposition into its constituents -fatty acids, etc. Finally, usually within two or three weeks following the injury, the debris resulting from the degenerative process is cleared away, nothing remaining of the nerve fiber but an empty tube-the neurilemma which remains intact. The nerve fiber on the proximal side of the point of division (i.e., on the side toward the nerve cell) also degenerates as far as the first node of Ranvier, and even the body of the cell itself may show degenerative changes (retrograde degen eration), consisting of the disappearance of the Nissl bodies (p. 507) and shrinkage of the nucleus

The degenerative changes just described are readily detected by treating the tissue with a suitable stain and examining thin sections of it beneath the microscope. Most of the knowledge which we possess of the origin and course of the various fiber tracts in the brain and cord (Chap. 12) has been gained from experiments upon animals in which a lesion is made in some or other region and the degenerated fibers traced microscopically through successive levels of the nervous system, or from the examination of nervous tissue of subjects who have died of certain nervous diseases which have caused the degeneration of nerve fibers. The location of the degenerative changes following experimental injury to fibers in one or other part of the brain or cord, or resulting from disease, indicates the direction from which the affected fibers originated. For example, if after section of a fiber tract in the cord degeneration occurs above the point of section, the cell bodies of the sectioned fibers must be situated below. Also, by destroying a group of nerve cell bodies in one or other situation and examining tissue taken from different parts of the nervous system for degenerative changes, the course taken by the fibers of the injured cells can be traced (see also Fig. 12.16, p 525). also Fig. 12.16, p 525).

also Fig. 12.16, p. 525).

Degeneration of the peripheral nerves may result from causes other than mechanical injury Peripheral nerve degeneration is seen in poisoning by certain chemicals (e.g., lead, arsenic, alcohol), in vitamin deficiency (A and B<sub>1</sub>), and in certain nervous diseases (e.g., anterior poliomyolitis which causes the destruction of nerve cells in the spinal cord). It is most important in such conditions, or following mechanical injury to a nerve trunk, to have some means of detecting the presence of degeneration, and of determining the extent to which the degenerative process has progressed Such information is afforded by testing the affected nerve or nerves with an electric current. The current is applied to the skin overlying the motor nerve trunk. When the nerve is completely degenerated, it fails to respond (as shown by the absence of a contraction of the muscle) to either the faradic (interrupted) or to the galvanic current. When the current is applied directly over the muscle, a sluggish response is obtained with the galvanic current, but none with the faradic. These abnormal responses of the nerve and muscle to electrical stimulation constitute the reaction of degeneration. degeneration.

Nerve regeneration. Provided that the two ends of a divided nerve are not too widely separated, the continuity of most nerves ultimately becomes restored—the nerve is said to have undergone regeneration. The neurofibrils of the proximal section of the nerve fiber (i.e., the part extending from the cell body to the point of section) grow distally and, entering the empty neurilemmal tube of the degenerated fiber, gradually extend throughout its entire length (Fig. 11.4). The nuclei of the neurilemma multiply. The neurofibrils later become embedded in a ground substance or matrix of protoplasm. Thus, a new axis cylinder is formed. The rate-of-growth-of-the axis cylinder varies in different types of nerve from 0.25 to 2.5 mm per day. A myelin sheath is formed later, and finally the function of the nerve may be completely restored. The sprouting neurofibrils sometimes cross a gap of several millimeters separating the two segments of the nerve—a truly remarkable phenomenon. The force or influence which directs or attracts them into their proper channels is unknown. A neurilemma is abso-



Fig. 11.4 Regeneration of nerve, three types of ending.

lutely necessary for the regeneration process. The fibers of the central nervous system and of the optic and auditory nerves, which do not possess neurolemmal sheaths, are incapable of regeneration after interruption of their continuity, whether this has been caused by disease or mechanical injury.

The regeneration of nerve fibers makes possible the junction by operation of one nerve with another. In facial paralysis, for example, the power to move the muscles of the face can be restored by suturing the proximal end of a sectioned hypoglossal nerve to the distal end of the degenerated facial nerve. The latter nerve is regenerated by the growth of neurofibruls which grow down the empty tube formed by the neurilenima of the degenerated facial nerve. If the operation has been successful and the neurofibrils have reached the facial muscles, the patient, after a period of re-education, can control the movements of the previously paralyzed side of the face.

Nerve cells (i.e., the cell bodies) never regenerate after injury; when destroyed, they are replaced by the nonspecific neuroglia.

The physiological properties of the nerve fiber. Excitability and conductivity are the outstanding properties of the nerve fiber. Though many living tissues possess these properties, they are developed to the highest degree in nerve. When the nerve fiber is stimulated by means of an electric shock, mechanically (as by a pinch), thermally (as by the application of a heated glass rod), or chemically (as by touching it with a crystal of common salt), a disturbance is set up at the point of stimulation which is called the local excitatory state. If the stimulus is sufficiently strong to raise the latter to a certain critical value, a disturbance spreads at high velocity along the nerve This propagated disturbance or wave of excitation, set up by stimulating the nerve fiber, is called the nerve impulse.

fiber, is called the nerve impulse.

When a constant (galvanic) current of moderate strength, such as that derived from an ordinary dry cell, is used as a stimulus, it is found




Fig 11.5 Illustrating stimulation of a nerve by means of a constant current.

that excitation of a nerve, muscle, or other excitable tissue occurs at the instant that the current is closed, and again when the current is opened. Though, of course, the current continues to flow during the intervening period, it does not excite the tissue, provided that its intensity remains unchanged. The closing or making current (make shock) causes excitation of the nerve or muscle at the cathode—that is, at the point where the current leaves the tissue. The opening or breaking current (break shock) causes stimulation at the anode—that is, where the current enters the tissue (see Fig. 11.5). The stimulus caused by making the current is the stronger of the two, so that, when the current strength is gradually reduced, a point is reached at which a response occurs only with a make shock.

Electrotonus. The passage of a constant current through a nerve causes changes in excitability and conductivity of the nerve fibers. These effects are generally referred to as electrotonus. When the cur rent is "made," both these properties of the nerve are increased in the region of the eathode, whereas they are depressed in the region of the anode. The changes produced at cathode and anode are called, re spectively, catelectrotonus and anelectrotonus. The effects are reversed when the current is 'broken, a momentary enhancement of excita birty and conductivity occurring at the anode and depression of these




Fig. 11.6 Diagram of a nerve muscle preparation illustrating electrotonus acrows indicate the direction of the current—ascending or descending from or to the muscle. Shortening and thickening of the muscle indicate its contraction Hatched areas on nerve near the muscle in the second and third sketches represent blocking of the nerve as described in the text.

properties supervening at the cathode. It has been mentioned above that with a moderately strong current, excitation of the nerve, which may be made evident by using a nerve attached to its muscle, causes excitation with both make and break shocks but, upon gradual reduction in the strength of the current, a point is reached when excitation occurs only at the make If however, the current is very strong, and the electrodes are so placed upon the nerve that the cathode is nearer to the muscle than is the anode—that is, the current is descending in the nerve toward the muscle (see Fig. 11 6)-excitation also occurs only on the "make," because on the "break" the depressed conductivity at the cathode blocks the impulse initiated at the anode. If, on the other hand, the electrodes are reversed—that is, the current is ascend ing in the nerve from the muscle-excitation occurs only on the "break," for the impulse arising at the cathode is blocked by the de pressed conduction at the anode occurring on the "make" of the current (see Table 11-1).

Table 11-1

Strength of current	Descending current		Ascending current	
	Make	Break	Make	Break
Very weak	С	0	С	0
Moderate	C	С	l c	l c
Very strong	c	0	0	С

The qualities of a stimulus The prime factors in excitation of nervous or muscular tissues are intensity, rate of change, and duration of the stimulus

- I Intensity or strength of the stimulus. An electric shock, which is the most convenient and commonest type of stimulus used in the laboratory, must be of a certain minimal voltage. The electric stimulus whose voltage is just sufficient to excite is said to be of threshold or liminal strength. A stimulus weaker than this is called subthreshold or subliminal.
- 2 The rate of change in the intensity of a stimulus. An electric stimulus which is just capable of exciting the nerve when its intensity increases rapidly, may be quite ineffective if it reaches its maximum value more gradually. In order to be effective at the slower rate of change, the strength of the current must be increased. In other words, the threshold of excitation rises as the rate of increase in the current strength diminishes. This relationship between the threshold of excitation of the nerve and the rate of change in intensity of the stimulus is referred to as accommodation. Rate of change is a factor of just as great importance in other types of stimulation. We are all familiar with the fact that changes in the environment, whether of temperature sound or movement, are more likely to be effective stimuli if they occurs usuddenly
  - 3 <u>Duration—chronaxie</u> Since, as mentioned above, a constant current excites only at the make and break, though the current continues to flow in the intervening period it had been thought that the duration of a current played no part in excitation But it is known now that the current must flow for a certain minimal time, measured in thousandths of seconds, in order to excite That is to say, if the time elapsing be tween the make and break of the current is too short, the tissue is not excited no matter how strong the current may be Thus, alternating currents of very high frequency but having a strength of several thou sand volts, do not stimulate when passed through the body

The length of time during which a current must flow in order to ex cite a nerve or muscle varies with the strength of the current the time being shorter with a strong than with a weak current. The excitability of a given nerve or other tissue may be measured by determining the minimal intensity of a current which if allowed to flow for an indefinitely long period will excite—that is the time factor is not considered. On the other hand the excitability can be measured by using a current of a standard strength and determining the minimal time during the




Fig. 11.7 Strength-durat on curve. The chronaxie is the time interval marked by the intersect on of the vert cal interrupted line with the upper horizontal line the latter representing a strength of current twice the rhoosase. The chronaxie value in this instance as indicated by the arrow is 002 second

ing which it must flow in order to excite The current of a strength just sufficient to excite but of indefinitely long duration is called the heteobase A current of double this strength is then employed as the standard and the minimal duration for excitation measured. This time factor is called the chronaxie of the tissue in question. In other words chronaxie is a measure expressing the shortest duration for excitation of a current having a strength twice the rheobase (Fig. 11.7.)

The chronaxies of the different types of excitable tissue and of nerves and muscles of the different species as well as of the same species show very wide variation—namely between 0 1 and 100 millisec onds (thousands of seconds) The chronaxie of the muscles of the hu man arm for example is around 0 1 millisecond whereas that of the smooth muscle of the frogs stomach is about 100 milliseconds. The chronaxie of the human sciatic nerve is about 100 milliseconds (see Table 11 2 p 469) Rapidly acting tissues such as voluntary muscle have shorter chronaxies than the more slowly acting such as smooth muscle. The tissues of warm blooded animals have in general shorte chronaxies than those of cold blooded species. The chronaxies of the nerve fibers which conduct at high velocity are shorter than those of fibers of smaller diameter and slow conduction. A nerve and the muscle which it supplies have chronaxies of nearly the same value and certain

Table 11-2

Species	Structure	Chronaxie (milliseconds)
Man	Flexors of arm	0 08-0 16
	Extensors of arm	0 16-0 32
	Retina	12
	Vestibular nerve	14-22
	Sciatic nerve	03
Dog	Muscle of	
_	ventricle of heart Auriculoventric-	20
	ular bundle	60
Frog	Stomach Gastrocnemius	100
	muscle	03
	Ventricular	
	muscle	90
Crab	Muscle of claw	12

drugs-such as curare-which paralyze voluntary muscle are be lieved by some to cause their effect by increasing the disparity be tween the chronaxie values of the muscle and its nerve

Electrical changes in nerve. The nerve impulse is accompanied by a change in electrical potential Any part of a nerve in the excited state is electrically negative, relatively, to all other parts of the nerve. The excited part is as the negative pole (copper) of a battery, any mactive part corresponds to the positive pole (zinc) If, therefore, a pair of

electrodes are in contact, one with an excited part of the nerve, the other with a resting part, and the Two electrodes connected through a sensitive electric recording ap paratus, a current flows and causes a movement of the indicator of the instrument. The direction taken by the current is from the unexcited part of the nerve (+) to the excited part (-), through the instrument, and from excited to unexcited part through the nerve Thus the electric circuit is completed (see B 1, galvanometer 2, stimulus 3, in




Fig 118 Diagram illustrating ac tion current A and current of injury Fig 118A) A similar electrical jured part of tissue (see text)

phenomenon occurs in muscle and other excitable tissue—any active region of a tissue being relatively negative to an inactive region.

The current thus set up in excited tissue is called the action current. Injured nerve or other tissue is also relatively negative to healthy tissue, the current set up when the injured and uninjured sections of the tissue are connected by a conductor is called the current of injury or demarcation current (see Fig. 11.8 B).

As we have seen, the excited state—the nerve impulse—travels rapidly over the nerve fiber. By placing electrodes upon a length of




Fig. 11.9 Showing the production and recording of a diphasic action current (see text) Stippled area indicates wave of contraction

nerve, and connecting them with a sensitive electric apparatus, such as the string galvanometer, or with an amplifier and eathode-ray oscillograph, it is possible to detect the passage of nerve impulses, whether these are set up by artificial stimulation or occur naturally. By means of special photographic apparatus, a permanent record of the action currents can be obtained (see Fig. 11.9), and their velocity and magnitude determined. For example, if the electrodes are placed upon two uninjured parts of a nerve and one end of the nerve stimulated, the impulse when it reaches the first electrode causes this to be relatively negative to the second electrode. The indicator of the electric recording apparatus swings away from the zero position. This causes a stroke to be inscribed above the base line in the photographic record (Fig. 11.9 A). When the impulse passes from beneath the first electrode, but has not yet reached the second (i.e., the nerve beneath each electrode is inactive and so of equal electrical potential), the indicator of the in-

strument returns to zero and a downstroke is inscribed in the record (see Fig. 11.9 B). When the impulse arrives at the second electrode this becomes relatively negative to the first, a current flows through the instrument in the opposite direction a curve below the base line appearing in the record (Fig. 11.9 C). Thus a diphasic curve is produced (Fig. 11.9 D).

If, instead of placing both electrodes upon uninjured tissue, the nerve beneath the second electrode is crushed as is the usual practice in recording the action currents the current resulting from the dif

ference in potential between the in jured and uninjured portions (cur rent of injury) undergoes a sharp reduction in strength when the im pulse reaches the first electrode that is the action current opposes the current injury. This variation in strength (negative variation) of the injury current appears as a single wave in the record. A second wave directed oppositely to the first does not appear, since the implaced upon the crushed section.



Fig 11 10 Action potential wave recorded by means of the cathode ray oscillograph (After Erlanger and Gasser)

first does not appear, since the impulse cannot reach the electrode placed upon the crushed section of nerve Such a record is called monophasic (see Fig. 11.10)

The most convenient and reliable means which we possess for in vestigating the properties of the nerve impulse is provided by the analysis of the electrical changes recorded from the nerve. The strength of the action current as indicated by the amplitude of the recorded curves, their frequency and velocity can be readily determined, such values are taken as corresponding to those of the nerve impulses themselves.

# The Nature of the Nerve Impulse

The "all or none" or "nothing" principle (or law) The 'all-or none principle as it applies to the contraction of heart muscle, has been already defined (p 159) This principle applies also to the nerve impulse and to the contraction of the individual fibers of skeletal muscle For example, a stimulus just strong enough to excite the nerve fiber causes the transmission of an impulse which as gauged by the action current, is of the same magnitude and has the same velocity as one set up by a stimulus of much greater strength. In other words, a stimulus

which is capable of causing a response at all causes the maximum response. Nor can the impulses in the nerve fiber, as can the contractions of a muscle fiber, be summed—that is, the impulses as measured by the electrical response, cannot be increased in magnitude by raising the rate of stimulation. Each impulse remains separate.

The nerve impulse may therefore be compared to a spark traveling in a fuse of gunpowder; for, of course, it is immaterial, in so far as the size of the spark and the rate of its transmission are concerned, whether the fuse has been ignited by a small or by a large flame (which is comparable to the stimulus applied to the nerve). For the sake of illustration other comparisons may be drawn between the nerve fiber and a fuse of gunpowder. The nerve impulse and the spark are self-propagated, both derive the energy for their transmission from the material through which they pass They, therefore, do not undergo any diminution with distance, differing in this respect from a sound wave, a wave in water, or from an electric current, each of which is reduced in magnitude as the distance from its point of origin increases.

If a section of the nerve fiber is treated with a narcotic—that is, ether or alcohol—conduction in the narcotized region is depressed; the impulse (as indicated by the action current) is reduced in magnitude by passing through this region. But the impulse, upon entering the unnarcotized section of nerve, regains its original value—further proof for the statement that the conduction of the impulse is dependent upon energy derived from the nerve fiber itself A wave in air or water, or an electric current, once having undergone a reduction in force as a result of some resistance in its path, is not restored to its original value after having emerged from the medium offering the resistance. The spark in gunpowder behaves, however, like the nerve impulse. If a section of the fuse is dampened, the spark is less intense and travels more slowly through this section, but at once flares up again upon reaching the dry section of powder beyond.

It is therefore evident that the nerve impulse, though inseparable from an action potential, is not merely an electric current transmitted along the nerve as along a wire.

Lillue's iron wire model of the nerve fiber. An electrochemical phenomenon most closely comparable to that of the nerve impulse is exhibited by an iron wire immersed in a solution of acid of appropriate concentration. A nerve model so constructed was devised by Dr. Ralph Lillie of Chicago, and is generally known as Lillue's iron wire model. If an iron wire is immersed in strong nitric acid, it becomes

coated with a layer of iron oxide. If the wire is then placed in a glass tube filled with a solution of acid suitably diluted, and "stimulated" by scratching its surface at one point so as to break the film of oxide, a disturbance consisting of effervescence and the formation of a dark compound is set up and travels rapidly along the wire; the film of oxide gradually re-forms in the wake of the propagated electrochemical change. The wire can also be "stimulated" electrically. The model shows several of the phenomena associated with the passage of the nerve impulse. An action potential can be recorded from the wire as the propagated disturbance passes over it. Electrotonic effects (p. 465) and a refractory period (see below) can also be demonstrated. The latter is that period following the "impulse" during which the oxide coating is being re-formed and the wire cannot be "stimulated."

The membrane theory of nerve excitation and conduction. The electrochemical change in the wire following its "stimulation" is quite evidently a surface phenomenon. The nerve impulse is thought to be of




Fig. 11.11 Illustrating the nerve membrane theory. Straight arrow indicates the direction taken by the impulse, curved arrow, the potential difference The part of the nerve marked by wavy lines is depolarized, and is negative to the polarized part The refractory.

periods in which the nerve is being repolarized are represented by hatching; following this again repolarization has occurred, and the ability of the nerve to conduct an impulse has been restored

the same nature, occurring at the surface of a membrane or film enveloping the nerve fiber. This conception has been embodied in what is known as the depolarization or membrane theory of the origin and spread of the nerve impulse. According to this theory, the nerve fiber in the resting state is polarized—that is, the ions in the fluids in contact with the two sides of the film or membrane are arranged with the cations on the outside and the anions on the inside (Fig. 11.11). Thus, positive and negative electric charges are balanced on either side of the membrane which remains impermeable to them so long as the nerve is at rest. Upon stimulation of the nerve, some chemical change is brought about which renders the membrane permeable to the ions; it

becomes depolarized at the point of stimulation. An electrical potential is set up between the adjacent polarized and nonpolarized regions of the nerve, which acting like the original stimulus causes the spread of depolarization down the nerve, thus the impulse is propagated automatically. Excitation of the nerve is not restored until the depolarized region is repolarized. The period during which the nerve is depolarized corresponds to the absolute refractory period.

The absolute and relative refractory periods of nerve. The nerve fiber after the passage of an impulse is for a short time unable to conduct. This is due if we accept the membrane theory to its being in the depolarized state. This brief interval following the passage of the im




Fig 11 12 Curve of the recovery in the sci atte nerve of the frog Two stimuli were appled to the nerve the second at mulus being separated from the first by various time inter vals and of just sufficient strength to excite Intensity of stimulus is plotted along the ord nate time along the

abscissa. The interrupted horizontal line indicates the threshold strength of urent required to excite the resting nerve. During the absolute refractory period (about 0.003 second in this instance) a stranglas however strong will not excite. The excitability returns gradually during the next 0.02 second (relative refractory period). (After Adrian)

pulse and during which the fiber will not conduct is called the absolute refractory period (see Fig. 11.12). The nerve fiber in this state may be likened to the gunpowder fuse after the spark has passed leaving only a trail of ash a fresh train of powder must be laid before a second spark can travel the same path. The duration of the absolute refractory period of frogs nerve is from 0.002 to 0.003 second and of manimalian nerve about 0.001 second. It is followed by a longer period during which repolarization of the nerve fiber is occurring. This is called the relative refractory period. It has a duration of from 0.01 to 0.02 second. During this time, the excitability and conductivity of the fiber is below normal a stronger stimulus than usual being required to set up an impulse which as judged by the action current is of smaller magnitude. These properties improve progressively reaching the values for resting nerve at the termination of the relative refractory period.

The velocity of the nerve impulses. In mammalian myelinated nerves, the impulse travels at a velocity of from 90 to 100 meters per second—that is, at about the speed of a revolver bullet. The impulses in unmyelinated mammalian nerve and in all nerves of cold-blooded animals, whether myelinated or unmyelinated, travel at much slower rates. Nerve fibers of large diameters transmit impulses at a higher velocity than those of small diameter. In the small nonmedullated fibers of some cold-blooded animals, the rates of conduction of the nerve impulse are less than 2 meters per second (Table 11-3).

### Table 11-3

### Nerve velocities

That there are at least two types of sensory nerve with different conduction velocities can be deduced from a common experience. When, for example, one stubs one's toe or receives a hard knock upon the shin, a sharp evanescent pain is felt which after a short interval is followed by a duller more enduring and much more unpleasant kind of pain. The impulses causing the initial pain are carried by nerves of relatively large diameter with a rapid conduction rate; the second pain is dependent upon nerves of smaller diameter and of slower conduction. Upon the nerves conducting at the more rapid rate (30 to 100 meters per second), the protective or withdrawal reflexes depend. Were consciousness not apprized of an injury until the slowly conducted impulses arrived, irreparable damage might be inflicted upon a part before the warning message to protect it had been received.

The frequency of nerve impulses. The length of the absolute refractory period determines the maximum frequency at which the nerve fiber is capable of transmitting impulses. In mammalian nerve, for example, the absolute refractory period has a duration of  $\tau_0^2 \sigma_0$  second. In other words, a period of  $\tau_0^2 \sigma_0$  second must clapse after the passage of an impulse before the fiber can again conduct. The maximum frequency is, therefore, around 1000 per second. Action currents of this frequency have actually been demonstrated by means of the cathoderay oscillograph A single fiber in the nerve trunk was separated from its neighbors by careful dissection and laid across a pair of electrodes connected to the recording apparatus.

The relation of the frequency of the impulses to the magnitude of the response. It has been mentioned above that the individual nerve impulse cannot, so far as is known, be varied in magnitude or in any other character by altering the strength of the stimulus. It may naturally be asked then, "Why does a stronger stimulus produce a greater effect? A strong stimulus for example, causes a more intense sensation than a weaker one when applied to a sensory nerve ending, or a more powerful contraction when applied to a motor nerve. The greater sensation caused by the stronger stimulus is due entirely to the higher fre quency of the impulses. Increasing the strength of the stimulus causes a rise in the frequency of the impulses discharged along each nerve fiber when applied to a nerve trunk the stronger stimulus also excites more nerve fibers than the weak one. These two factors, namely, the frequency of the impulses in each nerve fiber and the number of fibers involved—that is the number of impulses reaching consciousness in a unit of time—determine the intensity of the sensation.

irrequency of the impulses in each nerve incer and, the number of noers involved—that is the number of impulses reaching consciousness in a unit of time—determine the intensity of the sensation. Similarly, the magnitude of the muscular response when a motor nerve is stimulated is related to the frequency of the impulses discharged along the individual nerve fibers, as well as upon the number of fibers excited. Each nerve fiber of a mammalian motor nerve terminates by dividing into some hundred branches, each of which supplies a muscle fiber. Now, as we shall see (p. 488), when a muscle is stimulated by a single electric shock it gives a single contraction or inition, if stimulated by a series of such shocks in rapid succession, the resulting twitches become fused into a sustained contraction or itetanus which exerts a force considerably greater than that of the single twitch In a voluntary movement of the human arm for example, the impulses pass from the central nervous system to the muscles at varying fre quencies—from 5 to 50 or more per second. At the lower frequencies, the individual contractions are incompletely fused and only a proportion of the muscle fibers are activated, the movements are weak. Com plete fusion of contractions occurs at the higher frequencies a powerful movement resulting.

In some types of contraction, such as the tonic contraction of the muscles which maintain the body's posture, the impulses are transmit ted at moderately high frequencies, but only a proportion of the nerve fibers going to the muscle are active at one time—that is, the impulses are discharged from the central nervous system asynchronously along the several fibers composing the motor nerve. As a consequence, the

<sup>&</sup>lt;sup>1</sup> The single motor nerve fiber and the group of muscle fibers which it supplies is called a motor unit

fibers of the muscle are never all contracted at once; they contract in rotation. Since the muscle fibers respond in relays, some resting while others contract and vice versa, this type of contraction, though generally less powerful than that which results in movement, is able to be maintained for relatively long periods without fatigue

The nature or quality of sensation. The sensation aroused by stimulating a sensory nerve ending varies with the particular nerve ending excited. For example, stimulation of certain nerve endings in the skin causes a painful sensation, of others a sensation of touch, of others, as the contraction of the contraction of

The nature or quality of sensation. The sensation aroused by stimulating a sensory nerve ending varies with the particular nerve ending excited. For example, stimulation of certain nerve endings in the skin causes a painful sensation, of others a sensation of touch, of others, again, a sensation of heat or of cold. The optic nerve transmits impulses which arouse visual sensations. The acoustic and olfactory nerves and the nerves of taste, when stimulated, give rise to their own peculiar sensations. To what is this specificity of the sensory nerves due? There are two possibilities: (a) The fibers constituting the several types of nerve may differ in some fundamental way from one another, so that the impulses set up may have some distinguishing characteristic. In other words, each type of fiber may convey to consciousness a message peculiar to that type of fiber (b) The fibers themselves may be unspecific and the impulses which they transmit may be all alike, the different kinds of sensations—pain, touch, visual, auditory, etc.—being then dependent upon the particular region of the brain where the impulses are received. That is to say, the different regions of the brain interpret, each in its own way, the messages that come to them, though such messages are all identical in their general characters.

The latter conception appears to be the true one. All the evidence indicates that the impulses, along whatever type of nerve they are transmitted and though they may differ in minor details, are essentially the same in character For example, the action currents recorded from the optic nerve, from a motor nerve, or from a cutaneous sensory nerve, show no characteristic differences which might reasonably be considered as a basis for the several qualities of sensation. The center in the brain in which the impulses arrive is the determining factor. Impulses reaching one part of the cerebral cortex cause a visual sensation but, if these same impulses could be directed to the auditory center of the brain they would, according to this view, give rise to a sensation of sound. Or again, if it were possible to stimulate the receptors of the eye by a sound, and those of the ear by a light, the sensation to which such stimulation gave rise would still be one of light or of sound, respectively. For example, though the rods and cones of the retina are the receptors for sight, and are so specialized that light rays constitute the most effective, indeed the only really adequate, stimulus, we know

from our own experience that they, or the optic nerve fibers can be excited by mechanical means—e.g., a blow or firm pressure upon the eveball when a flash or a ring of light is seen

By way of summary it may be said that whereas the intensity of a sensation (the brightness of a light the loudness of a sound etc.) depends upon the frequency of the impulses the nature of the sensation is dependent upon the area of the cerebral cortex in which the impulses are received.

Chemical changes in nerve. Until some years ago attempts had failed to demonstrate any increase in oxygen consumption or of earbon dioxide production by nerve during its activity. Nor could any rise in temperature of the nerve be demonstrated during the passage of the impulse. It was therefore thought that the conduction of the nerve impulse was a purely physical process. By the application within recent years of more refined methods of investigation it has been shown quite conclusively however that conduction in nerve is essentially chemical in nature. A resting nerve produces a small quantity of heat (70 × 10<sup>-4</sup> calories per gram of nerve per second) which is increased by over 50 percent during activity. The resting nerve also consumes oxygen and produces carbon dioxide the grseous exchanges under going a definite increase when the nerve is stimulated.

Carbohydrate is not apparently the fuel from which the energy for the conduction of the impulse is derived. During stimulation no reduction in the glycogen content of the nerve occurs the disappearance of sugar is not accelerated and lactic acid is not produced. Nerve conduction differs therefore in this respect from muscular contraction (p. 497). The immediate source of the energy for nerve conduction is furnished by the explosive breakdown of phosphocrtatine. The phosphocratine undergoes resynthesis after the passage of the impulse—
i.e. during the refractory period—the conductivity of the nerve being thus restored. The source of the energy for the recovery process is unknown. That the recovery process is essentially different from that occurring in muscle is evident from the fact that it occurs in the absence of oxygen. A nerve placed in nitrogen will therefore continue to respond to stimulation for a relatively long time—3 hours or so

# The Physiology of Skeletal (Voluntary) Muscle

The minute structure of skeletal muscle has been briefly described in Chapter 2

Functional anatomy The char acteristic and outstanding prop erty of muscle is its contractility. and the most obvious function of skeletal muscle is to move the limbs head trunk and other parts The muscle fiber is a highly complex mechanism primarily di rected toward the conversion of energy derived from chemical re actions to mechanical energy which is exhibited as movement or same kind of force. Most skele tal muscles as the name implies are attached to bony structures though a few such as those of the face are attached to skin others are attached to cartilage and some to Ireaments or to other muscles One of the bones to which a mus cle is attached is usually held more or less stationary by the ac tion of other muscles the prime mover (see below) thus is pro vided with a fixed point for its contraction This attachment of the muscle is usually called its origin The other extremity of the muscle especially of a limb mus cle (e g the biceps of the arm) is most usually attached to a bone




Fig 11 13 Showing the action of a mus le crossing a joint When the muscle (in this case the biceps of the arm) contracts it pulls upon its insertion into the bone (radius) of the forearm and bends (flexes) the elbow joint

which moves upon some type of joint when the muscle contracts this attachment is called the insertion of the muscle. In the case of the muscles of the limbs and of certain other parts the muscles are situ ated with their insertions distal to their origins (i.e. more toward the fingers or toes) and when the muscle contracts its insertion moves toward its origin which remains stationary (Fig. 11.13). Most volun tary muscles are inserted not directly into the bone but through the medium of a strong touch inelastic fibrous cord called a tendon (or sinew in popular language. Fig. 11.14). Tendons vary greatly in length in man this may be from a fraction of an inch to more the foot. Sometimes a muscle has a tendon at each of its extremities.




Fig. 11 14 Showing the blending of voluntary muscle fibers with fibers of the tendon, and of the tendon with the periosteum (covering of the bone).

The form of muscle and the arrangement of its fibers. Muscles are of many shapes and sizes. Their shapes are variously described as fusi-form (spindle shaped), quadrilateral or rhomboid, triangular, strap-like (such as the long muscle on the front and inner side of the thigh and knee), and pennate (like a feather) in which the fibers pass from either side to a central tendon which they join obliquely (see Fig. 11.15). This arrangement of the fibers is called bipennate. In other muscles, only one half of a feather is represented, the fibers passing obliquely to only one side of the tendon; they are called unipennate. In most muscles, however, the fibers pass in a direct line—line of




Fig 11.15 Diagram showing four different types of arrangement of the fibers in voluntary muscles (Description in the text.)

Fig 11.16 Illustrating Glisson's experiment. (See text )

pull-from origin to insertion

When a muscle (e g, the biceps of the arm) contracts, it appears to increase in volume, and the ancient and medieval physicians thought that such actually occurred as a result of what they called animal spirits flowing into the muscle through its nerves Francis Glisson, an English physician (1597–1677), showed by a very simple experiment that the




Fig 11 17 Showing the branches of a single nerve fiber terminating in a group of muscle fibers of a motor unit

muscles of the human forearm did not increase in volume when they contracted, they actually diminished a little in bulk. He had a man insert his arm up to the elbow into a tall glass vessel provided with a side tube (see Fig. 11.16). The vessel was then filled with water to the brim and a bandage bound around the top of the vessel and the arm at this point in such a way as to prevent leakage. When the muscles of the arm contracted no rise in the level of fluid in the side arm occurred, a slight fall was observed.

Red (dark) and white (pale) muscles We are all familiar with the dark and light flesh of fowl In mammals also there are two types of muscle, the dark red and the light colored. The division into these two types is more pronounced in the cat and rabbit than in the monkey or man. The pale muscles (like the wing muscles of birds) contract more rapidly than the dark colored, and require a higher frequency of stim-

<sup>&</sup>lt;sup>2</sup> The slight reduction in volume was probably due to the expression of blood from the vessels by the compressing force of the contraction

ulation in order to telanize them (p. 488) The two kinds of muscle differ histologically the fibers of the dark variety have as compared with those of the pale muscles more abundant sarcoplasm are more opaque and show less distinct cross striations. The dark muscles as well as giving a slower contraction also fatigue less readily. They constitute in man mainty those muscles (e.g., the extensors of the tong) which are capable of sustaining a position of the body such as standing for a long period. In some human muscles both kinds of fiber are present toether.

The innervation of skeletal muscle motor end plate and motor units Each muscle cell receives both a motor and sensory innervation. Each nerve fiber of a spinal motor nerve which is an axon of a cell in




Fig. 11.18 Showing a nerve fiber ending in a muscle—the myoneural junct on (After Couteau redrawn)


the anterior horn of the spinal cord divides into some 100 branches which supply a corresponding number of muscle fibers. The anterior horn cell (including its axon the nerve fiber) is called a motoneuron and the group or bunch of muscle fibers which it innervates is designated a motor unit (Fig. 11-17). The muscle cell possesses a specially organized structure known as the motor end plate. Each axon branch of the motor nerve fiber as it approaches the muscle fiber for which it is destined loses its myelin sheath and terminates on the motor end plate as a flat expansion. This contact (there appears to be no actual union) between the terminal of the nerve fiber and the end plate is called the myoneural or neuromuscular junction (Fig. 11-18). It exhibits many of the properties of the synapse existing between neurons (or 510).

The sensory nerve fibers also lose their myelin sheaths and end in a special fusiform organ within the fiber known as the muscle spindle. The spindle consists of a group of muscle fibers differing from those of the rest of the muscle and enclosed in a well-defined capsule. It is a receptor of kinesthetic sense (p. 514 and Pl. 9A).

Muscular movements, Muscles which bend a limb at a joint—e.g.,

Muscular movements, Muscles which bend a limb at a joint—e.g., those which raise the thigh at the hip joint, or bend the elbow or knee—are called flexor muscles (or flexors). Those which straighten the limb—lower the thigh at the hip joint, or straighten the arm at the elbow or the leg at the knee—are called extensor muscles (or extensors). Muscles which move the limb or other part in a direction away from the mid-line of the body, or of a limb, are called abductors; and those executing the opposite movement—that is, bringing the limb or other part toward the mid-line of the body or of a limb—are known as adductors. There are also muscles which rotate a limb In the case of the movements at the ankle joint, confusion is avoided by adding the qualifying prefixes, plantar- and dorsi- to the word flexor. Thus, turning the foot toward the ground is called plantarflexion. Dorsiflexors and plantarflexors are corresponding terms given to the muscles which carry out these movements.

In performing a given movement—e.g., bending of the leg at the knee or of the arm at the elbow—the several muscles carrying out the movement act as a "team" In the bending of the arm at the elbow, for example, the muscles which execute the actual movement (biceps, etc.) are called prime movers, but we shall see (p. 517) that the oppositely acting (extensor) muscles must relax, the muscles which straighten the elbow (such as the triceps) are therefore called antagonists. The functions of these muscles are, of course, reversed when the elbow is extended, the triceps is now the prime mover and the biceps the antagonist. Fixation muscles are those which hold stationary the bone or bones giving origin to the muscle, thus enabling it to move the bone into which its tendon is inserted. Thus, in raising the arm from the side, muscles about the shoulder joint contract to hold the shoulder blade (scapula) and collarbone (clavicle) steady. Muscles that serve to steady a joint crossed by the tendon of the prime mover, and in this way enable the prime mover to contract to best advantage without loss of power, are called synergists. Thus, when the muscles (flexors of the fingers and wrist) which clench the fist contract, there is a tendency for the wrist to be sharply flexed. Normally this does not occur because the muscles on the back of the forearm (extensors of the wrist), whose



F g  $\,11\,19\,$  Sho ving apparatus for st mulating an isolated frogs muscle and the method of recording the contraction

tendons cross the back of the wrist contract and thus give rigidity to this joint

Method of recording contractions of an isolated muscle Our knowledge of the physiology of muscular contraction has been gained largely from studies of the isolated muscle of the frog The muscle of the calf—the gastrocnemus—is the one most generally employed This muscle is attached above to the lower end of the thighbone (femur) and below through a strong tendon called the tendon of Achilles (tendo Achillis or tendo calcaneus) to the bone of the heel In preparing the muscle for an experiment it is removed from the animal to gether with its nerve supply the sciatic and the lower end of the fe

mur The preparation is then suspended by fixing the section of bone in a clamp, and fastening the lower tendinous end of the muscle to a light lever, as shown in Figure 11 19. The writing lever is adjusted so as to make light contact with the surface of a revolving drum, motivated by clockwork and covered with smoked paper. This instrument is called a kymograph. As the drum revolves, the soot is removed by the point of the lever, which thus traces a white horizontal line so long




Fig 11 20 The inductorium A primary coil B secondary coil

as the lever remains stationary. Any shortening (contraction) of the muscle causes an upward movement of the lever which then inscribes a curve on the rotating drum.

A tuning fork of the required frequency of vibration (e.g., 1760, 270 per second) is arranged with a writing point to mark the time intervals at the bottom of the tracing, or some device run by clockwork may be used A signal magnet is also set up in the primary circuit of the inductorium (see below) to indicate the instant that the stimulating current, which also activates the signal, is made or broken

The inductorium. The induced electric current is usually employed in classroom experiments for stimulating muscle and nerve, it can be finely graded in strength, and a series of rapidly repeated shocks can be applied to the excitable tissue. The instrument employed for this purpose is called an inductorium, and consists of two cylindrical spools or coils of wire called primary and secondary (Fig. 11.20). The primary coil has an iron core around which are made a few turns of thick wire. It receives current from a dry cell, a switch or key being inter-

posed in the circuit. The secondary coil is composed of many turns of fine wire Its core being hollow, it is capable of being slid like a sleeve over the primary coil It runs on parallel metal bars and can be pivoted as well around its transverse axis When the current from the dry cell enters the primary coil a current is induced in the secondary coil The low electromotive force of the primary circuit is transformed to the high electromotive force of the induced current which is led off to stimulating electrodes fastened to binding posts one on either end of the horizontal bars along which the secondary coil slides. The primary circuit may be made and broken by means of the key or by the rapid movements of an automatic interrupter or vibrator on the instrument itself Nerve or muscle can then be stimulated by a series of rapidly alternating make and break induced currents. The break shock of the induced current is more effective as a stimulus than is the make shock because the primary current causes an induced current in the primary coil itself as well as in the secondary and its direction is opposite to that of the primary current as a consequence the intensity of the lat ter as well as its rate of rise is reduced (see p 467) In the case of the break shock the self induced current cannot weaken the primary cur rent since this has already ceased to exist. As a matter of fact, the di rection of the self induced current at the break is the same as that of the primary current

The current induced in the secondary coil is increased or diminished by sliding this coil toward or away from the primary coil. The induced current can also be graded by rotating the secondary coil upon its privot and altering the angle which it makes with the primary coil. When the secondary and primary coils have their long axes at right angles to one another there is no induced current. A current is induced as the secondary coil is moved from the right angle toward the hori zontal position, and increases steadily the farther it is turned in this direction reaching a maximum for any given distance from the primary coil when the long axes of the two coils are in line.

The simple muscular contraction or "twitch" Muscle, like nerve responds to any one of the four types of stimulus—electrical mechanical thermal or chemical. The muscle may be stimulated directly—that is by placing the electrodes in contact with its surface—or through its nerve. When stimulated in either of these ways and with a single stimulus—e.g. a make or break shock—the muscle contracts and having raised the writing lever to a certain height, relaxes again and allows the lever to return to its original position. Thus a curve such as that shown in Figure 11.21 is inscribed. This simple brief contraction is called a

muscle twitch: it is due, of course, to the shortening (contraction) in unison of the numerous fibers of which the muscle is composed. A simple contraction or twitch does not occur in the living animal under ordinary physiological conditions, but the reflex contraction of the extensor muscles of the thigh caused by a tap on the patella tendon (knee jerk) is of this nature. The curve shown in Figure 11.21 is marked off into three parts to represent corresponding phases of the muscular response The stimulus is applied at A, shortening of the muscle commences at B. and relaxation at C. The distance from A to B indicates the period elapsing from the application of the




Fig 11.21 Record of a muscle twitch A-B, latent period; B-C, contraction phase, C-D, relaxation phase Time indicated by lower tracing (\(\gamma\_{00}^{\dagger}\)) second) caused by a vibrating juming fork (After Stirling)

stimulus to the commencement of the contraction. This is called the latent period; its duration is about 0.01 second. The next period (B-C), during which the lever rises from the horizontal line (base line) to the summit of the curve, is the contraction phase, it lasts for approximately 0.04 second. The part of the curve from C to D represents the relaxation phase. This has a duration of about 0.05 second.

As determined by the method described above, the total duration of the contractile process of the frog's gastroenemus muscle, including the latent period, is found to be about a tenth of a second. The muscles of warm-blooded animals have considerably shorter contraction times.

The latent period of the muscle, as measured from the instant of stimulation of its nerve, is made up of: (a) the time taken for the impulse to travel along the nerve from the point of stimulation to the muscle; (b) the very brief interval during which the initial chemical changes are occurring in the muscle fiber and tension is being developed; (c) the inertia of the lever. The last mentioned, (c), can be almost entirely eliminated by refined methods of recording—e.g., some form of recording in which the contraction of the muscle moves a mirrored surface. This reflects a beam of light which is photographed. The light beam, which is substituted for the material lever, is free from mertia Furthermore, its length can be such that great magnification of the movements of the reflecting surface can be obtained. The true

latent period of muscle—that is, the interval (AB) between the stimulation of the muscle fiber itself and the commencement of the shortening process—varies between 0 0025 and 0 004 second

The effect upon the force of the contraction of increasing the intensity of the stimulus II a series of stimuli of gradually increasing strength is applied directly to the muscle, contraction curves of graded heights will be obtained (see Fig. 11 22). The weakest stimulus which will excite the muscle is called the minimal or threshold stimulus. As the strength of each successive stimulus is increased, a greater and greater response is obtained from the muscle. Ultimately, a point is reached where the muscle contracts maximally—that is, no further in




Fig 11.22 Record of a series of contractions caused by a succession of stimuli of graded strength. The muscle lever traced each of these lines while the drum of the kymograph was at rest and was then moved by hand to a fresh position after each contraction.

crease in the height of the contraction curve results from increasing the strength of the stimulus. This is called the maximal stimulus

It was stated on page 471 that muscle like nerve, obeys the 'all-onnone law The grading of the muscular response to stimuli of graded strength as just described appears to contradict this statement. The explanation is that a muscle is composed of a number of units—the muscle fibers. The force of the contraction of the whole muscle is dependent upon the number of fibers excited by the stimulus. A strong stimulus excites a larger number of fibers than does a weak one, and with a maximal stimulus the current spreads to involve all the fibers Studies made upon a single muscle fiber have shown that it does not respond by graded responses as the strength of the stimulus is increased. It responds maximally or not at all—that is, it obeys the "all or none" law.

Summation and tetanus. If a second stimulus is sent into an isolated frog's muscle within the latent period of the first, no obvious effect is produced If, however, the second stimulus is applied while the muscle is contracting in response to the first stimulus, it contracts again, the second contraction is added to the first, the shortening of the muscle being then greater than that caused by a single stimulus. The greatest

effect of the second stimulus is observed when it is applied to the muscle near the height of the contraction caused by the first. This phenomenon, whereby one contraction is added to a previous one to produce a greater total shortening of the muscle, is called *summation* (Fig. 11.23). A third contraction may be added to a second, a fourth to a third, and so on. When the stimuli sent into a frog's gastroenemius




Fig. 11.23 Illustrating summation and tetanus 1, the second stimulus x was applied after relaxation of the muscle, two separate contractions of equal amplitude result; 2, summation, the second stimulus x being applied at a shorter interval after the first than in 1, 3 and 4, incomplete tetanus caused by a series of stimuli applied at short intervals apart, 5, complete tetanus resulting from a still more rapid rate of stimulation, the individual muscular responses being completely fused. (After Stirling and Howell)

muscle are at a slower rate than from 15 to 30 per second, the individual contractions can be distinguished—that is, they are not completely fused. The contraction curve shows smaller or larger waves synchronous with the rate of stimulation. At rates of stimulation higher than this, the responses are completely fused, the contraction curve being perfectly smooth. The curve rises to a maximum height, which is considerably greater than that caused by a single stimulus, and remains at this height as long as the stimulation lasts or until the muscle becomes fatigued. A sustained contraction of this nature is called *complete tetanus*. The contraction caused by slower rates of stimulation, and in which the individual responses are distinguishable, is referred to as incomplete tetanus.

The rate of stimulation at which complete fusion of the contractions occurs varies with the speed at which a given type of muscle normally contracts. Thus, rapidly acting white muscles require a higher rate of stimulation to produce tetanus than do the more slowly acting red muscles, and the rate required by smooth muscle is slowest of all (5 stimula per second). In the rapidly acting muscles of the wings of m sects, contractions are not fused until the rate of stimulation reaches a rate of 300 per second and upward, and even a human eye muscle funternal rectus) requires a stimulation rate of 350 per second. The re



Fig. 11.24. Illustrating treppe contracture, and fatigue. The muscle was stimulated at intervals of 1 second. The contractions are very close together because the kymograph drum was moving slowly (After Howeli.)

quired rate for red muscle of the human calf, on the other hand, is only 30 stimuli per second. In man, the voluntary contractions of the muscles of the limbs are brought about by the discharge of impulses along the motor nerves at rates of from 5 to 50 per second. At the lower rates the contraction is an incomplete tetanus, but at the higher frequency complete fusion of the contractions results (see also p. 476).

Treppe or the "staticrase" phenomenon. When a muscle is stimuli.

Treppe or the "staircase" phenomenon. When a muscle is stimulated repeatedly at regular intervals, the first few contractions of the series increase successively in amplitude This phenomenon has already been described for heart muscle (p. 160). It is due, apparently, to the beneficial effect exerted upon the irritability of the muscle by the chemical products of the first few contractions, and by the rise in temperature (Fig. 11.24).

The effect of loading upon the contraction of a muscle. A muscle contracts more forcibly if suitably loaded The increase in the height of the contraction increases up to a point with each weight that is added to the scale pan suspended from the muscle, as shown in Figure

11 19, page 484 The force of the contractions then decreases again as the load is gradually increased (Fig. 11 29, p. 495). For an isolated frog's muscle, the maximum height is attained at a loading weight of between 30 and 50 grams. When a loaded muscle is stimulated to contract, the first effect is a sudden lengthening of the muscle fibers. This sudden stretch causes tension in the muscle which is immediately followed by shortening. The stretching of the fibers in some way enables the muscle to mobilize more energy than is possible for the unweighted muscle. We have seen the same phenomenon in the case of cardiac




Fig. 11.25. Showing the effect of oxygen lack upon the onset of fatigue. Records from a pair of sartorius muscles of frog stimulated at 1 second intervals. A in nitrogen. B in oxygen. (After Fletcher.)

muscle (p 193) Throughout the shortening of the muscle the tension remains constant. This type of contraction is therefore called isotonic (Gk 1808, equal + tonus, tension). But if the muscle is loaded with a weight that it is unable to lift or is attached to a stiff spring that it cannot bend only a very slight shortening of the muscle fibers can occur. Such a contraction is called isometric (Gk 1808, equal + metron, measure).

Fatigue. If direct stimulation of an excised muscle—that is with the electrodes placed upon the muscle—be continued after the contractions have reached their maximum amplitude the irritability of the muscle gradually becomes depressed due to the accumulation of lactic acid (Fig. 11.25). The concentration of lactic acid at which loss of irritability of the muscle occurs and at which it fails to respond to stimulation, is around 0.5 percent.

The onset and development of this state of fatigue are more rapid, the higher the rate of stimulation Isolated muscle, or one in the intact animal the blood supply of which has been occluded, also fatigues more rapidly than one receiving an adequate oxygen supply Oxygen

is required for the removal of the lactic acid produced during contraction. In the muscle deprived of its blood supply this, of course, does not readily occur. Lactic acid accumulates, and it is to such accumulation that the loss of irritability of the muscle is attributed (see p. 498).

When a muscle is stimulated, not directly, but through its nerve, fatigue supervenes much sooner—long before the lactic acid concentration has reached the value at which it depresses muscular irritability. We know that nerve is highly resistant to fatigue. There must, then, be



Fig. 11.26 Illustrating the ready fatigability of the myoneural junction—i.e., before the muscle itself. The muscle was stimulated through its nerve to the arrow, S, and then directly.

a third structure or tissue in the nerve-muscle preparation that is peculiarly susceptible to fatigue; this is the junction between nerve and muscle—the myoneural junction—which exhibits the properties of a synapse. The early fatigability of the myoneural junction can be demonstrated by a simple experiment. If the nerve to the muscle is stimulated repeatedly until the muscle fails to respond, a stimulus of the same strength as before applied directly to the muscle causes a vigorous contraction (see Fig. 11.26).

Contracture. Under certain conditions, relaxation of the muscle takes place very slowly; it may remain in the partially contracted state for a considerable length of time. This state of the muscle is referred to as contracture. If a muscle in a state of contracture is stimulated at relatively long intervals, it responds by further shortening, each contraction being then superimposed upon the sustained contraction caused by the previous stimulus. Contracture is brought about by the action of certain drugs, especially veratrine, and by fatigue. It also sometimes occurs in apparently normal muscle during the first of a series of contractions and then passes off (Fig. 11.24).

The effect of temperature upon muscular contraction. Warming a muscle causes it to respond more rapidly and vigorously to stimulation Contraction and relaxation phases are both shortened, the latent period is also of shorter duration. Lowering the temperature of the muscle exerts the opposite effect, the contraction being weaker and all its phases lengthened. Cold blooded animals, such as snakes and frogs, whose body temperature varies with that of the environment, move




Fig 1127 The effect of temperature upon the contraction of a frog s gastroenemus muscle Tracing should be read from left to right Figures are the temperatures (°C) of the salt solution in which the muscle was immersed (After Pembrey and Phillips)

sluggishly, therefore, in cold weather. The activity of mammals, on the other hand, whose muscles are maintained at a constant temperature of around 98 5°F, is not influenced by changes in the temperature of the environment. Nevertheless, the more efficient action of the muscles after the commencement of exercise is due, in part, to the local rise in temperature resulting from the earlier contractions "Warming up" to one's work is a familiar expression (Fig. 11.27)

The effect of temperature upon the speed of muscular contraction conforms with Van't Hoff's law, which states that the velocity of chemical reactions, which, of course, are the basis of muscular contraction, is increased to between two and three times by a rise in temperature of 10° C

Muscle is inexcitable at very low temperatures. Frog's muscle, for

example, fails to respond at temperatures of from 32 to 35° F. On the other hand at an excessively high temperature (105 to 110° F for frog's muscle) the muscle proteins are coagulated, and death of the muscle results. At the same time, the muscle undergoes pronounced shortening and loses its natural translucency. This state is called hear rigor. It resembles rigor mortis, a term given to the stiffening of the muscles of the body generally which occurs in from 1 to 7 hours after death.

Electrical changes in muscle. Like nerve and other excitable tissues an action potential is developed in muscle when it contracts, the active region of the muscle being negative to the mactive region. A current of




Fig 11 28 The rheoscopic preparation (See text )

injury can also be demonstrated when one electrode is placed on a crushed or cut end of the muscle the other upon the uninjured surface and the two electrodes connected through a galvanometer (see Fig. 11 8, p. 470). The action current or the current of injury in muscle can be used to excite the nerve of another isolated muscle. This so called rheoscopic preparation is shown in Figure 11 28. When the nerve of muscle A (upper sketch) is placed so that one part rests upon the cut surface of muscle B and another part of the nerve is made sud.

denly to touch a point on the uninjured surface of B, a slight twitch of A occurs each time the contact is made. Similarly, if B (lower sketch) is stimulated to contract, a twitch of A occurs as the wave of contraction passes beneath a nerve which touches the contracting muscle at two points, a potential difference being established as the contraction wave spreads between the active (contracting) part of the muscle and the inactive (resting) part

Muscular work and efficiency. Work, as defined by the physicist, is the product of the load (expressed in grams, kilograms, or pounds) and the vertical distance (in millimeters, centimeters, meters or feet) through which it is raised According to the units of distance and weight employed, the work is expressed in grammillimeters, gramcentimeters, kilogrammeters, or foot pounds

The isolated muscle can be made to perform work by stimulating it to contract and lift a weight A load of a known weight is attached to the lever at the point of attachment of the muscle, which is then stimulated The height above the base line of the tracing inscribed by the lever is measured This measurement, however, is not the actual distance through which the weight has been raised, for the shortening of the muscle has been magnified by the lever The distance as measured must therefore be divided by the magnification of the lever, which is usually five

The work performed by the gastrocnemius muscle of the frog is most suitably expressed in grammillimeters (gmm) For example, a muscle that lifts a weight of 1 gram to a height of 2 mm does 2 gmm



Fig 11 29 Diagram to show the method of measuring muscle work Heights of contractions along vertical lines, weights in grams along the horizontal line (After Mc-Kendrick)

of work. The muscle, when loaded with a heavier weight, may not raise it so high yet may do more work. Thus, if it lifts a weight of 4 grams only 1 mm it does 4 gram of work. Starting with a light load and gradually adding to it for a number of successive contractions, it is found that the work performed increases up to a maximum and then falls off. There exists, therefore, an optimum load—that is, a load with which the muscle performs the maximum amount of work (see Fig. 11.29).

Since the work, as just defined, is the product of the load and the lift, no work is done by an unloaded muscle when it contracts, nor by one that contracts against a weight that it is unable to lift. In both these instances, however, there is an expenditure of energy which is derived from chemical processes taking place in the muscle, all of this energy appears ultimately as heat

The fibers of a muscle contracting against a resistance, such as a weight that it cannot move or a stiff spring that it cannot bend, of course, do not shorten The tension within the muscles increases, however Such a contraction is called isometric (Gk itos, same + metron, measurement) An isotonic contraction, on the other hand, is one in which the muscle shortens, but its tension remains almost constant introughout the contraction. In the body both types of contraction take place The isometric type is seen in those muscles that maintain the body's posture, such as the extensors on the front of the thigh. These exert a constant pull upon the leg, and thus hold the limb in the extended position to afford a firm support for the body. The movements of the limbs, as in walking, lifting objects with the hands, or throwing a ball, are brought about by isotonic contractions.

Of the total energy expended by any machine, a part appears as mechanical work, the remainder being dissipated as heat. The efficiency of the machine is defined as the percentage of the total energy that appears as work. Thus

Efficiency = 
$$\frac{\text{work}}{\text{total energy (work + heat)}}$$

As mentioned above, a muscle contracting isometrically or without a load performs no work, all the energy is dissipated as heat. Contracting with an optimum load, on the other hand, muscle has an efficiency of from 25 to 30 percent, which compares favorably with that of the best type of gasoline engine. By measuring the heat produced by a muscle during its contraction and the work done, its efficiency can be readily calculated from the equation given above. In making the calculation, the energy appearing as work and as heat must, of course, be expressed in the same units. We may express the work in heat units (e.g., microcalories). Now, the heat equivalent of a grammillimeter is 00235 microcalorie. Therefore, if the heat produced is 8.4 microcalories, and the work performed is 1200 gmm, then the total energy expenditure expressed as heat can be calculated as follows.

Heat produced = 8 4 microcalories
Work = 1200 pmm

Work = 1200 gmm Heat equivalent of the work =  $1200 \times .00235 = 2.8$  microcalories

The total energy expenditure is, therefore, 8.4 + 2.8 = 11.2 microcalories.

The efficiency is  $\frac{2.8}{11.2} = 25$  percent

The chemical constitution of muscle. Fresh muscular tissue, which composes the lean part of beef, pork, etc, contains about 20 percent of protein and around 75 percent of water. The remaining 5 percent is made up of minerals and various organic compounds (glycogen, glucose, lipids, steroids, and nonprotein nitrogenous compounds). The principal mineral in the muscle cell (fiber) is potassium, which has a concentration of around 400 mg per 100 grams; magnesium, phosphorus, sodium, and calcium are present in much lower concentrations. The chief nonprotein nitrogenous substances are creatine, phosphocreatine, adenosinetriphosphate, carnosine, and urea. The main proteins composing the contractile elements of muscle (the micellae) are myosin and actin. These two proteins are in very close functional association in the micellae; the combination, known as actomyosin, is of essential importance in the contractile process

The pigment of muscle, called myoglobin or muscle hemoglobin, is closely allied chemically, as well as in function, to hemoglobin.

The chemistry of muscular contraction. Muscle derives the energy for its contraction, ultimately, from the combustion of carbohydrate; in the final step of this process, oxygen is consumed and carbon dioxide (CO<sub>2</sub>) and water (H<sub>2</sub>O) produced. The actual contraction of the muscle, however, is brought about by the explosive separation of a molecule of phosphoric acid from adenosinetriphosphate (abbr. ATP), a compound composed of adenylic acid (a nucleotide, p. 366) and three molecules of phosphoric acid. Adenosinediphosphate (ADP) is thus formed. This reaction is the very first to occur when the muscle fiber contracts. The molecule of phosphoric acid, adenosinediphosphate (ADP) with the acquisition of phosphoric acid, the glycogen splits off a molecule of phosphorylated glucose, called glucose phosphate, which is converted through a series of intermediate reactions to fructose diphosphate (i.e., fructose combined with two molecules of phosphoric acid). The fructose diphosphate passes through a series of complicated reactions to form lactic acid. After the transfer of phosphoris from ATP to glycogen and the formation of ADP, phosphoreatine splits into

creatine and phosphoric acid. The liberated phosphorus is taken up by ADP and ATP reformed. thus

phosphoric acid + ADP ---- ATP

The conversion of ATP to ADP as just stated is the immediate source of the energy for the contraction of the muscle fiber From the breakdown of phosphocreatine is derived the energy for the reformation of ATP from phosphoric acid and ADP Phosphocreatine is also resynthesized the energy for the resynthesis being derived from the breakdown of glucose phosphate to lactic acid

Of the lactic acid produced one fifth is oxidized to carbon dioxide and water. The remainder (four fifths) is resynthesized to glycogen in the liver muscles and other tissues of the body. Each of the several chemical reactions described above is dependent upon a specific en zyme. Only one of these need be refurred to and it will be spoken of later.

The removal of phosphoric acid from ATP and its resynthesis as well as the breakdown of phosphocreatine can be carried out in the absence of oxygen. The breakdown of glucose phosphate to lactic acid and the resynthesis of phosphocreatine can only occur in the presence of oxygen. There are therefore two phases in the contractile process—a nonoxidative anaerobic or contraction phase and an oxidative aero bic or recovery phase. This latter is necessary to restore energy producing substances and permit continued contraction of the muscle.

In so far as its energy relationships are concerned the muscle fiber has been compared to a submarine which while submerged is driven by energy stored in electric batteries oxygen being not required this period corresponds to the anaerobic phase of the process in the muscle. The batteries are recharged when the submersible reaches the surface oxygen being then necessary. This corresponds to the aerobic or recovery phase of the muscle.


Though it will be clear from the foregoing account that the reac tions leading to lactic acid formation are required for the continued ac tivity of muscle contractions nevertheless can proceed for a time after

A muscle deprived of oxygen (i.e. in an atmosphere of n trogen) is therefore capable of tending to 4 mulat on for a considerable time. Later a ed. however is not removed, by fordation and resynthes is to glycogen but accumulatine reduces the irritable by the first but of the concentration of later a ed at which the muscle fals to respond is around of Sevent This is called the facer of a notion in N then oxygen is supplied to a muscle used by repeated sit mulation in n trogen the lattic acid is removed and the irritable if y of the muscle restored.

the production of lactic acid has been entirely suppressed; the reaction is therefore not essential for the actual contraction. The production of lactic acid from the breakdown of carbohydrate in the isolated muscle can be prevented by treating it with a drug called sodium iodoacetate; yet the muscle so treated responds to stimulation for some time, or until the phosphocreatine store of the muscle is exhausted. But phosphocreatine cannot be reformed, since the energy for the resynthesis is derived from the glycogen to lactic acid reaction. A muscle poisoned by iodoacetate obviously cannot continue to respond for as long a time as can a normal muscle. The glycogen to lactic acid reaction, therefore, though not directly responsible for the contraction, serves, as it were, to "rewind" the contractile mechanism—to "set the trigger" for the next contraction. Carbohydrate, then, is the ultimate source of the energy for the activity of the isolated muscle

Table 11-4 presents a summary of the chemical changes occurring during and immediately after its contraction

Table 11-4



The transformation by the muscle fiber of chemical energy into mechanical energy resulting in movement. The shortening (contraction) of the micellae is the fundamental mechanical change that occurs in the muscle fiber when it is excited to contract. We must now consider the possible way in which the loss of a molecule of phosphoric acid from adenosinetriphosphate (ATP) brings about this change. The contraction of a whole muscle is simply the sum of the shortening taking place in the millions of micellae that the muscle contains, and it is strongly suggested that the shortening of the micellae is due, in turn,

Fig. 11.30 Showing a myosin mole cule in an extended (1) and in a folded or contracted state (11)

to the shortening by folding or rolling up, of the long chains of molecules constituting the muscle proteins (Fig. 11 30)

In the ultramicroscopic study of proteins of hair (eg, wool feathers, etc.), it has been found that their molecules are arranged in long roughly paral lel and more or less folded chains The hair can be lengthened by stretching it, when released it returns to its original length This property of such structures is due. it is believed to the unfolding and refolding, like a concertina, of the long amino acid chains of its con stituent protein When heated or treated with a strong chemi cal, the natural condition of the

the hair is permanently shortened and its inherent elasticity destroyed as a result of the tight irreversible folding of the amino-acid chains of its proteins. Thus is explained the shrinkage of woolen fabrics.

Now, myosin like the fibrous proteins of wool and other hairs, is composed of long chains of amino acid molecules. A chemical change in the environment of the molecules (brought about by the splitting of a phosphoric acid group from adenosinetriphosphate it is supposed) causes folding of the molecular chains and, as a consequence, shortening of the micellae. When the chemical conditions which induced shortening of the micellae have disappeared, the latter return to their previous length, the muscle fiber relaxes

A chamilia molecule of a fibrous protein in the folded (contracted) and in the unfolded (relaxed) state is shown in Figure 11 30

Myosin an enzyme The surprising discovery was made a few years ago (1939) that myosin—one of the essential proteins in the muscle fiber, and which with actin, largely composes the micellae—appears to be itself an enzyme required for the splitting, with lightninglike rapidity, of adenosinetriphosphate 4

\* Following the usual practice of naming enzymes myosin is also known as adenosinetriphosphatase

As a machine the muscle fiber is unique. There is no man-made mechanical device with which it can be compared satisfactorily. The muscle is a chemical machine and a comparison has been drawn between it and the internal combustion engine in which the firing of a mixture of gas and air moves a piston. The comparison would be closer if an engine were conceived in which the spark which ignites the gas mixture and the piston were one and the same. A piston raised to a high temperature would, in a way, be comparable to myosin, for this protein, being the moving part of the muscle as well as an enzyme through which the initial energy is liberated, corresponds to both piston and




Fig. 11.31 Diagram showing the stages of heat production by muscle during and following a short tetanic contraction. The three rectangles represent the initial heat. I, heat produced at the commencement of the contraction; 2, that produced during the maintenance of the contraction, 3, relaxation heat, 4, recovery heat. (After Evans, Recent Advances in Physiology.)

spark. But the muscle fiber is not, like the gasoline motor, a heat engine utilizing the pressure generated by heated gas molecules for power. Muscle gains the energy for its contraction largely from the bonds which hold certain phosphoric acid groups to organic compounds, such as adenosinetriphosphate and phosphocreatine.

Heat production in muscle. It has been mentioned that during the isometric contraction of muscle, the total energy expended appears as heat. The heat production is divided into two phases, termed the initial heat and the recovery heat. The initial heat comprises the heat produced in chemical reactions during the contraction of the muscle (contraction heat) together with a smaller quantity which appears during relaxation (relaxation heat). The latter results from the conversion to heat of the energy exhibited as tension during the contraction phase. The initial heat is produced during the anaerobic phase of the contractile process; it, but not the recovery heat, is therefore generated by a muscle contracting in nitrogen The recovery heat appears after re-

laxation is over, and is due mainly to the oxidation of lactic acid (Fig. 11.31).

The phases of heat production are summarized in the following

- 1. Initial heat = 45 percent of the total heat,
  - a Contraction heat 65 percent of the initial heat,
    - b. Relaxation heat 35 percent of the initial heat.
- 2. Recovery heat = 55 percent of the total heat.

## The Physiology of Muscular Exercise

Oxygen debt. In strenuous muscular exercise of brief duration, only a fraction of the oxygen required for the recovery phase of the contracting muscles is breathed during the course of the exercise. For example, the oxygen requirement for a race of 100 yards, which takes only a few seconds to run, is 6 liters or more. Yet the quantity of oxygen which can be absorbed in this time is less than I liter. Indeed, a short race can be run with the breath held. In other words, the sprinter goes into debt for oxygen during the race, and pays up later. Thus the oxygen consumption after the exercise period is much higher than during an ordinary period of rest. The extra oxygen is utilized in the removal of the lactic acid produced during the exercise. The size of the oxygen debt for any piece of work is determined by measuring the oxygen consumption of the after-period of exercise, and subtracting from the result the figure for the oxygen consumption of a corresponding period of rest. During very strenuous muscular effort, the oxygen debt may amount to 15 liters or more

The ability of the muscles to contract, without receiving the full oxygen requirement until the work has been completed, has an obvious advantage. Short periods of strenuous exercise can be undertaken which would otherwise be impossible, for the respiratory and circulatory systems are quite unequal to the task of furnishing, during the exercise, the great volume of oxygen which is ultimately used in the recovery process. The maximum quantity of oxygen that can be delivered to and consumed by the tissues of a large healthy man is not more than about 2 liters per minute and, of most persons, considerably less than this.

In light exercise no oxygen debt is incurred. The lactic acid is removed as it is produced; in other words, the body, in so far as oxygen consumption is concerned, "pays as it goes." Lactic-acid production and its removal are nicely balanced; a "steady state" is established. When the "steady state" has been reached, a person engaged in an athletic performance of some duration—such as running, rowing, etc.—breathes with less effort. As the expression goes, he has got his "second wind."

The source of the energy for muscular work. We have seen that the energy for the contraction of the isolated muscle is ultimately derived from earbohydrate. Glycogen is broken down through glucose phosphate and fructose diphosphate to lactic acid. In the numerous studies of the respiratory exchanges of the isolated gastroenemus muscle of the frog during contraction, respiratory quotients of 1.0 (p. 354) are obtained. In the intact animal carbohydrate is also the fuel for short bouts of muscular exercise After a sprint, for example, a fall in blood sugar may occur, and the ingestion of glucose prior to a race is now recognized as a valuable means of postponing fatigue and enhancing muscular performance. In prolonged and exhausting work the carbohydrate stores become depleted, fat is then burned to furnish the required energy. Protein apparently is not utilized in muscular exercise, or, if so, to a very small extent.

or, if so, to a very small extent.

The adjustments in the respiratory and circulatory systems in muscular exercise. During strenuous muscular effort, the quantity of air breathed is from 300 to 400 percent greater than during a resting period of the same length. The greater pulmonary ventilation is brought about by an increase in the respiratory rate, as well as by an increase in the volume of each breath. The increased breathing occurs at the very outset of exercise; indeed, it has been shown in experiments upon man that it may actually anticipate the exercise by a brief interval—that is, an effect upon the respirations may be noted at the instant a signal has been given to start some muscular act, and before the muscles contract. Chances in heart rate and blood pressure may also prea signal has been given to start some muschan are, and octobe the mus-cles contract. Changes in heart rate and blood pressure may also pre-cede the contractions of the muscles. The influence upon the respiratory cede the contractions of the muscles The influence upon the respiratory and circulatory systems at the commencement of or preceding the exercise is due to impulses from psychic levels of the brain, and probably also from the motor area of the cerebral cortex, upon the centers in the medulla. The efficiency with which the work is performed is also undoubtedly influenced by emotional factors These act most likely by bringing about more perfect coordination of respiratory and circulatory mechanisms, but also, probably, through a direct effect upon the muscular contractions themselves. A man is spurred to greater effort, and is a more effective worker, if the work interests him than if he is bored with it, or does it as a matter of routine. In emergencies or during excitement—"in the heat of the moment"—feats of strength of en504

durance are accomplished which, in the absence of the emotional factor, would be impossible 1t is likely that adrenaline liberated into the blood stream through nervous influences plays a role in some instances (pp. 420-1)

If the exercise is severe, chemical factors soon come into play to increase the pulmonary ventilation. The production of lactic acid and carbon dioxide in the muscles causes a slight increase in the hydrogen ion concentration of the blood which acts as a stimulus to the respiratory center. The greater carbon dioxide produced by the exercise is thus eliminated.

The adaptations of the circulatory mechanism in exercise are of a highly complex nature. The blood pressure rises to from 160 to 180 mm Hg as a result of constriction of the vessels in the splanching region. The carotid sinus depressor reflex is in abeyance or neutralized by pressor reflexes. The vessels in the muscles themselves dilate. A larger proportion of the blood volume is diverted, therefore, to the active muscles than during rest. The vasodilatation in the muscles is due, mainly, to chemical substances-namely, carbon dioxide and lactic acid, products of the contractile process-acting directly upon the vessels Not only do the muscles receive a greater proportion of the total blood volume during exercise than during rest but, as a result of the contraction of the spleen, the volume of circulating blood may be in creased by from 20 to 25 percent. The blood also circulates more rapidly, the output of the heart and, consequently, the quantity of blood pumped per minute through the systemic and pulmonary vessels is in creased many fold. In a large robust man performing arduous work, the output of the heart may amount to 35 liters (nearly 8 gallons) or more per minute. The greater cardiac output is the result of the greater venous return-that is, of the increased volume of blood flowing through the contracting muscles and carried along the great veins to the right side of the heart

The effect of muscular exercise upon the heart rate varies in differ ent persons. In those untrained to muscular work, the heart accelerates markedly, a rate of from 110 to 120 per minute being not unusual, whereas the heart of the athlete may show little or no acceleration during muscular feats of a highly exacting character, such as sprinting or rowing. Through the respiratory and circulatory mechanisms just described, the maximum load of oxygen is delivered to the contracting muscles. The oxygen supply is further augmented through the effect which the carbon dioxide and lactic acid produced in the muscle, and the local rise in temperature, exert upon the liberation of oxygen in

oxyhemoglobin. As mentioned on page 251, increased acidity and a i rise in temperature cause the hemoglobin to give up a greater part of its oxygen load.

Athletic training causes a moderate enlargement of the normal heart. The enlargement is purely physiological, and is commensurate with the accompanying increased bulk of the skeletal muscles. Enlargement (dilatation and hypertrophy) of the heart occurs when the heart is damaged; and athletes, of course, like anyone else, may be subjects of heart disease. Because some athletes have shown cardiac dilatation and hypertrophy or other evidence of heart disease, it used to be thought that athletics, if engaged in too enthusiastically, led to heart disease. It is now generally admitted, however, that the wellconditioned heart of the young adult is not damaged by even strenuous exercise. The skeletal muscles fatigue before a healthy heart does. In other words, the heart free from disease can perform the greatest task which is ever demanded of it. The work of the heart consists in discharging the blood conveyed to it, and the skeletal muscles are not capable of driving enough blood along the years to tax its powers. On the other hand, a person with a diseased heart, should he indulge in strenuous exercise, runs a serious risk, if not of inducing cardiac failure, of at least causing serious damage to his heart. In apparently

healthy persons after middle age, the state of the heart muscle is an unknown quantity, and for this reason excessive muscular effort should be avoided

## The Central Nervous System

The structure of nervous tissue. The brain and spinal cord are composed of nerve cells with their processes—the nerve fibers—and neuroglial cells. The bodies of the nerve cells, and their processes for a short distance from their origin, compose the gray matter of the central nervous system. The nerve fibers, collected into bundles or tracts, constitute the white matter. Neuroglial cells perform no essential nervous function; they form a supporting framework for the nervous elements proper. There are three types of neuroglial cells, called, respectively, oligodendroglia (or oligodendrocytes), astrocytes, and microglia (see Fig 12.1). The last mentioned are elements of the reticulo-endothelial system

The nerve cell or neuron is the structural unit of the nervous system It consists of a body and one or more processes. The cell bodies in different parts of the nervous system vary widely in size (4 to 25 microns) and in shape (triangular, multiangular, round, spindle- or pear-shaped) In respect to their internal structure, however, all possess certain characteristics in common. The typical nerve cell has a relatively large globular nucleus situated near the center of the cell and containing, as a rule, a single nucleolus. The cytoplasm, or pericaryon—that is, the part of the body of the cell exclusive of the nucleus—shows, after suitable staining procedures, fine fibrils—the neurofibrils. These course through the cell body from pole to pole, sweeping around the nucleus. Their presence in the nerve fiber has been mentioned on page 461. When stained with an appropriate basic dye, the cytoplasm of the nerve cell exhibits a striated or mottled appearance, due to the 504.




Fig 12.1 Types of nerve and neuroglia cells. A, motor cell from anterior horn of the spinal cord, B, unipolar nerve cell; C, giant pyramidal cell from the motor area of the cerebral cortex; D, Purkinje cell from the cerebellar cortex; E, oligodendroglia cell. F, astrocyte of neuroglia; G, microglia.

presence of irregularly shaped particles, resembling the chromatic material of the nucleus. These particles are called Nissi bodies or (from their tendency to be arranged in rows, thus suggesting a tiger's stripes) tigroid bodies or the tigroid substance (see Fig. 12 2).

In 1949 Dr. Murray Barr made the astonishing discovery that the nerve cells of females could be distinguished histologically from those of males. The nucleolus in the nerve cells of the former shows adjacent to it a minute body composed of chromatin. This "nuclear satellite,"

508




Fig 12 2 A drawing (semidiagrammatic) of a nerve cell All but one dendrite have been cut

as it was originally called is not found except very rarely in the nerve cells of males. The dot of chromatin has since been shown to be present in the cells of the other tissues of the body in which it lies near the nucleolar membrane it is now more usually referred to as the sex chromatin (see p 666)

A nerve cell possesses as a rule many processes which vary very





Fig. 12 3 Diagram to show how a few nerve cells may communicate with a great number of others in different parts of the central nervous system

widely in length, thickness, and general appearance, according to the type of cell. These processes are of two kinds called, respectively, dendrites and axons. The dendrites which are usually multiple, though they may be single or even absent, conduct impulses toward the body of the cell; the axon conducts from the cell body. A typical dendrite has many branches (Gk. dendron, a tree), whereas the axon, which usually arises from the opposite pole of the cell is, as a rule, a long slender fiber with no branches near the cell body; it branches near its termination, or it may give off twigs at intervals along its length and more or less at right angles to it. The axon is always single.

Fig. 12 4 End feet (represented by small black ovals) or pieds termineaux making contact with the body and the dendrites of a nerve cell; A, axon; D, dendrites.



The processes of nerve cells constitute the fibers—both motor and sensory—making up the white matter of the brain and spinal cord and, gathered into bundles, compose the peripheral nerve trunks. Thus some processes, such as those extending from the lower part of the spinal cord to the toes, are 3 or 4 feet long in man. Structurally, there is nothing to distinguish these two types of long nerve fiber from one another.

Neurons are linked together in the central nervous system axon to dendrite or to the cell body. Through the formation in this way of chains of two or more links long and often very intricate conducting paths are established in the nervous system (Fig. 12.3). A junction be tween the axon of one neuron with a dendrite or the cell body of an other is called a synapse.—There is no actual structural union of two neurons at the synapse—merely contact. The axon terminates in a small swelling called a synaptic nob end foot, or pied termineau through which it makes contact with the dendrite or body of another nerve cell (Fig. 12.4).

## Reflex Action

510

A nervous reflex is an involuntary act brought about by the stimulation of an afferent nerve ending or receptor (p. 514). Familiar examples of actions which are purely reflex in nature are the following the quick closure of the eyelid when some object touches the eyelashes or even suddenly approaches and threatens to strike the eye the sudden




Fig 12 5 Diagram of the simplest conceivable reflex are

withdrawal of the hand or foot when it is painfully stimulated, the sharp recovery of one's balance when, as a result of a slip the body's center of gravity is suddenly shifted and the jerk of the leg (knee jerk p 518) when the patellar tendon is tapped sharply Though such reflex acts are brought about involuntarily, we are aware that they have occurred, but innumerable reflex acts of which we are quite uncon

scious—e g, the movements of the gastrointestinal tract, variations in heart rate and respiration, changes in caliber of the small blood vessels, the secretion of glands, etc—are continually taking place in the body

The reflex arc. The structural basis of reflex action is the reflex arc. The latter may be described in its simplest conceivable form as consisting of two neurons, linked together in the central nervous system, axon to dendrite (see Fig 125) The two neurons are thus arranged, one with its axon, the other with its dendrite, directed toward the periphery The latter receives the stimulus, and transmits impulses to the central nervous system where, through a synaptic contact, the two nerve cells are linked, it is therefore called the receptor neuron, or the afferent limb of the reflex arc. The other neuron, which transmits impulses




Fig 12.6 Diagram of a spinal reflex are with its center in the spinal cord showing receptor and effector neurons connected by the connector neuron (in the spinal cord)

from the central nervous system to the peripheral organ, muscle, or gland, is called the effector neuron or the efferent limb of the reflex are. The region in the central nervous system (brain or spinal cord) where the two neurons form their junction (synapse) is referred to as the reflex center. Thus a stimulus applied to the terminal of the receptor neuron sets up an impulse or a volley of impulses which is transmitted to the reflex center, a discharge of impulses then occurs down the effector neuron to the peripheral organ or effector (e.g., muscle or gland)

For the sake of simplicity in illustrating the general principles of the reflex arc, one composed of only two neurons has been described, but it is unlikely that more than a very few such simple reflex arcs exist in higher animals. One nerve cell, at least, is, as a rule, interposed between the receptor and effector neurons. This is called the connector, internuncial, or intercalated neuron. In the great majority of reflex arcs in the central nervous system of mammals, not one but a chain of such connector neurons is present.


3 One of these is that of the knee perk (p 518)

512

A diagram of a simple reflex are having its center in the spinal cord is shown in Figure 12.6. It will be noted that the cell body of the receptor neuron is situated, not within the cord itself but just outside—namely, in the ganglion of the posterior root of a spinal nerve. The effector neuron has its cell body in the anterior horn of gray matter (anterior or ventral horn cell, p. 524); its axon leaves the cord in the anterior root of the spinal nerve, and is continued into a peripheral nerve. The connector neuron has its cell body in the posterior (dorsal) horn of spinal gray matter; its axon synapses with the dendrite of the anterior horn cell; its dendrite with the central process (axon) of the recentor neuron

In most reflex actions of higher animals, a great number of reflex arcs are involved. For example, should a person's arm be given a sharp and unexpected slap, he would jerk the arm away; associated movements of the shoulder and trunk would probably occur; he would turn his head and eyes toward the source of the injury and, most likely, utter an exclamation of some sort. A large number of muscles-those of the arm, shoulders, trunk, neck, eyes, tongue, larynx, and respiration-would therefore take part in the action Each muscle is supplied in turn by a large number of nerve fibers-effector (motor) neuronsand the stimulus must have excited a large number of afferent nerve endings—receptor neurons—in the skin. If instead of slapping the skin, which stimulates a relatively large area, a stab were made with a pin, the reflex response would probably be the same. A movement involving a large number of muscles would result Yet, as compared with the slap, the pin must have stimulated much fewer sensory endings. The number of effector (motor) neurons must have been greatly in excess of the sensory endings which had been stimulated. Each receptor neuron must therefore be connected within the central nervous system with a large number of effector neurons As a matter of fact, every receptor neuron is potentially in communication with motor neurons throughout the entire spinal cord and brain. The synapses, normally, offer a certain "resistance" to the passage of impulses which limits the spread of excitation within the central nervous system. In strychnine poisoning or in tetanus, however, synaptic resistance is greatly diminished; a very weak localized stimulus then sets up impulses which spread widely throughout the central nervous system, and may cause forcible convulsive contractions of virtually all the voluntary muscles of the body.

Not only is each afferent fiber in communication with a large number of effector neurons, but, conversely, each of the latter is connected with many afferent fibers. That is to say, stimulation of sensory nerves in widely separated parts of the body may bring about reflex contraction of one and the same group of muscles. An anterior horn cell of the cord is, therefore, a point of convergence of a number of afferent paths, each motor neuron serving as a pathway common to impulses from various sources (see Fig. 12.7). It is therefore called the final



common path. For example, scratching the back of a spinal dog (i.e., one whose spinal cord has been severed in the upper thoracic region some weeks previously) causes rhythmical contractions of the flexor muscles of the hind limb (scratch reflex). Painful stimulation of the hind paw also causes contraction of the flexor muscles of the limb, the paw being thus withdrawn from the injurious agent (flexor reflex). If one should attempt to elicit both these reflexes at the same time, there must obviously be a conflict between them. They are antagonistic and cannot both employ the final common path at the same instant. In such and other instances of two antagonistic reflexes competing for the final common path, the one of less biological importance gives way; it is suppressed. Thus, as compared with a painful stimulus which signals some injury to the animal, the momentary irritation of a flea is of little importance. It is otherwise with reflexes which are allied in na-

514

ture; they can utilize the final common path simultaneously; one reflex tends to strengthen or reinforce the other. For example, stimulation of the toes of the hind paw causes flexion of the corresponding limb. If the opposite forepaw is stimulated simultaneously, the flexion of the hind limb is strengthened.

Receptors. Most types of afferent nerve terminate at the periphery in specialized structures, known as receptors or sense organs. There are various types of receptor, each being especially adapted to respond to a particular kind of stimulus. Those of the skin, for example, respond to mechanical or to thermal stimuli, the rods and cones of the retina to light, the taste buds and olfactory cells to chemical stimuli, and those of the ear to sound vibrations. The energy of the stimulating agent is converted by the receptor, in each instance, into nerve impulses.

Receptors fall into three main classes—namely, exteroceptors, proprioceptors, and interoceptors. Exteroceptors are those which receive stimuli from the external environment. The receptors of the eye, ear, and skin, and those of taste and smell—i.e., those of the five senses come under this heading. The receptors of sight, hearing, and smell are sometimes also called distance receptors or telereceptors, since they make possible perception at a distance (see Pl. 9A).

Proprioceptors receive stimuli which arise within the body itself. The skeletal muscles, the tendons and joints, the lungs, heart, and abdominal viscera, as well as certain arteries (carotid and aorta, see p. 199) and veins, contain receptors which respond to changes in the activity of their immediate surroundings (internal environment). Most receptors in the viscera give rise to no sensation and serve reflex actions of which we are quite unaware. The proprioceptors situated in the skeletal muscles are stimulated by the contraction or stretching of the muscle fibers; these, called muscle spindles, have been described (p. 483) Of the impulses set up by the stimulation of proprioceptors in muscles, tendons, and joints, some reach consciousness; to such impulses is due our sense of the movement and position of our limbs—kinesthetic sense. Many impulses set up by muscular movement end, however, in parts of the central nervous system (e.g., spinal cord and cerebellum) which carry out their activities beneath the level of consciousness. The tonic contractions of skeletal muscle, as well as the smooth and coordinated nature of voluntary movement, are dependent upon impulses, both sensory and nonsensory, arising in the muscle proprioceptors.

The interoceptors are situated in the mucous membrane of the re-

spiratory and alimentary tracts. Though removed from direct influence by the external environment, these receptors, nevertheless, are stimulated by agencies originating in the outside world—namely, air and food.

Conduction over the reflex arc. Reflex conduction differs from conduction over nerve fibers in several particulars. In the first place, the velocity of transmission of impulses over the reflex arc is considerably lower than that over the nerve fiber. For example, if a muscle is caused to contract reflexly, the time elapsing between the application of the stimulus to the receptor and the response of the muscle is considerably greater than if the impulses had traveled the same distance over a nerve fiber alone. The difference is due to the synapse, where the impulses are delayed. Second, conduction in the reflex arc is always in one direction-to the reflex center in the afferent limb, away from the center in the efferent limb. When a nerve fiber is stimulated, impulses travel in both directions. If, for example, a motor nerve fiber is cut just after it leaves the spinal cord, and the central end stimulated, impulses are set up which travel along the efferent fibers to the spinal center. But these impulses, caused by the artificial stimulus, do not reach the afferent neurons; they are blocked at the synapses. The latter thus acts like a valve, permitting impulses to pass in one direction over the reflex arc, but not in the other. (See also p. 521.)

Third, reflex conduction is much more susceptible to lack of oxygen, Jatigue, and to the action of anesthetics. We have seen that the nerve fiber has a very low rate of metabolism (p. 478), and will continue to respond to stimulation for a considerable time when deprived of oxygen. The metabolism of nerve cells, on the other hand, is high, and the cells very soon lose their irritability when deprived of oxygen. The cells of the cerebral cortex, for example, are permanently injured if their blood supply is cut off for 5 minutes or so The lower levels of the central nervous system are somewhat less susceptible to lack of oxygen: the cells of the midbrain continue to function for 10 minutes after their blood supply has been arrested; the centers of the medulla oblongata for from 15 to 30 minutes; and those of the spinal cord for from 15 to 35 minutes.

In the presence of oxygen, the nerve fiber continues to respond to stimulation for long periods without showing fatigue, in marked contrast to the reflex arc, which is readily fatigued. In the intact animal, general anesthesia (e.g., by ether or chloroform) quickly abolishes reflex action, whereas little effect is produced upon the excitability of the nerve fiber. Even after an animal has been killed by an anesthetic

(death being due to paralysis, by the anesthetic, of vital centers in the brain—e.g., of heart and respiration), the nerves for some time later show little or no alteration in their responses to direct stimulation of course, when an anesthetic is applied to an isolated nerve, its excitability is very quickly abolished. Among other characteristic features of reflex action are summation, after discharge, and reciprocal inhibition

Summation It was pointed out on page 476 that the magnitude of the nerve impulse cannot be increased by increasing the rate of stimu lation. When a nerve receives stimuli in rapid succession, the impulses do not fuse and produce an impulse of greater magnitude, but always remain separate. Summation, on the other hand, is readily demon strated in the reflex are. A single stimulus for example, applied to an afferent nerve, though it sets up an impulse, rarely causes a reflex re sponse. If, however, two or more stimuli of the same strength as the single one are applied in rapid sequence, a response follows. Each impulse produces a state of some sort in the region of the synapse which is capable of being added to or summed with, the state produced by previous impulses. The nature of this state produced at the synapses—the central excitatory state as it has been called—is not clearly under stood but evidently it must be raised to a certain threshold value be fore excitation of the effector neurons can occur.

After discharge When a motor nerve is stimulated, the response of the muscle ends (i e, impulses cease to be discharged along the fibers of the nerve) the moment that the stimulus is withdrawn When, on the other hand, the muscular response is brought about reflexly, the muscle may continue to contract for a short time after the stimulus to the afferent nerve has ceased This reflex phenomenon is explained upon the basis that the central excitatory state, which has been built up at the synapse, takes an appreciable length of time to be discharged along the effector neuron and, therefore, persists after the stimulus has been withdrawn

Reciprocal inhibition Stimulation of a motor nerve results in the contraction of all the muscles supplied by the nerve, irrespective of their actions, muscles having antagonistic actions (e.g., flexors and extensors) contract and oppose one another. No purposeful movement of the part is brought about, the motor nerve does not contain inhibitory fibers. In a reflex action, on the contrary, the muscular response consists of a coordinate movement—flexion of the knee joint, for example (see flexion reflex, below). In such a movement, the flexors of the knee contract while the extensors undergo reciprocal inhibition (relaxation). The mechanism whereby extension or other antagonistic

action is inhibited has been much debated, but the inhibitory effect must be developed in the reflex center—that is, at a synapse between the afferent and efferent neurons—for, as just mentioned, the motor nerve itself does not contain inhibitory fibers. We therefore speak of a central inhibitory state as well as of a central excitatory state.





Fig 12 8 Illustrating reciprocal inhibition

Reciprocal inhibition is also seen in voluntary movement. For example, in flexing the arm at the elbow, the contraction of the biceps is accompanied by reciprocal inhibition of the triceps. On the other hand, when the elbow is extended, the triceps contracts while the biceps relaxes (Figs. 12.8 and 12.9). Such reciprocal muscular action is, of course, essential for the performance of any purposeful act.

Certain limb reflexes of special interest. The flexion reflex. This reflex consists of a strong contraction of the flexor muscles of the limb (together with inhibition of the extensors—reciprocal inhibition) when the afferent nerve of the same (ipsilateral) limb is stimulated. The limb is flexed at ankle, knee, and hip. The biological significance of this reflex is evident, in the intact animal the limb is withdrawn, auto



518

Fig 12.9 Reciprocal innerva ton Record from the leg mus cles of a decerebrate cat show ing contraction of fixors I7, and inhibition of the extensors E The two curves actually were inscribed synchronously but the lever for the extensor muscle is set a little to the right of that of the fixor muscle (From Sherrington)

matically, from an agent which causes pain and, therefore, threatens injury. This and other reactions which are initiated by injury, and serve a protective purpose, are called nociceptive (L. noceo I injure) reflexes.

The crossed extensor reflex This is a strong contraction of the extensors of the limb which results from stimu lating the corresponding afferent nerve of the opposite (contralateral) limb It is evident then, that this and the preceding reflex can be brought about simultaneously by the stimula tion of a single afferent nerve. In the everyday life of the animal, the two reflexes therefore compose a purpose ful act for, should one paw come in contact with something which stimu lates it painfully, it is raised from the ground, while the body's support is strengthened by extension of the op posite limb

The stretch reflex Stretching a muscle connected through its nerves with the spinal cord causes it to con tract. The response is purely reflex Proprioceptors in the muscle—the muscle spindles—are excited by a stretch stimulus. The impulses thus

set up are transmitted to the spinal reflex center, a discharge of impulses then occurring down the effector neurons. A stretching force that lengthens the muscle by a small fraction of an inch is sufficient to elicit this reflex, its importance in maintaining a posture or attitude of the body for a prolonged period will be spoken of presently

The knee jerk and other tendon (or deep) reflexes A light tap upon the tendon below the kneecap (patellar tendon), while the knee hangs

limply in a semiflexed position, causes a quick contraction of the extensor muscle on the front of the thigh (quadriceps muscle). The leg gives a sharp kick. This is called the knee jerk or patellar reflex; it is one type of stretch reflex, the tap upon the tendon causing a sudden stretch of the thigh muscle. In the investigation of a person with nervous disease, the knee jerk is tested as a matter of routine. The reflex is abolished in any condition which interrupts the reflex arc-namely, disease or injury of the peripheral nerves (efferent or afferent fibers) supplying the quadriceps muscle, or of the reflex centers in the lumbar region of the spinal cord. It is exaggerated in-disease or injury involving the corticospinal tracts descending from the cerebral cortex to the cells of the anterior horns of the spinal cord (p. 526). Other tendon reflexes of clinical importance are the biceps and triceps jerks (tapping the biceps tendon in front of or the triceps tendon above the elbow) and the ankle jerk (tapping the tendo Achillis-the tendon above the heel). Loss of the biceps and triceps jerks indicates disease or injury of the nerves of the arm or of the motor cells in the thoracic region of the spinal cord, exaggeration of these reflexes is seen in disease involving the fiber tracts connecting the cerebral cortex with the motor cells in the thoracic part of the spinal cord; abolition of the ankle jerk tells of disease or injury involving the nerves to the leg muscles, or of the nerve cells of the lumbosacral part of the spinal cord; exaggera-

Though they do not come under the present heading, it is convenient to menicin here two cutaneous or superficial reflexes of great clinical importance—namely, the plantar response and the abdominal reflex. The normal plantar response consists of a downward movement (plantar-flexion) of the great toe—i.e., toward the sole, when the skin of the sole is stroked. In certain nervous lesions (e.g., injury to the corticospinal tracts), the great toe instead of being flexed toward the sole moves

tion indicates a lesion of the motor tracts

of the cord at his level.




Fig 12.10 Illustrating the normal plantar response, upper sketch, and the extensor response (Babinsky), lower sketch.

upward (dorsiflexion). This abnormal response is called the sign of Babinski (Fig. 12.10).

The abdominal reflex consists in a contraction of the abdominal muscles caused by stroking the overlying skin. It is abolished in lesions

of the corticospinal tracts, of the peripheral nerves, or of the reflex centers in the thoracic part of the spinal cord

The axon reflex This is not a reflex at all, in the ordinary sense of the term, for nerve cells play no part, that is to say, no reflex center is interposed in the path of the impulses. The reflex depends upon the fact that impulses can pass in either direction along a nerve fiber, in the axon reflex, they travel from some or other receptive surface along one branch of the fiber (axon), and then down the other branch to an effector—e g, smooth muscle or gland cell (see Fig. 12 11) In the



520

Fig 12 11 Showing the structural basis of the axon reflex Note that no nerve cell is involved. The arrows indicate the course taken by the nerve impulse

lowest forms of multicellular animals, such as mollusks, worms, and in certain parts of insects, these reflexes play a prominent part in nervous responses. In higher animals, they have been studied most often in the conjunctiva, mucous membrane of the intestinal tract, and in the skin.

The following well known and simple experiment furnishes proof of two way conduction in nerve The frog muscle shown in Figure 12 12 is supplied by a nerve which divides into two branches, each of which goes to different parts of the muscle. When the muscle is divided between the two branches, stimulation of one branch causes contraction of both parts of the muscle. Impulses must have passed along the stimulated branch and, turning at the Y formed by the two branches, reached the part of the muscle innervated by the unstimulated branch.

In man cutaneous vasodilatation caused by stimulation of the skin is brought about largely through axon reflexes. One branch of an axon of a sensory nerve ends in the skin the other in or near the wall of a small blood vessel. Vasodilatation is brought about, apparently, not by

the nerve impulse directly, but by a chemical substance which it causes to be liberated from the end of the axon branch. The vascular reactions caused by irritation of the cornea, and the increased vascularity of inflamed tissues, are also largely due to axon reflexes. Such reflexes can be abolished in any part by blocking conduction in the sensory terminals (e.g., in skin or conjunctiva) as by the application of cocaine.

Skeletal muscle tone. During consciousness the voluntary muscles,

Skeletal muscle tone. During consciousness the voluntary muscles, though not engaged in any movement, are always maintained in a state of slight contraction which is referred to as tone or tonus. All voluntary




Fig. 12.12 Illustrating two-way conduction in nerve.

muscles show tonus to some extent, but it is seen pre-eminently in those muscles which maintain the posture of the body against the force of gravity—the so-called antigravity muscles. These are, chiefly, the extensor muscles of the lower limbs, trunk, and neck. Muscle tonus is entirely reflex in character. The muscles are stretched between their attackments, thus the constant stimulus of stretch applied to the muscle spindles initiates the reflex. Like any other reflex, it is abolished if the reflex are is interrupted at any point, such as by a lesion of the motor nerve, of the afferent nerve, or of the reflex centers. For example, in injury or degenerative disease of the motor nerves, in locomotor ataxia (which involves the afferent paths in the spinal cord), or in anterior poliomyelitis (commonly known as infantile paralysis, in which the anterior horn cells are attacked), the tone of those muscles implicated by the disease is completely lost.

Higher centers of the brain (their precise location is unknown)

exert an inhibitory influence upon muscle tone. Thus, in man a lesion of the corticospinal tracts, which connect the cerebral cortex with the spinal centers, results in an exaggeration of the tonic contraction, this is manifested by the greater resistance offered by the limbs to passive movement (that is, a movement of the patient's limb made by the examiner). The physician, in describing the muscles whose tone is increased in this way, says they are spastic, and refers to the hyper tonic state itself as spasticity. Muscle tone is increased to an extreme degree after complete separation of the lower centers from higher



Fig 12 13 Decerebrate rigidity (After Pollock and Davis)

control. For example, section through the brain of an animal, anywhere between the upper part of the midbrain and the ves tibular nuclei (p 528) in the medulla oblongata results in pronounced rigidity of those muscles that support the body or parts of the body (eg, head) against gravity (antigravity muscles) All four limbs are stiffly

extended, the head is held erect, and the tail elevated. The stiffence limbs will support the unconscious animal when it is placed upon its feet. The condition is called decerebrate rigidity (Fig. 12.13). Thypertonus is at once abolished by destruction of the vestibular nucle or section of the spinal cord, or of the posterior spinal nerve toots, for this latter operation interrupts the afferent limb of the reflex are

Destruction of the vestibular nuclei or section of the spinal cord of an animal, whether or not it has been previously decrebrated, also abolishes the normal tone of the muscles The muscles become completely relaxed (flacead) immediately after this operation. The limbs hang limply like those of an animal immediately after death and no reflex activity can be elected. Also, since the vasomotor centers in the medulla oblongata are separated from the spinal centers, there is a profound fall in blood pressure (p. 196). This state, called spinal shock, persists for a time which varies with the animal species. Recovery from spinal shock occurs within a few minutes in the frog but not for several weeks in the dog, and never completely in main. From these observations it is concluded that the spinal reflex path.

From these observations it is concluded that the spinal reflex path ways, upon which muscle tone is directly based, must be reinforced by impulses discharged from the vestibular nuclei. The latter, in turn are under an inhibitory influence from higher centers, especially in the





Fig 12.14 Diagram of the principal factors responsible for muscle tone 1, cerebral cortex or other higher cere bral region 2 vestibular nucleus 3 spinal cord, 4, muscle spindle, 5 anterior horn cell of the spinal cord

midbrain and cerebellum. The vestibular nuclei, when released by section from the restraint of higher levels become hyperactive, with consequent exaggeration of tonus and the production of the state known as decerebrate rigidity described on page 522. Though the spinal centers are dependent, normally, upon impulses from the vestibular nuclei for the maintenance of tonus they can, in animals at any rate, act independently when isolated from the higher influence. This is evident from the fact that spinal shock is recovered from after a time (see diagram, Fig. 12.14)

## The Spinal Cord

The spinal cord is composed of a central, deeply and irregularly fluted column of gray matter, surrounded by white substance. The latter consists of bundles of nerve fibers (see Fig. 12.15). In a cross section of the spinal cord the gray matter appears as a central mass shaped roughly like the letter H, or the wings of a butterfly. The anterior and broader limbs of the H are called the anterior or ventral horns, the posterior narrow limbs, the posterior or dorsal horns Running through the center of the spinal gray matter is a canal filled with cerebrospinal fluid. This is the central canal of the cord, above, in the medulla, it expands to form the floor of the fourth ventricle of the brain. The spinal cord is incompletely divided into two lateral halves.

by an anterior cleft and a posterior septum. The white matter of each half is marked off again by the anterior and posterior horns of gray matter and the corresponding spinal nerve roots into an anterior, a lateral, and a posterior column. In the thoracic and upper lumbar regions of the cord, the gray matter between the anterior and posterior horns shows a small projection called the lateral horn, it is composed of sympathetic nerve cells.




LATERAL Fig. 12.15 A secCOLUMN ton of the spinal
ord showing the
central mass of gray
MATERIOR
WE ROOT umns of white matter (ascending and
descending fiber
tracts).

The anterior or ventral horns of gray matter contain large multangular cells. These cells are points of convergence for impulses from various sources—from the periphery of the body along sensory nerves, and from the motor area of the cerebral cortex by the cortico-spinal tracts (p. 526), as well as from the cerebellum and other parts of the brain. The axons of the anterior horn cells of the cord constitute the only pathway (final common path, p. 513) from the spinal cord to the skeletal muscles; the anterior horn cells, therefore, relay the impulses which bring about voluntary movement, and are also the effector (motor) neurons of the spinal reflex arc. The posterior horns of gray matter contain the cell bodies of connector neurons; afferent impulses, after entering the cord, are relayed by these cells to the anterior horn cells, and to neurons which connect with higher levels of the central nervous system.

The spinal nerve roots. In man there are 31 pairs of spinal nerves. Each nerve of a pair arises from the cord by two roots, an anierior or ventral, and a posterior or dorsal. Each root is formed by the fusion of a series of filaments (rootlets) that emerge from the vertical grows bounding the lateral columns of the cord. The anterior roots are composed of efferent (mostly motor) fibers derived from the cells of the

anterior horns of spinal gray matter; other finer fibers are efferents of sympathetic nerve cells in the lateral horns. The posterior roots are composed almost entirely of afferent fibers: sensory fibers from the skin, muscles, viscera, etc., and other afferent fibers which transmit nonsensory impulses—that is, impulses which do not reach consciousness. The cell bodies which give rise to the fibers of a posterior root are situated in a small swelling on the root itself; this is called the ganglion of the posterior root (Figs. 12.16 and 12 17). Both dendrite and axon of a ganglion cell arise by a common stem which divides into the respective fibers a short distance from the cell body. One branch (den-

Fig. 12 16 Dagram to illustrate Walleran degeneration (modified from Halliburton). Section at A causes degeneration of all motor and sensory fibers of the peripheral nerve beyond the point of section (black area). In B, section of the anterior root causes degeneration of the motor fibers since the motor cell bodies lie within the spinal cord; in C, section of the posterior root is followed by degeneration of the sensory fibers of the peripheral nerve because the cell bodies of the sensory fibers lie in the ganglion, in D, section fibers lie in the ganglion, in D, section



of the posterior roof between the ganglion and the cord causes degeneration of the sensory fibers of the spinal stump of the root and of ascending fibers within the cord as far as the next cell station (See also p. 463.)

drite) of the T-shaped process passes peripherally, the other (axon) carries impulses into the cord where it connects (synapses) with intraspinal neurons or passes without interruption to the medulla oblongata. Certain cranial nerves possess similar ganglia from which their afferent fibers originate (see Fig. 12.32, p 544). The two roots of a spinal nerve unite within the intervertebral foramen. The trunk so formed divides again almost immediately into a large anterior and a small posterior part, called, respectively, the anterior and posterior primary divisions of the spinal nerves. Each division receives fibers (motor and sensory) from both spinal nerve roots. The branches of the anterior primary divisions fuse with one another, redivide, and continue again in a complicated fashion to form three nerve plexusesthe cervical, brachial, and lumbosacral plexuses-from which the peripheral nerves to the skin and muscles of the neck and limbs ultimately emerge. Thus, in the formation of the plexuses, the fibers of the original spinal nerve roots become intermingled, the peripheral nerve

526

trunks containing both motor and sensory fibers. The posterior prunary divisions break up into branches which supply the skin and muscles of the back.

The fracts of the spinal cord. The nerve fibers composing the spinal white matter have various origins and form longitudinal bundles, usually referred to as tracts or fasciculi. The fibers of some of these tracts arise from higher levels of the nervous system and conduct impulses




Fig 12.17 The spinal nerve roots

downward; others arise from lower levels and conduct in an upward direction. They are therefore classed into ascending and descending groups. Each tract is further given a specific name derived from the origin and destination of its fibers. A cross section of the cord showing the several tracts will be found in Plate OB

The descending tracts The corticospinal or pyramidal tracts arise from the large cells of the motor area of the cerebral cortex (p. 532) on each side of the brain. The fibers of each tract synapse with the anterior horn cells of the coposite side of the cord. The tracts descend through the substance of the brain,

ternal capsule of the cerebrum (p. 539), the midbrain, pons, and medulla oblongata They form two prominences on the anterior aspect of the medulla oblongata, called the pyramids. At the lower border of the medulla oblongata the greater proportion of the fibers of one side cross to the opposite side of the cord and descend in the lateral column. They constitute, on each side of the cord, the lateral cartico-pinnal tracts. The remaining uncrossed fibers—the anterior corticospinal racts—descend in the anterior columns, but these also ultimately cross to the anterior horn cells of the opposite side at various levels of the cord.

The corticospinal tracts transmit impulses which bring about voluntary movements of the limbs and trunk. A lesion involving these tracts in any part of their course, from the motor area of the central cortex to their terminations in the spinal cord, results in weakness or paralysis of the muscles on the opposite side of the body. The fibers conveying impulses to the muscles of the eyes, face, tongue, and throat also rise from the motor area of the cerebral cortex, but they form connections with nerve cells in the motor nuclei of the cranial nerves (third to the twelfth, p. 547). These fibers, though they are strictly analogous to the corticospinal fibers and travel with the latter for a part of their course.

#### Table 12-1

Effects of lesions of the upper motor neuron

(1) Paralysis of the spastic type. That is, the paralyzed muscles are hypertonic and offer a greater resistance to passive move-

- ments than normally. (2) The tendon reflexes (knee jerk,
- etc.) are exaggerated.
- stimulation. (4) Little muscular wasting.
- (5) Babinski response (p. 519); abdominal reflexes lost (p. 519).

Effects of lessons of the lower motor neuron

The paralyzed muscles are flaccid. They are hypotonic, offering little resistance to passive movement.

The tendon reflexes are absent,

(3) Normal responses to electrical 'Reaction of degeneration (p. 463).

Marked wasting of muscles.

Normal plantar response (1 e., Babinski absent); abdominal reflexes may be lost.

are referred to as the corticobulbar tracts. Like the corticospinal tracts, they convey impulses from one side of the brain to the muscles of the opposite side,

Every pathway for the transmission of impulses governing voluntary movements, whether of the limbs and trunk or of the face, eyes, tongue, and throat, consists, therefore, of two neuron links. That neuron which has its cell body in the cerebral cortex is called the upper motor neuron. The lower motor neuron is the anterior horn cell of the spinal cord or a cell in one or other of the cranial nuclei. The axons of the lower motor neurons form, of course, the peripheral motor nerves (cranial and spinal).

The effects following a lesion of the upper motor neuron differ in certain respects from those resulting from injury or disease of the lower. These differences are summarized in Table 12-1.

The rubrospinal tracts. These descend in the lateral columns of the cord to synapse with the anterior horn cells. They arise from the red nucleus in the midbrain (p. 545). After issuing from the red nucleus, the fibers cross to the opposite side (decussation of Forel). Thus the red nucleus of one side is connected with the anterior horn cells of the opposite side of the spinal cord.

The vestibulospinal tracts. These arise from the vestibular nuclei situated in the lower part of the medulla oblongata. The fibers of each tract connect with the anterior horn cells of the opposite side of the spinal cord These tracts relay impulses transmitted to the vestibular nuclei from the labyrinth (of the internal ear) and cerebellum. They are of essential importance in the maintenance of equilibrium, correlating the tone and movements of the muscles with the position of the head in space.

The chief descending tracts of the brain and cord are represented in the diagram. Plates 10B and 10A.

The ascending tracts. The fasciculus gracilis (tract of Goll) and faciculus cuneatus (tract of Burdach) occupy the posterior column of each half of the cord. They are composed of the axons of cells of the posterior root ganglia (see page 525) The fibers of these tracts transmit impulses of light touch and the sense of position and movement (kinesthetic sense) to the nucleus gracilis and nucleus cuneatus in the medulla oblongata; from here the impulses are relayed upward by other neurons to the optic thalamus, and thence to the cerebral cortex. When these tracts are injured, the subject, since he is not informed of the movements and positions of his limbs, has difficulty in executing muscular acts with orderliness and precision; his movements are jerky and poorly controlled Impaired muscular control of this type is called ataxia. The sense of touch is little affected, however, because touch impulses ascend also in the anterior spinothalamic tracts

The spinocerebellar tracts. These are two in number on each side. One, the dorsal spinocerebellar tract, reaches the cerebellum via the inferior cerebellar peduncle (p. 552); the other, the ventral spinocerebellar tract, ascends to the midbrain, entering the cerebellum through its superior peduncle. They ascend in the lateral columns of the cord and transmit nonsensory impulses from the muscles to the cerebellum. Injury or disease of these tracts also results in ataxia, because the cerebellum does not receive the impulses that enable it to exercise its function in steadying and strengthening voluntary muscular acts

The spinothalamic tracts ascend one in the anterior column, the other in the lateral column of each half of the cord. The anterior spino-

529

thalamic tract conducts impulses of light touch from the skin of the opposite side of the body to the thalamis, from the thalamis the impulses are relayed by other fibers to the postcentral gyrus of the cere bral cortex. The lateral spinothalamic tract is the pathway for impulses of pain, heat, and cold from the opposite side of the body (see Pl 10B).

The sensory pathways in the brain are described on page 547

# The Brain or Encephalon

The brain is that part of the nervous system enclosed by the skull It consists of the cerebrum, midbrain pons, medulla oblongata and cerebellum (see Fig. 12.18)

The cerebrum The cerebrum is divided incompletely by a longitudinal fissure (the superior longitudinal fissure) into two halves called the cerebral hemispheres. These are the large ovoid masses lying in contact with the vault and walls of the skull. The hemispheres constitute by far the largest part of the human brain, and of the brains of




Fig 12.18 Plan in outline of the brain as seen from the right side. The parts are represented as separated from one another considerably more than is natural so as to show their connections. A tegmentum B pedunctle

apes and monkeys, but are less prominent structures in the brains of lower vertebrates. The most highly developed functions of the nervous system—memory intelligence, moral sense, etc—and the centers for sight, hearing smell, taste, general body sensations, and the voluntary control of bodily movements are soated in the cerebral hemispheres.

The hemispheres are composed of a covering of gray matter, called the *cerebral cortex* and a central mass of white matter. The latter is formed of tracts of nerve fibers ascending to the cortex and descending




Fig 12 19 Diagram to show the general course of the association fibers of the right cerebral hemisphere. The stippled area is the corpus callosum which is composed of commissural fibers.

from the cortex to lower levels of the nervous system, these are re ferred to as projection fibers. There are also many fibers which arise and end within a hemisphere itself, they connect different parts of the cerebral cortex and are called association fibers (Fig. 12.19). Some of these tracts are short and connect neighboring gyri, others are long and run between the cortex of two lobes (e.g., the frontal and occipital). Through a third type of fiber, called commissional, the two hemispheres are connected across the mid line. These fibers form a felted mass of white matter lying at the bottom of the superior longitudinal fissure, known as the corrus callosum (Firs. 12.20 and 12.25 p. 537).

known as the corpus callosum (Figs 12 20 and 12 25 p 537)

The cerebral cortex. The cortex shows numerous infoldings which give the surface of the hemispheres an appearance not unlike the surface of the hernel of a walnut These folds are called convolutions or gpr the depression or furrow between two convolutions is called a

sulcus, or if especially deep and long, a fissure By such reduplications of the cerebral surface, the total cortical area is greatly increased About a third of the gray matter occupies the surface of the comolutions, the remainder lines the sulci and fissures

On the basis of the characters of their constituent cells, several welldefined layers can be distinguished in the gray matter of the cerebral cortex. The respective cell layers, however, do not show uniform char-





Fig 12 20 Vertical section through the brain passing between the cerebral hemispheres

acters throughout all cortical regions. On the contrary, certain layers exhibit special characteristics in certain areas—e.g., the motor area in the frontal lobe and the visual area of the occipital lobe. Upon such characteristics depend, undoubtedly, the special functions of the different parts of the cortex (Fig. 12.21)

A band of cortex running downwards and forwards, in front of an parallel to the fissure of Rolando, contains giant pyramidal nerve cel (Betz cells) The axons of these cells descend through the brain—in ternal capsule, midbrain pons, and medulla—to connect with the



532

Fig 12.21 The microscopical structure of the cerebral cortex The numerals refer to the different layers. The cellular structure of each layer is shown on the left the fiber structure on the right (After Economo).

motor cranial nuclei and the an terior horn cells of the spins cord These fibers constitute the corticobulbar and corticospini tracts (p 526) They transmi impulses governing voluntar movements This part of the cor tex is therefore called the moto area But the motor area is no confined to the lateral surface of the cerebrum. It turns over the upper border and descends on the medical aspect of the hemi sphere Commencing about half way down the medial surface, and running upward to the up per border and then turning downward on the lateral surface (see Figs 12 22 and 12 23) the voluntary movements of the various parts of the body are rep resented in the following order toes ankle (up medial surface), knee (at upper border), hir trunk, shoulder, elbow, wris hand, face (down lateral sur face) The disproportionatel large areas for the hand an

meant to indicate the relatively large cortical areas representing these parts that carry out the most complicated movements and, therefore involve the greatest numbers of neurons

It has been mentioned elsewhere that the corticospinal fibers ultimately cross to the opposite side of the spinal cord. It follows that the muscles of one side of the body are controlled by the motor area of the opposite side of the brain Injury to the motor area of one hemisphere or of the corticospinal fibers in their course through the brain above the vel of their crossing, especially in the internal capsule (p. 539), erefore, results in paralysis of the muscles on the opposite side of the ody.

The areas of the motor cortex have been explored and mapped out hen exposed in man during intracranial operations. Electrical stimution of circumscribed areas elicits motor responses from groups of uscles or, if the stimulus is sufficiently localized, of individual muses on the opposite side of the body.




Fig 12 22 Diagram illustrating the localization of function in the cerebral cortex 1, motor area, 2, sensory (somesthetic) area, 3, premotor area, 4, auditory area, 5, part of visual area concerned with visual association, S, fissure of Sylvius

Situated in front of the motor area is the so-called premotor area. The functions of this cortical area have not been fully elucidated, but appears to exert a controlling influence over the motor area itself, irobably synthesizing the more localized movements represented in he latter area into more complex acts. Movements of small groups of nuscles or even of a single muscle can be elicited by stimulation of the notor area; but excitation of the premotor area causes broader and nore or less purposeful well-coordinated movements. The premotor area is also connected by descending fibers with the corpus striatum p. 539) and, through the frontopontine tract, with the gray matter of he pons. Through the former connection, it is in indirect communication with the red nucleus, cerebellum, and spinal centers; through the rontopontine fibers, a connection is also established between the premotor area and the cerebellum.




Fig 12 23 Ver tical sections through the fron tal lobe of the cerebrum at the fissure of Rolando [central sulcus) showing the motor (above) and sensory rep-resentations The disproportionate sizes of parts of the manikin are meant to indicate the relative extent of the corresponding cortical areas (After Pen held and Ras mussen }

Concerning the functions of the remaining and greater part of the frontal cortex-that is, of the part not occupied by but lying in front of the motor and premotor areas-little is definitely known; this region is therefore spoken of as a "silent" area. It is sometimes the site of a tumor, and the surgeon in order to eradicate the growth is often forced to sacrifice a large part of the brain tissue. The frontal lobe of one side, except for the motor and premotor areas, has been excised in a number of instances, but such operations are followed by little or no functional defect. Even after the removal of both frontal lobes in front of the premotor areas, there is remarkably little mental disability. It has been thought that the frontal lobe was the seat of intelligence; the greater development of this part of the human brain, as compared with the brains of lower animals, has supported such an idea. However, in the light of the results just cited and of experiments upon animals, our views on the importance of this part of the brain must be revised. There is apparently no special region or center for intelligence. It depends upon the cortex as a whole; upon the degree of development of the various cortical sensory areas, and the richness of the association paths through which these areas are interconnected. The frontal cortex is simply a region of high associative ability, and as such contributes its share to the general intellectual capacity of the individual.

The parietal lobe. The area of cortex lying behind the fissure of Rolando is sensory in function. It is called the somesthetic area. Here sensations of touch, warmth, and coolness, and of muscular movements (kinesthetic sensations), are perceived and interpreted (see also p. 528). But pain does not appear to be represented in the cerebral cortex. Impulses giving rise to this sensation end in the optic thalamus. The sensory representation of the various parts of the body show definite localization, following the same order as that already described for the motor area (see Fig. 12.23); those sensory regions—especially the hands, lips, and tongue—upon which we depend most for our impressions of the world around us have the largest cortical representation. Taste is recorded at the lower end of this area.

Like the motor area, the somesthetic area has been mapped out by point to point electrical stimulation, through an operative opening in the skull bone. The patients were conscious and able to describe the sensations aroused.

The temporal lobe. The cortex bordering the fissure of Sylvius, together with that buried in the fissure itself, is the region for the perception of sound. Here the fibers of the auditory radiation (p. 621) terminate; it is therefore called the acoustic area or the center for hearing. The acoustic area of each side of the brain receives impulses from both ears Its unilateral destruction, therefore, does not cause deafness in either ear, though some dullness of hearing in both ears may result. The temporal cortex adjacent to the acoustic area is associative in function being concerned with the interpretation of sounds—that is with understanding the meaning and significance of the various kinds of sound (e.g. words music, etc.) and with the association of a particular sound with visual tactile or other sensations. It is therefore called the psychoauditory area.

The temporal lobe is thought to be the seat of memory, a region of the cortex from which auditory visual and other impressions stored in other regions (e.g., psychoauditory or psychovixual area) can be recalled to consciousness. This belief is based largely upon the fact that a tumor of the temporal lobe or an epileptic seizure commencing in this region of the cortex is not uncommonly associated with or preceded by reminiscences when scenes long past are brought vividly to mind

The occipital lobe. The cortical area for vision—the visual area—is situated chiefly on the inner aspect of the occipital lobe, but it also extends around the posterior extremity of the lobe to a small part of its lateral surface. Impulses from the right halves of the two retinas are transmitted to the visual area of the right hemisphere, impulses from the left halves of the retinas to the left visual area (see Pf. 13A). De struction of the visual cortex on one side of the brain, therefore, causes blindness in the corresponding halves of the retinas only (see p. 586). The remainder of the posterior part of this lobe is associative in function, upon its activity depend the recognition and interpretation of visual impressions and the integration of these with other sensations.

Fig 12.24 The electroencephalogram recorded from the occupital region of the skull Eyes closed up to O in tracing when they were opened to a broken view kept open to C, when again closed Note almost complete disappearance of the waves from O to C

Electrical activity of the brain, the electroencephalogram. It was discovered in 1929, by Berger, a German physician, that rhythmical changes in electric potential—electric currents of low potential—were occurring almost continuously in the cerebral cortex, and could be recorded in man by means of pad electrodes applied to the scalp (see Fig 12 24) They can also be picked up by needle electrodes in con

tact with the skull bone Such a record, obtained by a highly sensitive electric device (amplifier and oscillograph), and a photographic apparatus, is called an electroencephalogram; the recording instrument is known as the electroencephalograph. Two types of wave are to be





Fig 12 25 Horizontal section through the brain.

seen in the record of a normal person in the waking state but with his eyes closed; they are designated, respectively, the alpha and beta waves. The alpha waves have a frequency of about 10 per second and a voltage of around 50 microvolts. They persist when the eyes are open and the visual field is uniform, as when the subject looks at a blank wall or other surface without pattern. They disappear when the view is broken up by a pattern, such as one composed of several objects, or contain-

<sup>2</sup> A microvolt is one millionth of a volt, a millivolt is one thousandth of a volt.

ing contrasting areas of light and shade, etc. The beta waves are of higher frequency (25 to 50 per second) and of smaller amplitude—that is, of lower voltage, namely from 5 to 10 microvolts. The beta waves are recorded best from the skull overlying the motor area of the cerebral cortex, whereas the alpha waves are seen best in records from



rig 12.20 Verificat transverse section inrough the view

the occipital region—that is overlying the visual area in which they apparently arise

A third type of wave called the delta wave is rarely obtained in the electroencephalogram of a normal person while he is awake but appears during sleep. These waves have a very low frequency (1 to 5 per second) and very often, a relatively high voltage (up to 200 microvolts).

The electroencephalogram is not affected by dividing the brain be low the cerebrum (through the midbrain). The electric currents are not due, therefore, to afferent impulses ascending to the cerebrum from lower levels but appear to be a manifestation of an inherent electrical activity of the nerve cells of the cerebrum itself.

Departures from the normal configuration of the electroencephalogram are seen in several abnormal conditions of the brain, notably, epilepsy and cerebral tumors. A study of such records is often of considerable value in diagnosis.

The optic thalamus, corpus striatum, and internal capsule. The optic thalamus is a large mass of gray matter buried in each cerebral hemisphere just above the midbrain. The third ventricle separates the optic thalami of the two sides (Figs. 12.25 and 12.26). In sections of the brain a streak of white matter known as the internal capsule (Figs. 12.25 and 12.26) is to be seen on the outer side of the thalamus and separating it from the corpus striatum. The internal capsule is composed of bundles of ascending and descending nerve fibers (e.g., corticospinal and sensory tracts). It is a bottle-neck pass which all fibers must traverse in order to reach lower levels of the nervous system, and through which all sensory fibers destined for the cortex must ascend. An injury to the internal capsule, therefore, causes extensive damage to the conducting pathways; paralysis and often sensory loss on the opposite side of the body result. Paralysis confined to one side of the body is called hemiplegia. It most commonly follows rupture of a blood vessel supplying the internal capsule, with consequent interruption of the motor and sensory paths. The immediate effect of hemorrhage into the internal capsule is loss of consciousness and paralysis of all four limbs. This condition, called cerebral apoplexy or in popular parlance "a stroke," lasts for a variable time, the subject may die without regaining consciousness. If he survives, the paralysis on one side disappears as consciousness is regained, but loss of power in the muscles of the face and limbs on the side of the body opposite to that of the hemorrhage persists. The tone of the affected muscles is increased (spasticity), the tendon reflexes are exaggerated, and the normal plantar reflex is replaced by the Babinski response (p. 519). In other words, the signs are those of an upper motor neuron lesion (p. 527). The optic thalamus and corpus striatum are intimately connected by

The optic thalamus and corpus striatum are intimately connected by nerve fibers, and both these gray masses are connected, as well, with the cerebral cortex, and with the spinal cord, cerebellium, and other parts of the nervous system. All sensory paths conveying impulses of pain, temperature, touch, and muscle sense enter the thalamus. The sensations of pain, of rough contact, and extremes of temperature—heat above 45° C and cold below 25° C—are appreciated in the thalamus itself. The thalamus is, therefore, an organ of crude consciousness. The finer sensations—i e., light touch, moderate changes in temperature (warmth and coolness), and muscle sense—are perceived by the

540

somesthetic region of the parietal cortex. The impulses subserving these sensations are relayed to the parietal cortex by fibers which arise in the thalamus and ascend in the internal capsule. The cortex of the parietal lobe brings a discriminating and critical ability to bear upon the impulses which it receives, being capable of judging the lesser gradations in stimulus intensity, and of detecting minor qualitative differences in sensation. The thalamus possesses no such discriminating capability. The somesthetic area of the cortex is believed also to send impulses to the thalamus which restrain its activity. When released




Fig. 12 27  $\,$  Model of the internal capsule, thalamus, and corpus striatum

from cortical restraint, the thalamus reacts in an "all-or-none" manner—a diffuse, ill-defined but intensely unpleasant sensation being aroused by ordinarily mildly painful stimuli, or by moderate changes in temperature. Such sensory abnormalities occur in disease of this part of the brain.

The thalamus is not a simple relay station, for it integrates the impulses before passing them on to the cortex.

The corpus striatum (striate body) is a large gray mass bent upon itself in such a way that in a horizontal section of the brain it appears as two separate gray masses—the caudate nucleus and the lentiform nucleus (Figs 12.25, 12.26, and 12.27). The latter consists of two parts: the outer and larger part is called the putamen; and the inner part, the globus pallidus. The lentiform nucleus is separated from the thalamus by the posterior limb of the internal capsule, and from the caudate nucleus by the latter's anterior limb. The striate body has numerous fiber connections with other parts of the nervous system—

The term nucleus is used to denote any circumscribed group of nerve cells or isolated mass of gray substance within the central nervous system.

e g, with the premotor area of the cerebral cortex, the thalamus and the red nucleus, and through the latter with the cerebellum and spinal centers. Of the functions of the corpus striatum little is definitely known. It is concerned in some way with the control of skeletal muscle tone, it probably also exerts a steadying influence upon muscular movements. In disease of the corpus striatum or of its connections,




Fig 12 28 Vertical section through the base of the brain in the region of the hypothalamus

marked rigidity of the muscles—flexors and extensors—is seen The resistance offered by a limb to passive movements has suggested a comparison with that of a lead pipe ('lead pipe rigidity) Tremor or jerkness of the limbs, upon attempting a voluntary movement, is also a common manifestation of disease of the corpus striatum Both these signs—rigidity and tremor—are seen in paralysis agutans, the name given to one form of disease of the striate body

The hypothalamus. This is the region at the base of the brain lying behind the optic chiasma (p 586) and beneath the floor of the third ventricle It contains several groups of nerve cells—the hypothalamic nuclei—which constitute centers controlling the sympathetic and parasympathetic functions Axons from three groups of nerve cells—the supraoptic, paraventricular, and tuberal nuclei—form a bundle of nerve fibers known as the hypothalamico-hypothyseal tract. This tract

542

enters the neural lobe of the hypophysis (pituitary body) It governs the liberation of the hormones of this part of the pituitary (Fig. 12.28) Injury or stimulation of the hypothalamus affects one or more of the fundamental processes of the body—e g, vasomotor control, tempera ture regulation, fat, carbohydrate, and water metabolism interference with the development of the sex functions as well as gastrointestinal motility. It has been reported that electrical stimulation of the hypo




Fig 12 29 Transverse section through the midbrain

thalamus in animals induces a sleeplike state. It also appears to be a center from which the reactions expressing the primitive emotions are controlled. For when in an experimental animal all restraint of the higher centers of the brain is withdrawn by removal of the entire cere bral cortex, the reactions of an enraged fighting animal are exhibited. Thus a cat which before the operation was a placid friendly animal spits and bares its claws when approached, its fur bristles and its tail becomes bushy, as when a normal cat is attacked by a dog. This state of the decorticated animal has been appropriately termed sham rage. When, on the other hand, a tract of fibers connecting the frontal lobe of the ecerebrum with the hypothalamus is severed a naturally wild and unfriendly animal becomes docile and affectionate. This latter observation has been put to practical advantage in the treatment of certain

mental states in man. Section of this connecting nerve tract is often followed by improvement of the mental condition.

The midbrain or mesencephalon. The midbrain is the short, narrow, pillarlike portion of the brain, lying immediately below the optic thalami. It is traversed by a narrow canal called the cerebral aqueduct;




Fig 12 30 The thalami and brain stem from behind

this communicates above with the third ventricle of the brain and below with the fourth ventricle. The greater part of the midbrain lies in front of the cerebral aqueduct, and consists of the two cerebral peduncles (Figs. 12.29 and 12.31). Anteriorly the latter are separated by a cleft, and appear as two stout columns which emerge from the pons below, and plunge into the substance of the cerebrum-above. Each column, which is termed the base of the peduncle, transmits the corticospinal, frontopontune, and temporopontune-tracts of the corresponding





Fig 1231 The midbrain looking from behind and from the right, cerebellum re-



medulla oblonguta

side of the brain. Posteriorly the peduncles are fused into a single structure called the tegmentum (L. tego, I cover) which contains, on each side of the mid-line, an ovoid mass of gray matter—the red nucleus—and two smaller gray masses, the nuclei of the oculomotor and trochlear nerves. This part of the mesencephalon also transmits two bundles of sensory fibers—the lateral and medial lemnisci. The former is the pathway for auditory impulses, the latter for impulses of touch, pain, muscle sense, etc. (Fig. 12.29).

The part of the midbrain lying posterior to the cerebral aqueduct is called the tectum (L. tectum, roof, from tego, I cover). It consists of two pairs of rounded eminences called, respectively, the superior and inferior colliculi (or corpora guadrigemina) (Figs. 12.30 and 12.31). The superior colliculus receives impulses from the retina and is connected with the spinal centers through a tract of descending fibers—the tectospinal tract. It is a center for visual reflexes—e g, movements of the head and eyes in response to retinal stimuli. The inferior colliculus receives auditory impulses, and serves as an auditory reflex center—e.g., the "pricking up" of an animal's ears in response to sound.

A small oval eminence is seen on the outer side of each colliculus. These are called, respectively, the medial and lateral geniculate bodies. The former, which is connected by a strand of fibers (called the inferior brachium) with the inferior colliculus, receives the bulk of the fibers comprising the auditory pathway (the lateral lemniscus; see also p. 621). The lateral geniculate body receives most of the fibers of the optic tract (p. 587) and is connected by a band of fibers—the superior brachium—with the superior colliculus (see Fig. 12.32).

The red nucleus (nucleus ruber) is connected by fiber tracts with the cerebral cortex, corpus striatum, thalamus, cerebellum, and spinal cord. Its functions are imperfectly understood, but it appears to be an integral part of the nervous mechanism controlling the execution of skilled muscular movements. The fibers which descend to the spinal centers are called the rubrospinal tracts. Upon issuing from the lower part of the red nucleus, these fibers cross with the fibers of the opposite side. This crossing is called the decussation of Forel.

The pons (pons Varolii). This part of the brain stem lies below or (in quadrupeds) behind the midbrain and above or in front of the medulla oblongata. It transmits the corticobulbar, corticospinal, and rubrospinal tracts, and the medial and lateral lemnisci. It contains the sensory nucleus of the trigeminal nerve and the nuclei of the facial and abducent nerves. It is in communication with the cerebellum

through the middle cerebellar peduncle, and with the cerebrum through the frontopontine and temporopontine tracts. The fibers of these tracts synapse with nerve cells in the substance of the pons called the pontine nuclei. The axons of cells composing these nuclei relay impulses to the cerebellum.

The medulla oblongata. The medulla oblongata is the continuation upward of the spinal cord. It is somewhat conical in shape, broadening as it ascends, until at its junction with the pons its circumference is nearly double that of its lower end. A canal, continuous with the central canal of the spinal cord, runs through the center of the lower half of the medulla, but in the upper half the posterior wall of the canal opens out to form a lozenge-shaped space called the fourth ventricle. The floor of the fourth ventricle is composed of gray matter; this represents the prolongation upward of the anterior horns of the cord which, through the opening up of the central canal, have become spread out Upon the anterior aspect of the medulia are two vertical columns called the pyramids, formed by the corticospinal (pyramidal) tracts. At the junction of the medulla with the spinal cord, the pyramid of each side divides into a smaller and a larger part. The latter crosses (decussates) with that of the opposite side and descends in the lateral column of the cord. The smaller division descends uncrossed in the anterior column of the cord (see p. 526). The medial fillets (see below) cross (sensory decussation) a little above the decussation of the pyramids (motor decussation) and ascend behind the latter. The nuclei of the eighth, ninth, tenth, eleventh, and twelfth cranial nerves are situated in the medulla; it also contains the nucleus gracilis and the nucleus cuneatus, and the so-called vital centers-cardiac, vasomotor. and respiratory The medulla oblongata is connected through the inferior cerebellar peduncle with the cerebellum (p. 553).

Summary of sensory pathways in the brain. As mentioned, on page 528, impulses reaching the nuclei gracilis and cuneatus along the fibers of the posterior columns are relayed upward by secondary neurons. The axons of the latter, after leaving the nuclei and crossing to the opposite side (sensory decussation), ascend through the medulla, pons, and multorain forming a tract known as the medial fillet or lemiscus. The medial fillet is joined by the anterior and lateral spinothalamic tracts, as well as by fibers carrying impulses from the face (nigeminal lemniscus). These four sets of sensory fibers lead to the optic thalamus. From here all sensations except pain and extremes of temperature (see p 539) are relayed by other neurons to the postcentral (somesthetic) area of the cerebral cortex (p. 535).

547

The cranial nerves. The nerves arising from the brain are twelve in number on each side. They are designated by numbers in the order of their origin, from before backward. Roman numerals I to XII are usually employed in referring to them, or the words first, second, third, and so on (see Fig. 12.32). Each nerve also receives a name descriptive of its function or distribution. The numbers and the corresponding names of the cranial nerves are given in Table 12-2.

#### Table 12-2

#### List of the cranial nerves

I. Olfactory VII. Facial

II. Optic VIII. Acoustic (or auditory)
III. Oculomotor IX. Glossopharyngeal

IV Trochlear X. Vagus
V. Trigeminal XI. Accessory
VI. Abducent XII Hypoglossal

The olfactory nerves, or nerves of smell, are distributed to the olfactory mucous membrane covering the roof of the nose. They will be described in the section on the sense of smell (p. 641).

The optic nerve, or nerve of sight, is composed of the axons of the ganglion cells of the retina. The central connections of the fibers of the optic nerve will be described in the section on vision (p. 587).

The oculomotor nerve arises from a group of nerve cells—the nucleus of the oculomotor nerve—situated in the gray matter of the floor of the cerebral aqueduct (p. 543). The fibers course forward through the midbrain, emerging from the inner aspect of the cerebral peduncle. The oculomotor nerve supplies all the eye muscles with the exception of the superior oblique and external rectus, which are innervated, respectively, by the trochlear and abducent nerves. It also conveys parasympathetic fibers to the constrictor muscle of the iris, and to the ciliary muscle.

The trochlear nerve has its nucleus in the floor of the cerebral aqueduct, a little behind the oculomotor nucleus. It emerges from the brain at the lateral border of the pons. The trochlear nerve supplies the superior oblique muscle of the eyeball.

The trigeminal nerve contains both sensory and motor fibers. Its motor fibers are the axons of a group of cells situated in the upper part of the pons. This collection of gray matter is called the motor nucleus of the trigeminal. The sensory fibers originate in the trigeminal (semi-lunar) ganglion, which lies upon the floor of the skull and is homolo-

gous with the posterior root ganglia of the spinal nerves. Like those composing the ganglia of the posterior spinal nerve roots, the cells of the trigemnal ganglion are unipolar—i.e., each gives off a single process that divides into two. One branch passes centrally, the other peripherally. The central fibers form a short trunk—the sensory root of the trigemnal nerve—which enters the brain in close association with the motor root. Within the brain, the fibers of the sensory root divide into ascending and descending groups. The former end in a collection of gray matter—the superior sensory nucleus of the trigemial—situated in the pons close to the motor nucleus. These fibers convey the discriminative qualities of sensation—namely, light touch, localization, knesthetic sense, etc —from the face. The descending fibers terminate in the spinal nucleus of the trigeminal nerve—an elongated mass of gray substance, extending from the lower part of the pons to the upper part of the spinal cord. The fibers entering the spinal nucleus transmit impulses of the crude forms of sensation from the face—namely, pain and extremes of temperature.

namely, pain and extremes or temperature.

The peripheral processes of the cells of the trigeminal ganglion form three trunks called the first, second, and third divisions of the trigeminal nerve or the ophthalmic, maxillary, and mandibular nerves, respectively. The first two divisions (ophthalmic and maxillary nerves) are composed entirely of sensory fibers, but the third division (mandibular nerve) is united with the motor root of the nerve. The first division, or ophthalmic nerve, supplies the skin of the forehead and anterior part of the scalp, the structures within the eye socket (orbit), and the skin of the side of the nose and upper lid. The second division, or maxillary nerve, is distributed to a part of the dura mater, certain structures within the orbit, the lower eyelid, the upper lip, and to the upper teeth. The third division, or mandibular nerve, contains both sensory and motor fibers; it sends sensory fibers to the lower teeth and the lower lip, to the skin of the side and lower part of the face, and to the mucous membrane over the anterior two thirds of the tongue and the floor of the mouth. Its motor fibers supply the muscles of mastication (see Fig 12.33).

The trigeminal nerve may be the seat of a most severe and stubborn form of neuralgia. All three branches may be involved, or the pain may be confined to the area of distribution of one branch. Surgical division of the sensory root of the nerve, or of one or other branch, may be required in order to relieve the pain.

The abducent nerve. The abducent nucleus is situated in the lower part of the pons beneath the floor of the fourth ventricle. The nerve

emerges from the anterior aspect of the brain in the groove lying between the pons and the upper end of the pyramid of the medulla oblongata. It supplies the external rectus muscle of the eyeball.

The facial nerve has a large motor and a small sensory root. The fibers forming the motor root arise from the motor nucleus, which lies in the lower part of the pons. This root also transmits secretory and vasodilator (parasympathetic) fibers to the submaxillary and sublingual glands; these fibers arise from a separate group of nerve cells—

the superior salivatory (or salivary) nucleus-lying in close proximity to the motor nucleus. The motor root of the nerve leaves the anterior aspect of the brain at the lower border of the pons. The fibers of the sensory root (also called the nervus intermedius) are the axons of cells situated in the facial (geniculate) ganglion which lies within a canal in the temporal bone. The sensory root enters the brain in close association with the motor root. The peripheral processes of the ganglion cells are distributed to the anterior two thirds of the tongue (see Fig. 13.61, p 640). They transmit impulses of taste to a nucleus in the medulla oblongata named the tractus solitarius. From the latter the



Fig. 12 33 Distribution of the trigeminal nerve in the face. Ophthalmic division, stippled, manilary division, hatched, mandibular division, black.

impulses are relayed to the thalamus and thence by tertiary neurons to the center for taste in the cerebral cortex (p. 535). Both taste and parasympathetic fibers (secretory and vasodilator) leave the facial trunk in its chorda tympani branch.

The motor fibers of the facial nerve form a stout trunk which leaves the cranial cavity through a small opening in the floor of the skull. The nerve curves forward below the ear in the substance of the parotid gland to reach the face, where it breaks up into numerous branches. These supply the muscles of the face, lips, eyelids, forehead, and anterior part of the scalp.

The acoustic or auditory nerve is entirely sensory in function. It consists of two distinct sets of fibers, the cochlear and vestibular nerves. The courses of these nerves and their central connections are described on pages 621 and 632, respectively.

The glossopharyngeal nerve contains motor, sensory, secretory, and

vasodilator fibers. The motor fibers issue from the upper part of the nucleus ambiguus situated in the medulla oblongata, and are distributed to a single muscle (the stylopharyngeus). The sensory fibers con duct impulses of taste from the posterior third of the tongue to the lower part of the tractus solitarius. The taste impulses are relayed to the optic thalamus, and thence to the cerebral center for taste in the lower part of the somesthetic area. Sensory fibers are also distributed to the mucous membranes of the pharynx, tonsil, and palate, they convey impulses of ordinary sensation, touch, temperature, etc. The secretory and vasodilator (parasympathetic) fibers leave the glossopharyn geal nerve in its tympanic branch and are distributed to the parotid gland. The secretory and vasodilator fibers have their origins in the inferior salivatory (salivary) nucleus which lies below the superior salivatory (salivary) nucleus (p. 296). The glossopharyngeal nerve also supplies a sensory filament—the sinus nerve, page 199—to the carotid sinus.

The vagus nerve contains motor, sensory, secretory, and vasodilator fibers. The motor fibers are of two types—voluntary and autonomic. The fibers to voluntary muscle originate in the lower part of the nu cleus ambiguus (see accessory nerve p 551). These fibers are, actu ally, derived from the cranial root of the accessory (ninth cranial) nerve, they enter the vagus nerve and supply the cricothyroid muscle of the larynx and the inferior constructor muscle of the pharynx. The autonomic motor fibers are distributed to the muscle of the bronchi heart, esophagus stomach gall bladder, pancreas, small intestine, and first third or so of the large intestine. The fibers going to the heart are inhibitory in function (p 187). The involuntary motor fibers the se cretory fibers (to the gastric glands and pancreas), the cardiac inhibitory fibers, and the vasodilator fibers belong to the parasympathetic division of the autonomic system. They arise from the dorsal nucleus of the vagus. This nucleus is a mixed one, for it is also a terminal for sensory fibers. Some of the latter connect with motor autonomic neurons within the dorsal nucleus which thus functions as a reflex center.

The vagus nerve emerges from the lateral aspect of the medulla ob longata as a series of rootlets which soon join to form a stout trunk. This leaves the skull through the jugular foramen, and passes down the neck in close relation to the internal and common carotid arteries. Two ganglia are situated upon the vagus trunk, the upper one lies within the jugular foramen at the base of the skull, the other just below the point where the nerve issues from the skull. The sensory fibers of the vagus have their origin within these ganglia.

The accessory nerve is entirely motor in function, and consists of a cranial and a spinal root. The cranial fibers arise from a nucleus lying below, and continuous with the lower end of the nucleus ambiguus. They leave the side of the medulla oblongata as four or five delicate strands which, after uniting, pass from the skull through the jugular foramen. The spinal fibers are the axons of anterior horn cells situated in the upper five cervical segments of the spinal cord. This part of the nerve enters the cranial cavity through the foramen magnum and reenters the neck through the jugular foramen. Within the latter fora-




Fig 12 34 The superior aspect of the cerebellum

men, it joins the cranial part, but becomes separate again almost immediately.

Most of the muscles of the pharynx, larynx, and soft palate are supplied by the cranial part of the accessory nerve. The fibers pass into the pharyngeal and laryngeal branches of the vagus, through which they are distributed to voluntary muscles in these structures (cricothyroid and pharyngeal constrictors). The spunal part sends fibers to certain muscles of the neck and shoulder (sternomastoid and trapezius).

The hypoglossal nerve is distributed entirely to the muscles (intrinsic and extrinsic) of the tongue. Its fibers arise from the hypoglossal nucleus in the medulla oblongata, and appear as a series of rootlets in the groove situated on the anterolateral aspect of the pyramid The rootlets leave the skull through the hypoglossal canal and unite just below the base of the skull The trunk so formed curves forward in the upper part of the neck to reach the cavity of the mouth

The cerebellum. The cerebellum is situated behind the brain stem (midbrain, pons, and medulla), and beneath the posterior portions of

the cerebral hemispheres (Fig. 12.18, p. 529). It is composed of two lateral masses, the cerebellar hemispheres (Figs. 12.34 and 12.35) and a central elongated wormlike structure, called the verms (L. worm). The white matter of the cerebellum forms a central branching framework within the substance of each hemisphere and in the center of the vermis. The gray matter covering the terminal stems of this structure constitutes the cerebellar cortex. The cortex of the hemispheres is thus constructed of a number of Jeaves flattened against one




Fig 12 35 The inferior aspect of the cerebellum

another with only their edges appearing on the surface. The cerebellar cortex is composed of several layers of cells, but unlike the cerebral cortex it is of uniform cellular structure—i e, the cells at all depths are of similar type. Large cells with flask-shaped bodies and extensive dendritic arborizations—the cells of Purkinge—are peculiar to the cerebellar cortex (Figs. 12.36 and 12.1, p. 507). Three masses of gray matter—the cerebellar nuclei—are to be found in the depth of the cerebellum. The largest of these is known as the dentate nucleus. The nerve fibers streaming to and from the cortex and entering or leaving the cerebellar nuclei constitute the white matter of the cerebellum.

Each half of the cerebellum is connected with the rest of the central nervous system through three compact bundles of nerve fibers, called the cerebellar peduncle of brachild confirmation of the cerebellar peduncle (or brachild conjunctivum) plunges into the midbrain and transmits im

pulses from the cerebellum to the red nucleus and thalamus and, through the latter, to the motor area of the cerebral cortex (see Figs. 12.37 and 12.38). It also transmits, in the reverse direction, the ventral spinocerebellar tract which conveys impulses to the cerebellum from the spinal cord (p. 528). The impulses that leave the cerebellum by its superior peduncle originate in the cerebellar cortex. They pass first to one or other of the cerebellar nuclei, from which they are relayed by secondary neurons. The dentate nucleus contributes most of the fibers to the superior peduncle. The other two cerebellar nuclei are






Fig 12 36 A section through the cerebellar cortex and adjacent part of the white matter

called, respectively, the nucleus emboliforms and the nucleus fastigii. The latter receives impulses from the organ of equilibrium in the internal ear via the vestibular nuclei of the medulla oblongata.

The middle peduncle (or brachium pontis) transmits impulses from the pons to the cerebellum, its fibers arise from the pontine nuclei which, as stated elsewhere, receive impulses from the frontal and temporal regions of the cerebral cortex (frontopontine and temporopontine tracts). No fibers leave the cerebellum by this peduncle.

The inferior peduncle (or restiform body) connects the cerebellum with the lower part of the medulla oblongata. It is composed, mainly, of fibers of the dorsal (direct) spinocerebellar tract and of fibers arising in the vestibular nuclei in the medulla oblongata (vestibulacerebellar tract). Thus, through the inferior cerebellar peduncle, the cerebellum receives proprioceptive impulses from the labyrinth (p. 630) and the skeletal muscles. The inferior peduncle also transmits impulses from the cerebellum to the vestibular nuclei (cerebellovestibular tract).





The functions of the cerebellum are concerned with voluntary movements, yet they play no part apparently in initiating such movements. An animal deprived of its cerebellum shows no actual paralysis, but its movements are shaky, jerky, and poorly controlled. In man, injury or disease of the cerebellum causes the same lack of muscular control; movements are not executed smoothly and evenly, and with the nicety of direction and force characteristic of normal muscular action. For example, the subject of cerebellar deficiency, if asked to touch his nose with his finger, moves the arm jerkly and fauls to hit the mark. The gait is staggering in character; the muscles are hypotonic, and the limbs show a coarse tremor upon attempting any movement. Ataxia is the term applied in general to disorders of voluntary muscular control; those resulting from disease of the cerebellum are therefore grouped under the designation cerebellar ataxa.

As a result of numerous experiments upon animals and of studies of cerebellar deficiency in man (gunshot wounds or disease), it is now generally believed that the chief function of the cerebellum is to blend or synergize the actions of the different muscles engaged in a given movement. Through cerebellar activity, the contractions of the individual muscles, or groups of muscles, are so timed and graded in force that their combined action results in a smooth and effective movement. The anatomical basis for such function has been outlined above. Its superior peduncles connect the cerebellum with the motor area of the cerebral cortex on the one hand and, on the other hand, through the red nucleus and the rubrospinal tracts, with the anterior horn cells of the spinal cord. Through the spinocerebellar tracts, on the other hand, the cerebellum is kept constantly "informed" of the position of the limbs and movements of the muscles Through the vestibulocerebellar tracts, it receives impulses from the labyrinth acquainting it with movements of the head in space.

The impulses reaching the cerebellum give rise to no sensation; the cerebellum carries on its activities entirely beneath the level of consciousness.

# The Membranes (Meninges) of the Brain and Spinal Cord

The central nervous system is enclosed within three concentric membranous sheaths. The outermost of these is relatively thick and tough and is therefore called the *dura mater* (L. hard mother). The inner-

most is very delicate, and is applied closely to the surfaces of the brain and spinal cord. It is known as the pia mater (L. tender mother). The membrane between these two has been named the arachnoid mater, or simply, the arachnoid (Fig. 12.39). Its name is derived from




Fig 12 39 The membranes (meninges) of the brain Upper drawing, a vertical transverse section, label of dura mater omitted. Lower drawing, a vertical longitudinal section to show the partitions (septa) of the dura mater.

the meshwork of filaments that connect it with the underlying pia mater, and which somewhat resembles a spider's web (Gk. arachnē, a spider).

The dura mater consists of two layers, the outer of which serves as a lining (periosteum) for the skull bone, the inner one as a protective covering for the neural structures; it carries the larger blood vessels. The dura mater gives off a sickle-shaped process, called the fats cerebri, which descends vertically between the two hemispheres; a hori-

zontal sheet, the tentorium cerebelli, which projects forward between the cerebrum and the cerebellum; and a short vertical crescentic spur, the falx cerebelli, that extends forward in the notch between the posterior poles of the cerebellar hemispheres. The narrow interval between the dura mater and the arachnoid is called the subdural space. The arachnoid, a much thinner membrane than the dura mater, forms a loose investment for the central nervous system. With one exception—namely, the deep fissure between the hemispheres—it does not dip into the fissures and sulci, and other depressions in the contours of the brain, but bridges over them The pia mater, on the contrary, adheres to the underlying nervous tissue, following all its irregularities faithfully, even into the depths of the fissures.


The subarachnoid space and the ventricular system. The interval between the arachnoid and the pia mater which, as mentioned above, is interlaced by a delicate network, is known as the subarachnoid space. It is filled with a thin Impid liquid called the cerebrospinal fluid. This fluid is similar in composition to plasma or lymph, but its dissolved constituents (sodium chloride and bicarbonate excepted) are in lower concentration. Its composition is given below. It will be observed that the protein concentration is extremely low, whereas that of sodium chloride is higher than in plasma (see Table 12-3, and pages 57 and 58).

Table 12-3

### Composition of cerebrospinal fluid in milligrams per 00 cc

Protein	15-55
Glucose	50-80
Calcium	4-6
Potassium	12-17
Sodium chloride	720-750
Bicarbonate	
(Volumes percent CO <sub>2</sub> )	40-60
Inorganic phosphate	1.25-2.0

Very few cells are to be found in the cerebrospinal fluid, and they are of the lymphocyte variety. The cerebrospinal fluid is under pressure. In the cisterna magna, in the lying down position, it is around 90 mm of water, but falls to zero or a little below when the subject stands. In the lower part of the spinal subarachnoid space (lumbar region), the cerebrospinal fluid pressure in the lying-down position is about the same as that in the cisterna magna, but rises to between 500






Fig 12 41 Diagram of the ventricles of the brain

and 600 mm of water in the standing position. This is about the hydrostatic pressure of a column of fluid extending from the cranial cavity to the lumbar region. Coughing, sneezing, blowing, straining at stool, or lifting weights raises the pressure in the veins of the thorax. The augmented pressure is transmitted through the veins of the neck, and thence to the cerebrospinal fluid. The volume of fluid is kept constant at about 130 ce (in man) by the nicely balanced processes of formation (secretion by the choroid plexus) and reabsorption through the arachnoid granulations (see below). There is no communication between the subarachnoid space and the subdural space; the latter, like the pleural and peritoneal cavities, is a potential space only, a thin layer of lymph lying between the two membranes.

The cerebrospinal fluid serves a protective function, serving to shelter the delicate nervous structures from mechanical shocks that may be applied to its bony encasement. It gives buoyancy to the brain and spinal cord; the weight of the brain in its normal situation is only about one thirtieth of its weight after removal from the cranium.

Since the pia mater clings to the surface of the brain, dipping into all the fissures and sulci and following closely all the contours of the cerebrum, brain stem, and cerebellum, whereas the arachnoid forms a relatively loose investment, there are certain regions-such as at the junction of the pons with the medulla, and of the latter with the cerebellum-where the subarachnoid space is widened. In such regions, there are relatively large collections of cerebrospinal fluid. These are called cisternae (L. a cistern or reservoir). The principal cisternae are shown in Fig. 12.40.

Hollow spaces filled with cerebrospinal fluid are to be found within the brain itself. They are call




Fig. 12.42 Cast of the ventricular system of the brain. (Redrawn after Retzius.)

within the brain itself. They are called ventricles, and are four in number. The largest are the lateral ventricles, one in each cerebral hemisphere (Figs. 12.41 and 12.42). The third ventricle is a much smaller space situated between the optic thalami. The fourth ventricle lies between the cerebellum behind and the pons and upper half of the me-

dulla oblongata in front. The posterior aspect of the latter two parts of the brain stem really form its anterior wall which, however, is usually referred to as its "floor". The superior cerebellar peduncles and parts of the cerebellum enter into the formation of its posterior wall or roof. Highly vascular fringes of pia mater project into each of the four ven tricular cavities. They are covered with epithelium which secrets the cerebrospinal fluid and are known as the choroid plexises. All the ventricles of the brain communicate with one another, as well as with



Fig 12 43 Internal hydrocephalus in a young child (Drawn from a photograph after Purves Stewart) The shaded area at the dome of the head represents the part of the skull (anterior fontanelle) which owing to the wide separation of the bones, is covered by scalp and membrane only In a normal child of the same age, this fontanelle should be completely, or very nearly, closed and even at birth it is no more than about I inch in its transverse dameter.

the subarachnoid space and the central canal of the spinal cord. The third ventricle communicates with the lateral ventricles through the foramina of Monro, and the third ventricle with the fourth through the cerebral aqueduct. Through an aperture in a recess on each side, the fourth ventricle communicates with the subarachnoid space and, at its inferior angle, it is continuous with the central canal of the spinal cord.

Arachnoid granulations is the name that has been given to small excressences of the subarachnoid mesh which, carrying a covering of dura mater before them, project into the venous sinuses of the brain (Fig. 12.39). They filter fluid from the subarachnoid space into the blood stream A smaller amount of fluid passes into the capillaries of the pia mater. Thus, by this process of reabsorption, the cerebrospinal fluid is maintained at constant volume. Any block, such as in the cere bral aqueduct or at the foramina of Munro (which may occur as a consential defect, or from disease in infancy or, though less commonly, in later life) will upset the balance between secretion and reabsorption.

Fluid will continue to be formed by the choroid plexuses on one side of the block, but reabsorption of the secreted fluid by the arachnoid granulations will be prevented. Damming back of fluid results, and the part of the ventricular system involved (this includes the third ventricle and usually the lateral ventricles) becomes greatly dilated. As a consequence of the increased fluid pressure, the brain substance becomes thinned and the skull bones separated. There is usually mental deterioration. This condition is known as internal hydrocephalus (Fig. 12.43).

External hydrocephalus may be caused by a blockage at some point in the subarachnoid space, as may result from meningitis, tumor, etc. The cerebrospinal fluid is then increased in any part of the subarachnoid space cut off from the arachnoid granulations. Cisternae, so isolated, are enlarged and the ventricular system may be expanded. There is a pronounced rise in intracranial pressure. But this type of hydrocephalus may occur though no blockage exists, and may then be due to an increase in the production of cerebrospinal fluid by the choroid plexuses (as in meningitis) or to an elevated pressure in the venous sinuses of the brain.

The withdrawal of cerebrospinal fluid 'from the subarachnoid space in the lumbar region through a specially designed wide-bore needle introduced between the fourth and fifth lumbar vertebrae (lumbar puncture) is today a common and invaluable procedure in the diagnosis of meningitis, brain abscess, tumor, etc. Lumbar puncture is also employed for the introduction of a spinal anesthetic or of a curative drug. The chief abnormaluties in the cerebrospinal fluid that are of diagnostic importance in neurological disease are a cloudy appearance, an increase in the protein content (ten times or more), and a rise in the number of leucocytes.

#### The Autonomic Nervous System

The autonomic (Gk. autos, self + nomos, law, i.e., self governing) or involuntary nervous system governs those functions which are carried out automatically, and which under ordinary circumstances do not obtrude upon consciousness. These functions include the control of the rate of the heart, the movements of the gastrointestinal tract, the caliber of the small blood vessels, the contraction or inhibition of smooth muscle in various other structures (e.g., skin, urinary bladder, gall

<sup>4</sup> A sample of cerebrospinal fluid may also be obtained in a comparable manner from the cisterna magna (cisternal puncture) or from a lateral ventricle.

bladder, and bronchi) and the secretion of the sweat glands and of various digestive glands.

The efferent nerves of the autonomic nervous system arise from groups of cells situated at different levels of the central nervous system, from the midbrain to the sacral region of the spinal cord. This fibers, after issuing from the brain or cord, make connections with nerve cells situated either in a ganglion or in the innervated organ itself (muscle or gland). Thus, every autonomic pathway consists of two neurons. The axon, whose cell body lies within the central nervous system, is called the preganglionic fiber; that of the outlying (ganglionic) nerve cell, the postganglionic fiber (see Pl. 12A).

The autonomic nervous system is divided upon an anatomical as well as upon a physiological basis into two parts, called the parasympathetic and sympathetic divisions (see Pl. 11).

The parasympathetic division. The parasympathetic division is subdivided into a cranial and a sacral part or outflow. The preganglionic fibers of the cranial outflow arise from groups of cells situated in the midbrain, pons, and medulla oblongata The midbrain fibers originate in the oculomotor nucleus and connect with cells in the ciliary ganglion; postganglionic fibers are distributed to the iris, and transmit impulses which constrict the pupil The cells in the pons are in close association with the nucleus of the facial nerve; the preganglionic fibers are conveyed in the chorda tympani branch of the facial nerve to ganglion cells situated in, or in close relation to, the submaxillary and sublingual (salivary) glands From here the impulses are transmitted by postganglionic fibers to the gland cells. The fibers in the chorda tympani nerve are vasodilator as well as secretory in function.

In the medulla, the cells that give rise to the parasympathetic fibers are situated in the glossopharyngeal and vagus nuclei. The former enter the glossopharyngeal nerve and pass to the otic ganglion; from here gostpanglionic fibers conveying vasodilator and secretory impulses are relayed to the parotid gland.

The vagus nerve is composed in the main of parasympathetic fibers; these have a very wide distribution—to the heart, bronchioles, esophagus, stomach, small intestine, and first third of the large intestine, and to the pancreas, liver, gall bladder, and bile duets. The preganglionic fibers of the vagus may, therefore, be 2 feet or more in length, whereas the postganghome fibers are very short. The latter form a rich plexus—Auerbach's plexus—in the walls of the gastrointestinal tract, bronchioles, and biliary duets. Groups of (ganglionie) nerve cells are scattered throughout the plexus of fibers.

The cells of the sacral outflow are situated in the second, third, and fourth sacral segments of the spinal cord. The fibers leave the cord by the anterior spinal nerve roots. Separating again from the other anterior root fibers, they combine to form the pelvic nerve. This nerve supplies motor fibers to the distal two thirds of the large bowel and to the wall of the urinary bladder, and vasodilator fibers to the penis and clitoris. It also contains unhibitory fibers to the internal anal sphinicter and to the internal sphinicter of the bladder. The postganglionic fibers of the sacral outflow arise from small ganglia situated in close proximity to, or in the walls of, the innervated organ.

The sympathetic division. The preganglionic fibers of the sympathetic nerves are the axons of cells in the lateral gray horns of the spinal cord (p. 524), from the first thoracic to the second or third lumbar segments, inclusive. The sympathetic is, therefore, commonly referred to as the thoracicolumbar outflow of the autonomic nervous system. The preganglionic fibers leave the cord by the anterior roots of the corresponding spinal nerves. The ganglia of the sympathetic are in two main groups, the vertebral and the prevertebral. The vertebral ganglia are situated on either side of the vertebral column (Pl. 12A). They appear as a series of 22 or 23 swellings on each side, connected together to form a long, beaded cord. This extends from the base of the skull to the coccyx, and is called the gangliated cord of the sympathetic. At their lower limits the two cords come together and end in a single (unpaired) ganglion called the ganglion impar.

The prevertebral ganglia are larger than the vertebral; they lie in front of the spinal column and in close relation to the aorta and its branches.

The sympathetic fibers destined for the supply of the limbs (blood vessels, sweat glands, and smooth muscle of the skin) soon separate from the anterior nerve roots and enter the vertebral ganglia. These fibers (preganglionic) are seen issuing as slender glistening strands, one from each anterior spinal nerve root from the first thoracic to the second or third lumbar segment. They are called the white rami communicantes. The fibers of which each of these strands is composed enter the vertebral ganglion corresponding to the segment of the spinal cord from which they arise; here synapses are usually made, by at least a small proportion of the fibers, with rerve cells (Pl. 12A). But the fibers of the white rami communicantes do not necessarily terminate around nerve cells in the ganglion which they first enter; they may pass up or down the gangliated cord to synapse with ganglion cells at a higher or lower level. The axons of the cells in the ganglia (post-

ganglionic fibers) leave the ganglia as short yellowish-pink filaments, known as the gray rami communicantes, which join the spinal nerves. The sympathetic fibers are thus distributed to the periphery with the ordinary motor and sensory fibers. All the spinal nerves receive postganglionic fibers, though the anterior roots of only a proportion of the spinal nerves (first thoracie to second or third lumbar) give rise to white rami (preganglionic fibers). That is to say, the spinal nerves above the first thoracie segment or below the third lumbar have gray rami but no white rami. The sympathetic fibers above or below these levels must travel up or down the gangliated cord to the appropriate vertebral ganglion and then pass as gray rami to the spinal nerves (see Pl. 12A).

The sympathetic fibers to the viscera and blood vessels of the abdomen do not connect with the vertebral chain of ganglia, but pass to the prevertebral ganglia—celiac, superior mesenteric, etc. The postganglionic fibers form plexuses around the branches of the abdominal aorta, from which the vessels and viscera receive their sympathetic supply. The fibers passing from the spinal cord to the prevertebral ganglia are collected on each side into three well-defined strands called the greater, lesser, and least splanchnic nerves.

Functions of the autonomic nervous system. The sympathetic division exerts a regulating influence on a great number of structures. Through the cardiac accelerator nerves, it increases the rate of the heart; through the splanchnic nerves, it inhibits the movements of the intestinal tract, maintains the tone of the arterioles of the abdomen, hastens the formation of glucose from glycogen by the liver, and causes the liberation of adrenaline from the adrenal medulla. Through the part of the sympathetic in the neck—the cervical sympathetic—impulses are conveyed to the pupil (causing dilatation), to the salivary glands, and to the blood vessels of the head and neck. Sympathetic impulses cause relaxation of the wall of the urmary bladder, but contraction of the internal sphincter of this organ Sympathetic fibers in the cutaneous nerves transmit motor impulses to the smooth muscle of the skin and secretory impulses to the sweat glands.

Many structures—such as the iris, heart, intestines, urinary bladder, salivary glands, and pancreas—receive fibers from both divisions of the autonomic nervous system. The actions of the two upon a given organ are antagonistic and belanced one against the other, the activity shown by the organ at any moment being the resultant of the two opposing influences. The parasympathetic fibers (vagus) to the heart, for example, are inhibitory, the sympathetic excitatory. On the other hand,

Table 12-4

Organ	Parasympathetic effects*	Origin of sympathetic postganglionic fibers	Sympathetic effects
Heart	Inhibition	Superior middle and inferior cervical ganglia	Acceleration
Vessels Cutaneous Muscular Coronary Salivary glands Pulmonary	Constriction D latation Constriction	Various vertebral ganglia Various vertebral ganglia Cervical ganglia Superior cervical ganglion Thoracic vertebral ganglia	Constriction Dilatation Dilatation Constriction Constriction and d latation
Cerebral Abdominal and pelvic viscera	Dilatation Dilatation	Superior cervical ganglion Prevertebral ganglia	Constriction Constriction
External genitalia	Dilatation	Prevertebral ganglia	Constriction
Eye Iris Cihary muscle Smooth muscle of orbit and upper lid	Constriction Contraction	Superior cervical gangl on Superior cervical gangl on Superior cervical ganglion	Dilatation Relaxation Contraction
Bronchi	Constriction	Thoracic ganglia	Dilatation
Glands Sweat Sahvary Gastric	Secretion Secretion	Vertebral ganglia Superior cervical ganglia Celiac ganglion	Secretion Secret on Inhibition Secretion of mucus
Acini Islets Liver Adrenal medulia	Secretion Secretion	Celiac ganglion Celiac ganglion Cel ac ganglion No postganglionic fibers	Glycogenolysis Secretion
Smooth muscle Of skin Of stomach wall	Contraction or in	Vertebral ganglia Celiac ganglion	Contraction Contraction or inhibition
Of small intest ne	Increased tone and	Celiac and superior mesen	Inhibition
Of large intestine	Increased tone and	teric gangl a Inferior mesenteric and hy	Inhibition
Of bladder wall (detrusor muscle)	Contraction	pogastric ganglia Inferior mesentene and hy pogastric ganglia	Inhibition
Of trigone and	Inhibition	Inferior mesenteric and hy pogastric ganglia	Contraction
Of uterus pregnant	Nil	Inferior mesenteric and hy pogastric gangl a	Contraction
Of uterus nonpregnant	Nil	Inferior mesenteric and hy pogastric ganglia	Inhibition

With certain exceptions (e.g. those supplying the sublingual and parotid glands and the sphincier pupiliae) the postgangl one fibers of the parasympathetic arise from cells situated in, or in close proximity to the innervated organ itself

## THE CENTRAL NERVOUS SYSTEM

566

motor fibers (excitatory) to the wall of the intestine are derived from the parasympathetic the inhibitors from the sympathetic The walls of the urinary bladder and intestine receive their motor innervation from the parasympathetic (vagus and pelvic nerves) whereas the sympathetic is inhibitory. The internal sphincters of the bladder and anus are innervated in a reverse manner, they are excited by the sympa are innervated in a reverse manner they are content of the state thetic and inhibited by the parasympathetic

A summary of the actions of the parasympathetic and sympathetic

fibers upon a number of structures is given in Table 12 4

The sympathetic and the hormone of the adrenal medulla are closely similar in their actions. The two acting in conjunction constitute what is referred to as the sympathoadrenal system and play an important role in the regulation of the internal environment of the body-te the composition and temperature of the fluids bathing the cells of the the composition and temperature of the most butting the cent state. It is through its effect upon the blood vessels sweat glands and smooth muscle of the skin the sympathetic controls heat loss through its action upon the blood vessels it also varies the distribution of water between the vascular system and the tissues Through an ac of whiter petween the vascular system and the fissues. Through sin at too upon the liver either through sympathetic nerve impulses or the liberation of adrenaline the sugar of the blood is raised. The sympathoadrenal system through its various activities increases the body's efficiency in times of stress. Many manifestations of an animal when in danger or when its powers are being taxed to the utmost are those of sympathetic stimulation—e.g. dilated pupils rapid heart action contraction of the spleen and the erection of hair or rufflino of feath ers (due to contraction of cutaneous smooth muscle see also p 423)

The sympathoadrenal system highly important though it is can nevertheless be dispensed with It is not essential to life nor even to well being provided the animal is not exposed to some environmental hazard Animals from which the entire sympathetic and the medulla of both adrenals have been excised live in perfect health in the shell tered surroundings of the laboratory. They cannot however with sind cold and are less well equipped than is a normal animal to meet an emergency which demands the marshalling of its resources either to defend itself or to fly from the threatened danger

The transmission of autonomic effects by chemical substances Re search of recent years has disclosed the amazing fact that many sym pathetic and parasympathetic effects are not brought about directly by the nerve impulses themselves but by chemical substances which the impulses cause to be I berated at the nerve indings. In treating of the control of the heart at has been mentioned that when the vagus is stimulated, acetylcholine is liberated, and is the direct cause of the inhibitory effect upon the heart muscle Similarly, adrenaline (p. 420), is liberated at the terminals of the cardiac accelerators, and from certain other nerves of the sympathetic system (see pp. 188 and 425). These facts, which have led to a revolutionary change in our ideas of peripheral nervous action, have been supplemented within the last few years by a number of observations of great interest and importance

It has been shown for example, that acetylcholine is liberated at the terminals of such parasympathetic nerves as those going to the iris (in the oculomotor nerve), to the salivary glands (in the chorda tympani), to the stomach and intestines (in the vagus), and to the bladder (in the pelvic), as well as from certain sympathetic nerves—e.g., from those to the sweat glands and from those that cause dilatation of the vessels of the skeletal muscles

Acetylcholine liberation also plays a part, though its precise mode of action is not altogether clear, in the transmission of nervous effects across sympathetic and parasympathetic synapses—that is, from preganglionic fibers to the ganglionic cell. Still more extraordinary is the discovery that acetylcholine liberation in the region of the motor end plate is an essential event in the contraction of skeletal muscle.

Following the suggestion of Sir Henry Dale the English physiologist, it is now customary to speak of those fibers that liberate an adren alinelike substance (see sympathin) as adrenergic, and those that liberate acetylcholine as cholinergic. All preganglionic fibers (sympathetic or parasympathetic), all postganglionic parasympathetic fibers, and certain postganglionic sympathetic fibers (e.g., to sweat glands), as well as the fibers of voluntary motor nerves are cholinergic (see Pl. 12C).

#### **Conditioned Reflexes**

The type of reflex (unconditioned reflex) that has been considered (pp 297 and 510) is carried out entirely through centers situated in the spinal cord or lower (subcortical) levels of the brain Unconditioned reflexes are inborn, they are not dependent upon past experience, education, or training In conditioned reflexes, on the other hand, the activity of the cerebral cortex plays an essential role. Conditioned reflexes were first demonstrated and studied by Pavlov, the Russian physiologist, and most of the knowledge that we possess of this type of response is the outcome of his investigations. An understanding of




Fig 12 44 Upper drawing showing arrangement for carrying out a conditioned reflex experiment. The open ng of the parotid duet has been transplanted to the skin of the cheek so that the saliva as it is secreted may be collected and measured. The an mal is separated from the experimenter by a part tion in order that extraneous types of stimulation (e.g. movements or sounds made by the experiment.) shall not arouse the animals interest and thereby interfere with the conditioning process. (From Paylov Lectures on Conditioned Reflexer. In ternational Publishers. N.Y.) Below d agram to illustrate the theory of the mechanism underlying conditioned reflexes. I taste buds 2 salvary center. 3 salvary gland. 4 occipital (visual) cortex. S. uncond to need at mulus C.S., conditioned stimulius in A the animal is fed and a stimulus shaped to the taste buds In B a conditioned visual stimulius (e.g. a flash of 1 ght) is applied at the same time as the uncond tioned stimulius. A pathway from the occipital cortex to the salivary center is thus established. In C., the cond to ned stimulius applied alone. Impulses pass from the visual cortex to the salivary center and evoke salivary secretion.

the conditioned response can best be gained by giving a few examples of the results of Professor Pavlov's experiments

When a newborn puppy is given milk to drink there is a secretion of saliva due to the stimulation of the taste buds in the mouth. The reflex are in this instance is constituted of the nerves of taste the salivary centers in the medulla and the secretory fibers (chorda tympani or glossopharyngeal) to the salivary glands. This is a simple or unconditioned reflex It is an inherent response and quite independent of previous experience gained through any of the organs of special sense. Now, as the puppy grows older it associates the appearance or smell

of the milk, or both, with its taste and the pleasure of gratifying the appetite. The mere sight or smell of the milk will then elicit a secretion of saliva; such a response is called a conditioned reflex. Pathways have become established in the brain between the cortical center for sight or for smell and the salivary center. In other words, the visual and olfactory stimuli-conditioned stimuli-set up impulses which, impinging upon the salivary centers, are capable alone (i e., without stimulating the nerves of taste or other nerves in the mouth) of exciting them (see diagram, Fig. 12.44). If, on the other hand, a puppy which has never tasted meat is offered some (but is not allowed to taste it), there is no secretion of saliva. The association paths between the higher centers of the brain and the secretory centers have not been developed. Conditioned reflexes become established not only with respect to feeding, and the secretion of saliva, but in many other similar ways in the everyday life of the growing animal Such processes are essentially psychic in character and dependent upon experience They, therefore, form the basis of training, and are of the greatest biological importance. The animal, through the conditioning process, reacts appropriately to the various stimuli-beneficial or injurious-in the environment.

The Pavlov school has shown that conditioned responses can be developed experimentally to an extraordinary degree. For example, if an animal is fed a number of times, during or shortly after the ringing of a bell, a secretion of saliva occurs merely upon ringing the bell. The contact of food with the taste buds constitutes the unconditioned stimulus; the sound of the bell, the conditioned stimulus. The flash of a light, the sound of a buzzer, horn, or ticking metronome, a particular odor, a touch upon the skin, or the passive movement of a limb into a certain position, and many other types of stimuli may serve for the establishment of the conditioned response—that is, may act as conditioned stimuli.

Inhibition of conditioned reflexes. There are two main kinds of inhibition affecting conditioned reflexes: external and internal. External inhibition is that which causes an established conditioned response to be suppressed when the animal is distracted by something which he sees or hears, or by any event extraneous to the experiment itself. Internal inhibition is the failure of a conditioned reflex to be elicited after it has been repeated a number of times without the application of the original unconditioned stimulus For example, if a salivary conditioned response is evoked several times without giving the animal food, it becomes weaker and weaker with each repetition and finally fails

completely The response is said to have been extinguished. In order to maintain the conditioned response at full strength the unconditioned stimulus must be employed after every few repetitions. This re application of the unconditioned stimulus is called reinforcement.

Another type of internal inhibition known as conditioned inhibition is best explained by an example. Let us say that a salivary conditioned reflex has been firmly established to the sound of a tuning fork another stimulus—e.g. a flash of light—is then combined with it but whenever the two are in combination reinforcement (feeding) is omit ted. Then the single stimulus—namely the sound of the tuning fork—if regularly reinforced causes salivary secretion but if combined with the flash of light does not. Thus the latter stimulus has an inhibitory effect upon the original conditioned response.



Fig. 12-45 Examples of different figures which were successfully of fiferentiated in experiments upon a dog. The letter T shown in the upper left hand corner of the figure served for the post ive stimulus the obligate, figures and the who te letter T were differentiated from the post ve stimulus from which the other cross was the post ve stimulus from which the other which figures were different ated (From Paylov Condition of Reflegere, Oxford University Press).

The conditioning process can be so developed that stimuli closely similar in character can be differentiated (as indicated by the secretion of saliva) with astonishing precision. A dog in which a conditioned salivary reflex has been established to a sound of a certain pitch will not respond to a sound of the same quality and intensity but varying in pitch from the original by only two or three double vibrations per second.

Differentiation of visual stimuli can also be developed to a phenom enal degree Figures of various shapes—a cross square or circle—are readily differentiated (see Fig. 12.45). A circle can be differentiated by the dog from an ellipse whose diameters have a ratio no greater than 9 to 10. That is if a salivary conditioned reflex has been established to a circular object such as a white cardboard disc it alone evokes a response. A disc precisely the same in all respects except that it is not quite circular but has diameters in the ratio of 9 to 10 will not

cause a secretion of saliva Olfactory tactile and proprioceptive stimuli can likewise be differentiated with remarkable precision

It is evident that conditioned reflexes afford a valuable means for studying perception in animals. It has been established for example that the dog possesses the ability of discriminating between slight variations in the intensity quality and pitch of a musical note between various odors and by sight between objects of different sizes and shapes and between lights of different intensities. Colors on the contrary cannot be differentiated nor can a colored object be distinguished from a colorless one of the same size and shape. It is concluded therefore that the dog is totally color blind its surroundings being perceived only in white gray and black.

### The Physiology of Sleep

During sleep most of the functions of the body are carried on at the lowest levels possible in health. Heat production is from 10 to 15 per cent below the so called basal level. The mechanisms regulating the body temperature are less sensitive than in the waking state and are depressed by from 0.5 to 1.0 degree Fahrenheit, the rate of the heart is reduced in frequency by from 10 to 30 beats per minute, and a decline in blood pressure of about 20 mm occurs in quiet restful sleep. The turine volume is considerably reduced but its concentration in solids is increased. The cutaneous vessels are usually dilated and the secretion of sweat is stimulated. The tone of all the skeletal muscles is lessened the deep tendon reflexes are depressed or abolished and there may be a Babinski (extensor) response of the great toe (p. 519). The eyes are usually rolled upward and the pupils constricted. Gastric motility and secretion have a tendency to be greater during sleep than in the waking state.

Seriously detrimental effects may be exerted upon the nervous system by lack of sleep Experiments upon human subjects in which wake fulness was enforced for long periods caused profound psychological changes. In one series of experiments, the volunteers went without sleep for periods of a duration up to 112 hours. Loss of memory, irritiability hallucinations and even schizophrenic manifestations were observed within from 30 to 60 hours of sleeplessness.

The required hours of sleep depth of sleep. The amount of sleep required at different ages is given in the table below. Adults vary greatly in the hours of sleep which they find necessary to "knit up the ravelled sleeve of care."

	Hours
New born	 18-20
Growing children	10-12
Adults	6-9
Aged persons	5-7

The depth of ordinary restful sleep fluctuates throughout the sleeping period. In most adults, sleep deepens through the first hour, after which it lightens rather sharply, and then more gradually until morning or until the usual time of wakening. In growing children, however, sleep deepens a second time for a little while in the eighth or ninth hour. In these observations the depth of sleep was judged by the amount of movement of the subjects.

Theories of sleep Many theories have been advanced to explain the temporary loss of consciousness which we know as sleep Their very multiplicity emphasizes the difficulty of finding a satisfactory explanation of the physiological processes underlying the phenomenon Probably the oldest theory is that sleep is induced by a reduction in the blood supply to the brain or at least to the conscious centers. This is the so-called ischemic theory. Even the ancient Greek physicians had an idea that the carotid artery was in some way concerned with the onset of sleep. The name itself expresses this belief (Gk. karonides carotid arteries from karoo put to sleep). In modern times the drows ness after a meal (the blood being diverted presumably from the brain to the digestive organs) was pointed to in support of the ischemic theory.

Some have favored a chemical theory, in one form or another Chemicals such as lactic acid, acetylcholine, bromide, or a specific fatigue toxin' were supposed to accumulate during the waking hours which acted like a drug upon the nerve cells of the higher centers of the brain. But no sound evidence can be cited in confirmation of this theory. It would carry us too far afield to discuss the various theories, nor would it be of much profit to do so. But three theories will be briefly reviewed.

The first of these places a sleep center in the hypothalamus (p 541) A state, at least closely resembling sleep has been induced in animals by the electric stimulation of this region of the brain I is also true that many of the bodily changes in sleep are manifestations of activity of hypothalamic nuclei (especially of parasympathetic centers), such as constriction of the pupil, cutaneous vasodilation reduced frequency of the heart beat, increased gastric tone and secretion, etc.

The second theory to be outlined is that which has as its chief basis the reduction in the discharge of the afferent impulses from the periphery to the conscious centers. During waking hours, these centers are being continually 'bombarded by impulses from the organs of special sense, as well as from the skeletal muscles (impulses of kin esthetic sense) and semicircular canals. The centers are being con stantly 'alerted'. But when the sum of these impulses is reduced to a minimum, consciousness becomes dulled and sleep is induced. That muscular relaxation, and freedom from visual, auditory, and cutane ous stimulation are conducive to sleep needs no emphasis. We require a dark and quiet room, a soft bed and an even, comfortable tempera ture in order to sleep. But the reduction in the flow of afferent impulses to the brain can be no more than a contributory factor in the onset of sleep for the presence of all the desirable conditions mentioned will not always bring sleep. On the other hand sleep may supervene in the face of the most adverse conditions, in a bright light and noisy sur roundings and even on hard ground.

Payloy, as a result of his study of conditioned reflexes in dogs pro posed a theory of sleep based upon cortical inhibition (p. 569) caused by the repeated elicitation of a conditioned response without reinforce ment The conditioned inhibition thus established in one area of the cerebral cortex may spread to associated areas and finally involve the entire cortex Such widespread cortical inhibition is sleep, according to Pavlov It may be induced by various types of conditioned stimulation, if repeatedly applied without reinforcement. It is a familiar experience that a monotonous sound, an unchanging view, a boring lecture, or any recurring event for which our interest flags is conducive to drowsiness Counting sheep the least exciting of animals, is a well known device resorted to by those suffering from insomnia Pavlov described some experiments in which the animal subject would fall into a sleeplike state standing in its harness and even while it was be ing stimulated All attempts to elicit the conditioned response failed On later occasions, it would fall asleep when merely brought into the room where the previous experiments had been conducted. The well known surroundings had themselves acted as conditioned inhibitory stimuli There is little doubt but that the familiar appointments of our bedroom (we know very well that it is more difficult to go to sleep in a strange room), the preparation for bed, the feel of the bedcoverings, and even the time of retiring, act in a similar way upon ourselves. Having been associated innumerable times with sleep, they act as inhibitory conditioned stimuli, and induce in the cerebral cortex an inhibi tory influence which gradually diffuses over broad areas

All three of the theories which have been briefly sketched have in them each a degree of truth, but no one alone is entirely satisfactory

# The Special Senses

## The Physiology of Vision

General description of the eye. The adult human eye is shaped like a globe and has an approximate diameter of 1 inch (24 mm running anteroposteriorly and transversely, and 23.5 mm vertically). It is actually made up of the segments of two spheres. The anterior part is transparent and is called the cornea. It forms about one sixth of the globe and is a segment of a smaller sphere than is that of the posterior five sixths (see Fig. 13 1). The posterior wall of the globe is lined by a light-sensitive tissue called the retina A cylindrical bundle of nerve fibers—the optic nerve—passes from the posterior pole of the globe to the brain. The eye, except for the anterior fifth or so of its circumference, is enclosed in a bony case-the eye socket or orbital cavity-but a thick layer of aerolar tissue is interposed between the eyeball and the bone, which serves as a cushion to buffer it against external violence. The eyeball is also protected from injury by the eyelids which, as we know, close reflexly, in an instant, to prevent dust or other particles from coming into contact with its surface. The exposed part of the eyeball is covered by a delicate membrane called the conjunctiva which is continued forward on to the inner surfaces of the lids. When the lids open and close, the opposed conjunctival surfaces slide over one another. The surfaces are lubricated by a thin film of tears secreted by the lacrimal gland, which lies under the shelter of the bone forming the upper and outer part of the wall of the eye socket. The tears, after flowing over the surface of the eye, are drained from its inner angle into the nose by two small tubes-the lacrimal ducts (see Fig. 13.2). If it were not for the continual washing and lubrication of the eyeball by the tears, the delicate protective membrane would soon become dry

and inflamed; ulceration of the corneal surface would result. The tears also contain a material known as lysozyme which destroys bacteria.

The coats of the eyeball. The wall of the eye is composed of three layers or coats—an outer, a middle, and an inner (see Fig. 13.1).

The outermost layer or sclerotic coat is fibrous in character; it preserves the form of the eyeball and protects the more delicate vascular




Fig 13.1 Horizontal section through the eyeball The dotted lines, HC, mark a narrow canal in the vitreous body (hyaloid canal) which, in the embryo, lodged the hyaloid artery

and nervous coats within. The sclerotic is transparent in front, where it forms the cornea. It also composes that opaque exposed part of the eyeball surrounding the cornea, and commonly called the white of the eye.

The middle layer or vascular coat, contains the main arteries and veins of the eyeball. It completely surrounds the globe, except for a small circular opening in front called the pupil. The vascular coat consists of the choroid, ciliary body (p. 591), and tris (p. 607) which

is the annular colored band surrounding the pupil. The choroid comprises the posterior five sixths or so of the vascular coat.

The innermost layer, retina, or nervous coat contains the receptors for sight—that is, those elements highly specialized to respond to stimulation by light, and to convert luminous energy into nerve



Fig 13 2 The lacrimal apparatus. 1, lacrimal ducts, 2, lacrimal sac; 3, nasolacrimal duet marked off by interrupted line indicates the position of the lacrimal gland.

impulses. The retina is developed as an outgrowth of the primitive brain; it is, therefore, essentially nervous in structure and function. It consists of several layers; the chief of these from within (i.e., from the interior of the eyeball) outward are as follows:

- 1. Layer of optic nerve fibers
- Layer of ganglion cells
   Inner plexiform layer
   Inner nuclear layer
- 5. Outer plexiform layer
  6. Outer nuclear layer
  7. Layer of rods and cones

- 8. Pigment layer

The layer of nerve fibers is composed of the axons of the ganglion cells of the next underlying layer (see Fig. 13.3). The fibers turn horizontally a short distance from their origins and, converging toward the posterior wall of the globe, form the optic nerve.

The cells of the layer of ganglion cells are of different sizes and shapes. The larger cells are called giant ganglion cells, the smallest ones, midget ganglion cells. The central processes (axons) of the cells of this layer are the fibers already mentioned as forming the innermost layer and the optic nerve. Their dendrites pass outwards and form synapses with cells of the inner nuclear layer.

The inner nuclear layer contains relatively small, oval, and flattened cells whose central processes synapse in the inner plexiform layer with the peripheral branching processes (dendrites) of the ganglion cells. Their peripheral (outward running) processes synapse in the outer plexiform layer with the fibers of the rods and cones. The outer nuclear layer is composed mainly of the fibers and nuclei of the rods and cones. The plexiform layers are composed of fibers of the cells in the nuclear layers (or ganglionic layer) on either side, where, as just mentioned, synapses are formed.

The rods and cones (Fig. 12.3) are the receptors of sight. They are




Fig 13.3 The layers of the retina upper drawing below larger drawings of the rods and cones and the pigment cells, surface view

modified nerve cells. When light strikes them, impulses are set up and transmitted to the nerve cells of the inner nuclear layer, and thence the ganglion cells, and by the axons of the latter (which constitute the optic nerve) to the visual area of the cerebral cortex (see p. 536). It is evident from their position that light, in order to reach the rods and cones, must penetrate all layers of the retina except the outermost one of pigment cells. It should also be mentioned that, though the visual receptors are stimulated most effectively by light, they respond also to mechanical stimula A blow, or even light pressure upon the eyeball will cause a visual sensation in the form of a flash circle, or star of light. The sensations of light produced by pressure or other stimulus applied to the eve with the lds closed are called phosphenes.



Fig 13 4 The blind spot Close the right eye hold the figure about 6 inches in front of the left eye and look steadily at the white

disc. Move the book slowly toward the eye until the cross disappears. When this occurs the image of the cross has fallen upon the entrance of the optic nerve from which rods and cones are absent, it is therefore insensitive to light.

The layer of pigment cells consists of a single stratum of hexagonal epithelial cells. The cells send our protoplasmic processes which are insinuated between the rods and cones. The inner zones of the bodies of the cells contain numerous round and rod shaped granules of a dark brown pigment. The pigment serves, like the black paint on the inside of a camera to absorb light which otherwise would be reflected and diffused causing blurring of the retinal image. In the eyes of the frog and other cold blooded forms, a strong light causes the pigment gran ules to migrate into the cell processes surrounding the bases of the rods and cones, and thus to prevent more effectively the diffusion of light from one receptor to another. In dim light, the granules become concentrated toward the cell centers.

The fundus oculi, The posterior part of the interior of the eyeball is called the fundus (fundus oculi). It can be examined in the living subject by means of an instrument known as an ophthalmoscope/which throws a beam of light through the pupil onto the retina. The optic papilla is situated near the center of the posterior wall of the eyeball (see Pl 12B). It appears as a pale disc, and is produced by the convergence, as into a vortex, of the visual fibers to form the optic nerve The retinal blood vessels piecre the papilla near its center, and cross its face to reach the retina. The papilla is composed entirely of nerve

fibers, the rods and cones and other retinal layers being absent. It is therefore, insensitive to light—that is, an object is invisible if its image falls upon this small area. The optic papilla is for this reason, called the blind spot of the retina. The reader is referred to Figure 13.4 for a demonstration of the blind spot in his own eye.

The macula lutea. The macula lutea (yellow spot) is a small yellow.

The macula lutea. The macula lutea (yellow spot) is a small yellow ish area of the retina situated a little to the outer side of the optic papilla (see Pl 12B) The color of this area is due to the presence of a yellow pigment. In the center of the yellow spot is a minute depressed area known as the fovea centralis (Fig. 13.5). All the layers of the




Fig. 13.5 Microscopical appearance of the fovea centralis the region of most acute vision at contains cones but an man no rods

retina except the rod and cone and pigment layers are extremely thin in this area, it is only about  $\frac{1}{4}$  mm in diameter  $(\frac{1}{2^{10}}$  inch). The fovea is the region of most acute vision, when we look at an object, the eyes are directed so that an image of the object falls upon the fovea of each eye Vision in the rest of the retina (called the peripheral or extrafoveal retina) is much less sensitive, and enables us to gain only a dim, ill defined impression of our surroundings. The reader can demonstrate this fact for himself if the looks straight ahead and tries to define objects on either side or above and below his line of vision.

The crystalline lens and iris (Fig. 13.1) and the muscles of the eye ball will be described when the functions of these structures are being considered.

The duplicity theory of vision. According to this theory, which is now universally accepted the rods and cones have different thresholds of excitation and also show differences in function. Vision in bright light (photopic or daylight vision) and the perception of color are functions of the cones The rods, on the other hand, having a much lower threshold than the cones, make it possible to see in dim light (scotopic or twilight vision) That is to say, the rods respond to light of much lower intensity than is required to excite the cones. But the rods do not give rise to any color sensation, this is a sensation entirely dependent upon the cones Rod vision is in black, white, or a mixture of the two, namely gray 1 Thus it is that at twilight the color of the landscape fades, or seems to be brushed over with a gray wash The cones, though having a higher threshold to light stimulation than the rods, are capable of much more acute vision, making it possible for us to perceive the more precise form and finer detail of objects Some species of birds though they possess very acute vision, have few or no rods in their retinas the part of the human retina that is capable of the most acute vision-namely, the fovea centralis-contains only cones, or at the most, only a scattering of rods When we look at an object the eyes are so turned that the image falls upon this small cen tral area of the retina Surrounding this rod free region-the extra foveal or peripheral retina-there are both rods and cones, but the cones become progressively less numerous and the rods more so from the center of the retina to its more peripheral parts until, at the extreme limit rods are present alone

The retinas of night birds, such as the owl, and certain mammals such as the rat, contain only rods

The greater sensitivity of the peripheral part of the retina (rods) in dim light as compared with the foveal region (cones) is sometimes demonstrated in an interesting way when on a dark but starry night we may see a small star out of the "corner of the eye" (ie, when its image falls on the extrafoveal part of the retina) which disappears when our eyes are turned directly toward where it was seen, its image has now fallen on the fovea which in the prevailing light is blind

If a subject is seated in a dark room and a colored object is placed before him which even after his eyes have become accustomed to the dark, he cannot see, and the object is then very gradually illuminated he sees it without color As the illumination is further increased, and the threshold of the cones is reached, the object is seen in finer detail

<sup>&</sup>lt;sup>3</sup> It is thought that the rods though they possess no power of color distribution, do nevertheless give a uniform blue tust to a d may lighted scene. The blue cast gives to our surroundings in the moonlight, in which rods alone are functioning gives support to this belief.

and in color. The interval between the threshold of the rods and that of the cones—that is, the interval from when the object is first seen to the perception of its color—is called the colorless or achromatic interval. The rods, like the chemical on a photographic film, are almost insensitive to deep red. The achromatic interval for this color is, therefore, very short. That is to say, in a dim light a deep red object appears nearly or quite black, for black is a sensation caused by complete absence of retinal stimulation. That is the reason that at twilight in the garden the red flowers lose their color, appearing as dark blotches, whereas yellow and blue flowers, though they appear without color, stand out as light patches against the background.

The Purkinje phenomenon. The evidence for the duplicity theory of retinal function is overwhelming. The Purkinje effect, for example, clearly shows that two types of visual receptor are present in the human retina. The Purkinje phenomenon is the shift of the brightest part of the spectrum with changes in the intensity of illumination. If the spectrum (the series of colored lights from red to violet, as in the rainbow, into which white light is split by a prism) is viewed at high illumination, its brightest part is in the yellow, but, when the light intensity is reduced to that of twilight, its colors fade. The brightest part is now in the green; it has shifted to the right. The cones are stimulated maximally by pure yellow light, the rods by light green.

Dark and light adaptation. It is a familiar experience that we are for a time quite blind after passing into a dark room from a bright one. After a short time we begin to see dimly the objects around us, and it almost seems that a famt light is gradually illuminating them. The effect, however, is due entirely to changes occurring in the eye which, it is said, becomes dark adapted. The process is called dark adaptation.

<sup>&</sup>lt;sup>2</sup> There may be some confusion here in the reader's mind with regard to the statement that deep red light does not simulate the rods, or does so very slightly. This color, when the illumination is below the threshold of the cones though above that of the rods, does not cause a sensation of light at all, whereas green, blue, and other colored objects appear as varying shades of gray, at low light intensities. In other words, an object, which in bright light appears dark red, causes, when viewed in dim light, little or no photochemical reaction, therefore, it appears black, which is the negative "sensation" resulting from the absence of all retinal stimulation. This fact was made good use of during the late war. It was, of course, of the utmost importance that in air raids at might over enemy territory, the visuo of the crew should be at maximum efficiency. Their eyes must be fully dark adapted. But it would be most riksome for the crew to be kept in the dark, for an hour or so before they took off on their mission. They were therefore confined in a room illuminated by red light, where they were able to amuse themselves playing cards, or even reading. The breakdown of rhodopsin in this light was negligible, might vision was therefore maintained at maximum efficiency. See visual purple, p. 582.


Seeing with dark-adapted eyes is called scotopic vision or rod vision Dark adaptation is due to,

- Increased sensitivity of the rods as a result of the regeneration
  of the light-sensitive substance visual purple (see below), and
   Dilatation of the pupil which permits more light to enter
- the eve

We also have experienced the dazzling effect of a bright light after our eyes have been accustomed to the dark. But after a time we can look at the bright objects around us without discomfort. The eyes have now become light adapted, the processes bringing this about are referred to as light adaptation. Vision in the light-adapted eye is termed photopic and, since it is due to the cones, may also be called cone vision. The events which take place in the eye are the opposite of those occurring in scotopic or rod vision-namely,

- (1) Bleaching of visual purple and reduced sensitivity of the rods, and
- (2) Contraction of the pupil.

Visual purple or rhodopsin. The outer segments of the rods contain a rose-colored pigment which has been named, though somewhat inaccurately, visual purple. It also goes by the name of rhodopsin (Gk. rhodon, a rose + ops, the eye). This substance is of protein nature, with a molecular weight around 300,000 A pigment of the carotene group (p. 385) is linked to the protein. This complex compound (protein + pigment) is bleached by light rays, the pigment being split from the protein and converted to vitamin A. These products are colorless. An intermediate step in this photochemical change is the formation of an orange-yellow pigment called retinene. Thus,



In the dark, rhodopsin is resynthesized from vitamin A and proteig!

A photochemical substance—violet colored—is also believed to be present in the cones, but much less is known about it. It has been named iodopsin (Gk. iodes, violetlike).

The bleaching of rhodopsin by light can be demonstrated in a rabbit's eye which has been recently excised in a darkroom. When the eye is directed toward some brightly illuminated object, such as a window against the sky, the outlines of the window frame can be made out upon the retina, when the eye is opened and examined in a dim light. Where the light fell upon the retina, the visual purple was bleached, but was left unchanged where the bars of the window cast their shadow (Fig 13 6) The excised eye can thus be made to behave as a camera, the visual purple acting like the chemical in the

is called an optogram It is evident then that the initiation of im pulses in the nervous elements of the retina is dependent primarily upon a photochemical proc ess Defective vision in dim light (twilight)-an abnormality known as night blindness or nycta lopia-is a manifestation of vitamin A defi ciency or is due occasionally to a congenital lack of visual purple (see also p 386)




Fig 136 An optogram The circular area with radiating vessels is the optic

The sensitivity of the retina. The retina of the disc (blind spot) dark adapted eye is capable of responding to an almost infinitesimal amount of radiant energy (1e, light with wavelengths between 4000 and about 8000 millimicrons) It is some 3000 times more sensitive than a rapid photographic film A single rod it has been calculated, will give a response when the smallest possible quantity of light (green) energy falls upon it Dr Pirenne has esti mated that the energy liberated when a pin drops to the ground from a height of 1 inch, if converted to light energy, would be sufficient to give a faint visual sensation to every man who had ever lived! The rods are not equally sensitive to light of all wavelengths, they are most sensitive to light green rays (wavelength around 5000 millimicrons) The cones show their greatest sensitivity for yellow Sensitivity for both types of receptor is much less for the longer (red) and shorter (violet) light waves It has already been mentioned that the rods are almost insensi tive to deep red

Those who have had even a very limited experience in photography know that, with respect to the scene that we propose to photograph, two factors must be taken into account—namely, (1) the time of exposure (1 e, the interval between the opening and the closure of the shutter during which the light enters the camera) and (2) the size of the opening or "stop" of the diaphragm. It is entirely a question of the quantity of light to which the sensitive film is exposed. So it is, within limits, with the retina, a light stimulus of a certain intensity and dura tion that is just capable of causing a visual sensation becomes inef

fective if either its duration or its intensity is reduced. Or a light stimulus which is just beneath the threshold becomes effective if its duration or its intensity is increased. A similar relationship holds between the size of the retinal area that is stimulated and the intensity of the light. That is to say, a larger object dimly illuminated will be seen, whereas a smaller one even though somewhat more brightly illuminated may be invisible. If the intensity of the smaller object is increased so that the total quantity of light energy is equal to that falling upon the retina from the larger body, it then arouses a visual sensation.

The reciprocal relationship between duration and light intensity



Fig. 13.7 Showing the reciprocal relationship between the intensity and duration of a visual stimulus Records are from a single isolated fiber of the optic nerve. (After Hartline and Graham.)

#### 0.20

has been beautifully shown by Dr. Hartline and his associates. A flash of light of varying duration and intensity was thrown into the eye of the king crab, and the impulse frequency recorded from a single optic nerve fiber (see p 475) It will be observed in Figure 13.7 that the frequency of the impulses (intensity of response) is the same when the product of the duration and the intensity of the stimulus (duration × intensity) are equal, though one or other factor was varied.

Visual acuity. The ability of the eye to determine the precise shape and detail of any object, or to recognize the separateness of two small objects placed close together, is referred to as the visual acuity. It is exhibited in the highest degree by the fovea centralis which, in the human retina, is composed entirely of cones. It is seriously reduced by refractive errors (myopia, etc., p. 611). As a test for visual acuity in the diagnosis of refractive defects of the eyes and to aid in their correction by means of glasses, the oculist most usually employs Snellens test type (Fig. 13.8). These consist of heavy blook letters printed in black upon a white ground. The letters are of graded sizes, arranged in nine horizontal rows, the largest letters in the top row, the smallest in the lowermost.

The subject is seated 6 meters (approximately 20 feet) from the letters and asked to read the rows of type down to the smallest which

it is possible for him to distinguish. The visual acuity is expressed as a fraction, the numerator of which is the distance in meters at which the subject can read a given line of letters, and the denominator is the distance at which a person with normal average vision can read the same line. For example, a person with normal vision can read the seventh line at 6-meters; if the examinee can read this line, his vision is  $\frac{e}{6}$ , or normal. If, on the other hand, he can read only the third row which can be read by a person with normal vision at 24 meters, then his vision is only  $\frac{e}{4\pi}$ , or  $\frac{1}{4}$  normal.

The examiner, in testing for defective vision in this manner, fits removable lenses (trial lenses) before the subject's eyes while he reads. By using trial lenses of different kinds and power, and noting the improvement or lack of it in the patient's sight each time, the examiner is able to determine the type and degree of the refractive error that exists (pp. 610–11). He then prescribes the appro

T B
D L N
P T E R
F Z B D E

Fig. 13.8 Snellen's test type.

OFLETO

exists (pp. 610-11). He then prescribes the appropriate convex, concave, or other form of spectacle lens required to correct the defect. Optical action currents: The electroretinogram. When the retina is

stimulated by a beam of light thrown into the eye, action currents are set up which can be recorded by a galvanometer. The current is led off to the galvanometer by means of electrodes, one placed upon the cornea of the eye in its usual forward-looking position, and the other on any moist surface of the body. A record of such changes in electrical potential is shown in Figure 13.9. It is called the electroretinogram. Four waves are to be seen; they are designated a, b, c, and d. The wave a is a negative wave—i e., below the base line; the others are positive. The waves a, b, and c are inscribed when the light is flashed into the eye; d occurs when the light is cut off.

Action currents can also be demonstrated in the optic nerve when the retina is stimulated by light. A discharge of impulses then occurs which increases in frequency with the intensity of the illumination,

Fig. 13.9 The electroretinogram. Waves a, b, and c occur when a light is flashed into the eye; d occurs when the light is turned off.

but no change occurs in the amplitude of the waves, no ruatter how intense the light

stimulus may be. The relationship between frequency of the impulses and the intensity of the stimulus, which we have seen to be a characteristic of other kinds of nerve, has been demonstrated in a single nerve fibers separated by dissection from the other fibers of the optic nerve (see Fig. 13.7, p. 584).

The visual fields. The part of the outside world seen by one eye at any moment is called the visual field of that eye. In man, the ape, and other species whose eyes are placed in the front of the head, the visual fields of the two eyes overlap In other words, a large part of the outside world within the range of vision is seen by both eyes at the same time. This type of vision is called brocular. In the rabbit, horse, and other animals with eyes placed laterally in the head, the visual fields overlap to a very small extent; in these, vision is almost entirely managingly.

In those species possessing binocular vision, light rays from the outer or temporal half and from the inner or nasal half of either visual field fall upon the nasal half and temporal half, respectively, of the cor-responding retina (see Pl 13A). Now, the optic nerves come together a short distance behind the eyeballs to form the optic chiasma. Here, the fibers which have originated in the nasal half of one retina cross with those coming from the nasal half of the other—that is, fibers from the nasal half of the right retina cross to the left side of the brain, those from the nasal half of the left retina to the right side of the brain. Fibers from the temporal halves of the retinas continue uncrossed Thus, the visual pathway, on each side, from the optic chiasma to the cortex of the occipital lobe (visual center) transmits impulses from the nasal half of one retina and the temporal half of the other. Owing to this peculiar course taken by the visual fibers, the loss of sight following injury or disease of one optic nerve differs from that resulting from a lesion of the fibers anywhere between the chiasma and the occipital cortex, or of the visual center itself If the function of one retina or of one optic nerve is destroyed, complete blindness of the corresponding eye results, but the sight of the other eye is unaffected. Interruption of the fibers in their course from the chiasma to the visual area (optic tract), or a lesion of the visual area itself, causes blindness in one half of each retina. For example, if the injury involves the left optic tract or left visual cortex, blindness of the nasal (left) half of the right retina and of the temporal (left) half of the left retina (opposite halves of the visual field) results Blindness in one half of each eye is called, hemianopia (half blindness). The nasal half of one retina and the temporal half of the other are called homonomous halves. The type of

blindness just described is therefore called homonomous hemianopia Rarer types of hemianopia result from bilateral involvement of the temporal fibers alone at the chasma or of the nasal fibers alone, these types are given the qualifying terms of binasal and bitemporal respec tively (the designations binasal and bitemporal refer to the visual fields not to the retinas)

The shape and extent of the visual fields in man can be mapped out by means of an instrument known as a perimeter which is shown in




Fig 13 10 The perimeter This is an instrument used for mapping the visual fields. It consists of a metal band curved in a large arc of a circle near the center of which is a rest for the subject s chin The metal arc carries a small object which can be moved to or away from the center The are itself can be rotated into any position (meridian) it is rotated into different meridians in succes sion and at each new position the test object is moved slowly from a peripheral point beyond the subject s field of vision toward the center. The examinee indicates the moment that he sees the obsect ( out of the corner of his eye as it were) The points in each meridian so determined are

marked upon the chart shown on the instrument and in detail in Figure 13 11 c chin rest e position of subjects eye

Figure 13 10 A chart or map of the visual field of one eye is given in Figure 13 11. The instrument is often of great value in the diagnosis of diseases of the retina or of the visual pathways.

From the optic chiasma the visual fibers—that is the axons of the ganglion cells from the nasal half of one retina and the temporal half of the other—proceed on each side of the brain to a collection of nerve cells in the midbrain. These latter form on each side a small eminence known as the lateral geniculate body. The optic fibers from the chiasma to the lateral geniculate body are known as the optic tract. The fibers of the latter terminate by forming synapses with the cells of the lateral geniculate bodies which are therefore primary visual centers in lower animals but not in man visual perception is possible to a variable degree through these centers alone—that is, after the occipital




Fig 13.11 Chart of the normal visual field of the right eye Note the restriction of the visual field on three sides, caused

The first of these has its cell body in the inner nuclear layer of the retina; the second is the ganglion cell of the retina; and the third is a

by the nose, brow, and cheek

nerve cell in the midbrain (Fig. 13 12). Identical or corresponding retinal points. Objects in the binocular field of vision must, obviously, cast an image on each retina. Yet the two retinal images do not cause a double sensation; we do not see things in duplicate. The explanation for this fact is based upon the theory of identical or corresponding retinal points. Points in the nasal half of one retina are paired with corresponding points in the temporal half of the other. The ocular muscles direct the axes of the eyes so that an image falls upon each of these corresponding points. As mentioned above, the optic fibers from the homonomous halves of the retinas proceed to one side of the brain, the left halves (nasal of right eye and temporal of left) to the left visual area, the right halves (nasal of left eye and temporal of right) to the right visual area. The two images are fused in consciousness into a single sensation. If, for any reason, the eyes are not directed in such a way that the two images fall on corresponding retinal points, double vision-diplopia-results. Diplopia as a permanent condition occurs most commonly as a result of paralysis or weakness of the ocular muscles. It is seen as a temporary state in alcoholic intoxication, being then due to imperfect control and, in consequence, to unbalanced action of the muscles of the two eyes One can cause double vision in oneself by simply pressing upon the outer side of one eye, and thus displacing the eyeball slightly from its

cortex has been destroyed. The axons of the cells of the lateral geniculate bodies relay the visual impulses through the inter-nal capsule (p. 539) to the highest visual center situated in the cortex of the occipital lobe of the cerebrum. The nerve fibers from the lateral geniculate body on each side to the corresponding occipital cortex are called the optic radiation. A nerve impulse set up by stimulating the visual receptors in the retina, therefore, travels over three neurons to reach the cortex of the occipital lobe.




Fig 13.12 Diagram of the visual pathway.

normal position. Two images are also seen if an object, such as a pencil, is held about 3 inches in front of the root of the nose, or if the pencil is held at arm's length, and the eyes then directed to look at a point some distance beyond it. In the first instance mentioned, it is not possible to converge the eyes sufficiently to bring the two images on to corresponding points on the retinas; in the second instance, images of the near object fall upon noncorresponding points of the retina, because the convergence of the eyes when looking at distant objects is less than that required for near vision.

Elementary optical principles. A ray of light passing from one transparent medium, such as air, into another, such as water, is bent. That is the reason that an upright object, such as a log standing in water, gives the illusion of being broken where the air and water meet. This bending or deflection of the rays of

light is called refraction. Similarly, light rays passing through a glass prism (Figs. 13.13 and 13.14) are bent toward its base. A glass lens having both its surfaces convex, such as the crystalline lens, acts, in respect to the refraction of light, as though it were composed of two prisms placed base to base. Therefore, the rays of light reflected from any object must, after passing through such a lens,




Fig. 13.13 Showing the bending of light rays by a prism. Parallel rays, continuous lines; diverging rays, interrupted lines.




Fig 13 14 The bending of parallel rays of light by a biconvex lens and bringing them into focus

meet at a point (focus) on the other side. Light rays from every point in an object are thus brought to a focus, an image of each being formed. If the lens does not bring the rays to a point, circles or haloor light are formed instead of sharp points. The entire image of the object normally constituted of innumerable images of bright points is blurred accordingly. The failure of a sharp focus to be obtained may be due to the surface upon which the image is cast intercepting the light rays, either before they have come to their meeting point or after they have met and crossed as shown in Figure 13.15.

Light rays coming from infinity, or for practical purposes from any distance over 20 feet are parallel, whereas those reflected from a near object are divergent—that is, they radiate from every point in the object (see Fig. 13-16). In order, then to bend the rays from a near object so that they will meet—ie, be focused—at the same distance be hind the lens as will parallel rays, the lens must be able to bend the rays more acutely in the former instance than in the latter, in other words, its refracting power must be greater. The refracting power of a convex lens increases with the degree of its convexity.

A concave or a biconcave lens (i.e., one with two concave surfaces) acts like two prisms placed apex to apex (Fig. 13 17) and since the rays are bent toward the bases of the prisms, such a lens diverges or spreads the rays of light

A line drawn horizontally through the center of a lens is called its





Fig 13 15 Showing rays focused at S and intercepted at S1 and S1

Fig 13 16 Showing parallel rays of light (A) from a distant object be ing brought to a focus by a biconvex lens, and diverging rays (B) from a near object being focused at the same distance behind the lens. In the latter instance, however, the lens bends the rays more strongly This it is able to do because it is more convex



principal axis, and any line drawn diagonally through the center is called a secondary axis Rays of light passing through the principal axis or any secondary axis are not refracted The crystalline lens. The crystalline lens is a transparent biconvex

disc, about I inch in diameter, situated within the globe of the eve immediately behind the iris. It is enclosed in a thin membrane called the capsule The central part of the lens lies a little distance behind the pupil It is suspended in this position by a delicate annular ligamentthe suspensory ligament-which blends with the lens capsule, and is attached circumferentially to the interior of the globe through the ciliary body The latter also connects the iris with the choroid, and con sists of involuntary muscle fibers and a number of processes-the ciliary processes. The latter-about 70 in number-are arranged in the form of a circle to which the suspensory ligament is attached (see Fig 13 18) The iris and ciliary body divide the part of the eyeball in front of the lens into an anterior and a posterior chamber (Fig 13 1, p 575) Both chambers are filled with a clear limpid fluid called the . aqueous humor, a gelatinous transparent substance called the vitreous body . fills that part of the eyeball lying behind \_ the lens

Though the healthy crystalline lens is perfectly transparent, in the condition known as cataract (p 615) it becomes semitransparent or opaque, causing blindness in the affected eye




Fig 13 17 Parallel rays of light being diverged by a bi concave lens




Fig 13 18 The eye ball from in front, cornea and iris removed 1, suspensory ligament 2, lens, 3, choroid, 4 iris cut edge 5, ciliary processes (black) (Re drawn and modified from Schultze)

The refraction of light rays by the eye—the formation of an image upon the retina. There is a very close resemblance between the eye and a camera In both, the rays of light are refracted (bent) and brought to a focus upon a light sensitive surface—the photographic film and the retina respectively The refracting medium in the camera is, of course, the lens In the eye, light rays are not only refracted by the crystalline lens, but by the cornea as well and, to a slight degree, by the aqueous humor and vitreous body The image of an object falls upon the photographic film in an inverted position and reversed from side to side, the retinal image is similarly inverted and transposed Yet, we do not see objects upside down, because experience has faught us to interpret the retinal messages in accordance with reality (See Fig 13 19, lower sketch) In consciousness, the image is again inverted,




Fig 13 19 Upper drawing the formation of an inverted image upon the return Below, diagram showing the formation of an image by a convex lens 1, principal aixs 2 secondary aixs 3, optical center of Iens 4 principal focus, 5 screen Note that rays passing through the principal and secondary aixs are not refracted.

and reversed from side to side, so that it appears to us in its true position. We have come to know by experience (i.e., previous information received through other senses, especially of touch and hearing) that stimulation of the upper or the lower part of the retina, or of one side, is caused by light reflected from an object situated, respectively, in the lower, upper, or other side of the visual field. We are scarcely conscious even that the retina has been stimulated in the way that we feel and localize a touch upon the skin. We "see" the object at a distance and in a position the reverse of that which the image of the object occupies upon the retina. In a word, the object's image is projected in consciousness to its true position in space—i.e., in the visual field.

consciousness to its true position in space—i e., in the visual field. In Figure 13.19 (upper sketch) are shown the paths taken by rays of light coming from an object (represented by the large arrow) placed a little in front of a biconvex lens. The diverging rays are refracted by the lens, and brought to a focus on the screen S, which may be taken to represent the retina. Rays from each point or the object are brought to a point upon the screen, but in a reversed position, those from the upper part of the object being directed to the lower part of the screen and vice versa (only rays emitted from points at the head and tail of the arrow are shown). Similarly, those from the lateral halves of the object pass to opposite sides of the screen. Thus, a small and inverted image of the object is formed. One should note (Fig. 13.16) that the periphery of a lens refracts more strongly than parts more centrally placed, and that rays of light passing through either the principal axis or the secondary axes are not refracted. In the lower sketch in Figure 13.19, the formation of an image upon the retina is illustrated.

The retinal image can be observed in the excised eye of an albino rabbit. When the eye is directed toward a lighted candle and viewed from behind, an inverted image of the flame is seen shining through the unpigmented wall of the globe. This observation was first made by Christopher Scheiner in 1619. It is worth remembering that the image formed upon the retina is minute; the image of the smallest visible object is little more than 10,000 mm) in diameter

Accommodation of the eye. We have seen that rays of light coming from a distant object are approximately parallel; those from a near object are divergent. Therefore, with a convex lens of a given refracting power, rays from a near object cannot be focused at as short a distance behind the lens as can those coming from a distance (see Fig. 13.16, p. 591), which means that near and far objects cannot be focused upon the retina at the same moment. In order to turn the eyes from a far scene and focus them on a near one, either a stronger lens (that is,

one with a greater convexity) must be used, or the distance between the screen and the fens must be increased

The latter means is employed in photography. In order to take a photograph of a view less than 12 feet or so from the camera, the lens must be moved forward so as to increase the distance between it and the film. Though the eyes of certain species of fish have a mechanism for accommodating the eye to near vision based upon this principle, in man and the higher animals the first mentioned method is adopted, the




Fig. 13.20. Illustrating the mechanism of accommodation of the eye for near vision. The horizontally shaded lens and the unshaded irs show the position of the parts when at rest the vertically shaded lens and ins show the position during accommodation for a near point C, ciliary muscle S1, suspensory ligament (Redrawn from Landolt)

crystalline lens can alter its refracting power in order that either a far or a near object can be distinctly seen. When the eyes are directed to a distant view, the crystalline lens is least convex, being flatter, its refracting power is at the minimum. But when the observed object is within about 20 feet from the eyes, the convexity of the lens increases in order to bring the divergent rays from the near object to a sharp focus upon the retina. These together with the other adjustments made by the eyes in order to see either near or far objects clearly are termed ocular accommodation.

The far and near points of the eyes. The farthest point from the eyes at which objects can be focused upon the retina and, therefore, seen distinctly is called the far point of vision (punctum remotum). The far point is at infinity, for the visibility of a distant object is limited only by its size, the clearness of the atmosphere, and the curvature of the earth, for practical purposes it is taken as any distance beyond 20 feet, at this distance the lens is, as just mentioned, at its least curvature. The shortest distance from the eyes at which an object can be seen clearly

is called the near point (punctum proximum), it lengthens from in fancy to old age, in accordance with the gradual diminution in the elasticity of the lens, being between 7 and 9 cm for the former age and 40 cm for the latter (see Presbyopia, p 610) For the normal eye of the young adult, the near point is about midway between these two extremes—namely 25 cm approximately At any distance between the eyes and the near point an object forms a blurred image upon the retina because the lens is unable to increase its refracting power (convexity) any further. The reader can verify this for himself by bringing the tip of a pencil close to his eyes.

The mechanism of accommodation. The increase in the convexity of the lens and, therefore in its refracting power are brought about by




Fig. 13.21 Illustration by means of a mechanical model of the man ner in which the lens becomes more convex when the eye recommo dates for near vision right half of figure. The springs. S and S represent the elastic choroid and the radiating lines the suspensory ligament

the inherent tendency of the lens to assume a more convex shape when the restraint offered by the suspensory ligament is released (see Figs 13 20 and 13 21) The suspensory ligament (which as already stated extends outward from the capsule of the lens to be attached to the citi ary processes and through these and the citiary muscle to the inner coat of the eye) is taut when the eyes look into the distance. The ten sion this exerted upon the lens reduces its curvature. When the eye accommodates for near vision the ciliary muscle contracts and draws the choroid forward, the ciliary processes are brought closer to the lens—i.e., they form a smaller circle. The suspensory ligament is slackened thereby and the tension on the lens diminished. The lens by virtue of its inherent elasticity bulges forward (Figs. 13 20 and 13 21). The convexity of the central part of the anterior surface is increased to a much greater extent than the more peripheral part of the lens. Little change in the curvature of the posterior surface occurs.

Scheiner's experiment. That the crystalline lens alters its refractive power when the eyes are turned from a distant scene to look at an ob-

596

ject nearer to the eyes was first deduced by Christopher Scheiner (1579-1650), a German priest and scientist, from the following experiment

A card is pierced by two pinholes about 2 mm apart horizontally (Fig 13 22), a needle, N, is held in front of the card, C, while a convex lens placed behind it throws a clear image of the needle on the screen S at n (upper cut, Fig 13 22) If the screen is moved forward (5'), it intercepts the rays from the needle before they have come to a focus, two blurred images are formed at n' and n' If the screen is




Fig 13 22 C, perforated card, L, lens (Explanation in the text)

moved to a position behind the original one (S''), it is now in the path of the rays after they have crossed at the focus n two blurred images are formed at n'', n'. But if we had a series of lenses graded in refracting power, the image of the needle could be focused on the screen placed at varying distances

The lower cut of Figure 13 22 depicts the pierced card held at a distance of about 5 inches in front of one eye. The other eye is closed. The pinholes should be separated by a horizontal distance a little less than the diameter of the pupil. When the subject looks at an object, 0, at some distance beyond the needle, N, two blurred images of the needle are cast upon the retina (dotted lines), because the rays, as at n' and n' above, have not been refracted sufficiently. If the subject now looks directly at the needle, the two images fuse into a single clear one. Since we know that in the latter instance the retina cannot have moved back to the point of focus it must be concluded that the crystalline lens has increased its refracting power.

Purkinje-Sanson images The change in the form of the lens during

accommodation for near vision can be demonstrated by another simple experiment. If one looks into another's eye in a dark room while a lighted candle is held a little in front and to its outer side, one sees three reflected images of the flame (Fig. 13.23). Of these, one is bright and upright; it shines from the surface of the cornea which acts as a convex mirror. A larger but dim upright image is formed by the anterior convex surface of the lens. The third image is inverted, bright and smaller than the others; it is formed by the posterior surface of the lens which acts as a concave mirror. If the positions of the images are noted while the subject gazes at a distant point, and he then

Fig. 13 23 Purkinje-Sanson images A, during far vision; B, during accommodation for near vision (Redrawn and modified from Williams)



accommodates for near vision, the image reflected from the anterior surface of the lens will be observed to become smaller, and to move toward that reflected from the cornea which, of course, does not alter in size or position. As a rule, no change can be observed in the reflection from the posterior surface of the lens. The change in the position and size of the image on the anterior surface during accommodation can only mean that this surface becomes more convex.

The other adjustments for near vision are convergence of the eyes, which is necessary in order to bring the retinal images on to corresponding points (p. 588), and constriction of the pupil. The narrowing of the pupil permits light to pass only through the more central part of the lens which, as just mentioned, is increased in its convexity to the greatest degree during accommodation.

The appreciation of distance and depth—stereoscopic vision (Gk. stereos, solid + skopeo, I view). Our visual judgment of solidity—that is, our recognition that an object has depth as well as height and width—is due largely to the fact that the two retinal images are slightly dissimilar. If the reader will look at some object in front of him, first closing one eye and then the other, he will find that the view seen by the right eye is slightly different from that seen by the left. The right éye is able to see more of the right side of the object, the left eye more

of the left side (Fig. 13 24). The two slightly dissimilar images are fused in the brain, yet the composite image has, hidden within it, some thing of each separate one, upon this the stereoscopic effect largely depends. The fusion of the dissimilar images by the brain, and the impression of depth and solidity produced thereby, lies in a field of psychology of which little is known.

The instrument known as a stereoscope produces an illusion of solidity by making use of the principle of simultaneous stimulation of corresponding retinal points by slightly dissimilar images. A photo




Fig 13 24 Showing two views of the interior of a long room or corndor (upper figure) View A shows more of the right hand wall than does view B The former A would he seen by the left eye of a person looking into such a room and B by his right eve The fusion of the two similar but not identical images gives a visual sen sation of depth. In the lower figures two views of a block are shown A in which more of the right

side of the block is shown than in B would be the view seen by the right eye B would be the image of the block formed by the left eye. Fusing of the images in the two eyes is an important factor in our perception of the solidity or three dimensional nature of such an object.

If the reader will place one edge of a card about 5 inches wide vertically be tween the two sets of figures with the opposite edge of the card touching his nove and then gaze steadily with accommodation relaxed until the figures of each pair fuse into one a visual sensation of depth (upper pair of figures) and of solidity (lower pair) will be experienced.

graph taken with an ordinary camera appears flat because identical images are formed upon the retinas. A stereoscopic photograph on the other hand, is taken by a camera provided with two lenses which are set like the eyes a short distance apart. Thus two slightly dissimilar views are taken, which when looked at through the stereoscope are projected by means of prisms, one to each eye, so as to fall on corresponding retinal points.

Stereoscopic perception cannot however be explained entirely upon the basis of dissimilar retinal images, for the sensation of depth (or distance) is not abolished when one eye is closed; other factors are concerned. These are:

- 1. The apparent size of various objects in our field of vision. We know from experience the approximate size of the objects which we see, but the image which an object casts upon the fovea diminishes as its distance increases. For example, a church steeple at a distance casts an image upon the retina no larger, perhaps smaller, than does a pencil held a few inches from the eyes. We know the relative sizes of the two objects and, therefore, infer that the steeple must be far away and the pencil near.
- 2. The apparent change in color of an object with distance. The atmosphere is not perfectly transparent or equally so for all wavelengths. Tree-clad hills, which we know to be green, appear blue in the distance; the colors of many other objects appear to fade with distance, their detail and outline being dimmed by haze.
- 3. The blocking out of parts of a distant view by objects between it and the eyes gives a sensation of depth
- 4. Mathematical perspective. Straight lines running into the distance which are actually parallel (or objects along imaginary straight lines) are convergent in the retinal image. When we look down a railway track, for example, the rails appear to converge toward some point beyond the horizon. This arrangement of lines in the image formed upon the retina we have come to associate with distance. Perspective is an elementary principle in drawing, used to create an illusion of depth or distance. The artist draws objects along imaginary lines which run toward a point in the background of his picture.
- 5. Parallax. When one moves in any direction, near objects appear to move in the opposite direction, those in the background in the same direction as ourselves. This apparent movement of near objects in relation to ones farther away is called parallax; it is also produced by a movement of the head or eyes, even though the body remains stationary. Now, involuntary movements of the eyes are continually taking place with the production of parallax; this is believed to be one of the most important factors in giving us a sense of depth.
- The distribution of light and shade over the surface of an object and the shadow which it casts upon its surroundings are also important elements in the production of the stereoscopic effect,
- Color vision. White light is in reality a combination of colored lights—red, orange, yellow, green, blue, and violet—which can be separated by means of a glass prism. The series of colored lights so produced is called the (visible) spectrum (see Pl. 13B). The wavelengths of the

lights of the visible spectrum run from 8000 millimicrons for dark red to around 3000 for violet. Rays somewhat longer than the red are called infrared or heat waves; those shorter than the violet are called ultraviolet; neither, of course, gives a sensation of light. The prism refracts the long red rays less than it does the orange, the orange less than the yellow, the yellow less than the green and, so on, in the order of diminishing wavelengths. A beam of white light is thus split into its constituent colors. The separated colors can be recombined to produce white. But, in order to obtain a sensation of white light, it is not necessary to employ all the prismatic colors; not only white, but all colors can be produced from three—namely, red, green, and violet. These are called the primary colors. If we were to set up three lanterns, one of which emitted red light, another green, and a third violet, we could, by blending one, two, or all three of them upon a screen, produce white or any colored light we chose For example, the three colors in nearly equal proportions would produce white; green and violet would give bule; red and green and a very little volet would give yellow or orange. Several theories of color vision have been proposed. According to

Several theories of color vision have been proposed. According to Young's theory, which is very widely accepted, three kinds of cones exist in the retina, each type containing a substance sensitive to light of a different color. One kind is stimulated by red rays, one by green, and another by violet; white light falling upon the retina stimulates all three nearly equally. When a yellow color is seen, the sensation is due to the stimulation of equal numbers of the red and green, and a very few of the violet elements; if violet and green sensitive cones are stimulated, a sensation of blue results; and when the red, green, or violet type of receptor is excited exclusively, we see red, green, or violet, respectively. See Plate 14A

The qualities of color. There are three qualities of color—hue, saturation or intensity, and tone or brilliance. Hue is determined entirely by the wavelength of the light, thus red differs from green or blue because its wavelength is greater. Saturation or intensity, also called purity, depends upon the proportion of white mixed with the color—pink has a lower saturation than red Tone or brilliance, also referred to as value by artists, varies with the proportion of black mixed with the color. A vivid scarlet pigment, for example, can be reduced to a deep cherry by the addition of black.

Retinal areas sensitive to color. The extreme peripheral part of the retina devoid of cones is insensitive to color. It gives impressions of white, black, and gray only. Color perception is confined to the more central retinal regions where cones are present exclusively, as in the

601

fovea centralis, and in the extrafoveal regions where these receptors are mingled more or less plentifully among the rods. Nor is sensitivity to the different colors uniformly distributed over the retina. The area sensitive to blue is the largest, that sensitive to red is next in order of size; then comes the margin of a much smaller area sensitive to yellow. This latter area is little larger than the area sensitive to green which is the least extensive of all. The four areas may be compared to four transparent saucers, respectively colored and graded in size, placed one inside the other. Light passing through the small green saucer must pass through all the others as well; the area of the retina which it represents is, therefore, sensitive to all four colors. Light transmitted through the outer part of the yellow saucer will also pass through the red and blue, but not through the green saucer; the corresponding area is, therefore, sensitive to yellow, red, and blue, but not to green. The outer part of the red area is sensitive to red and blue; the outer part of the blue saucer transmits only blue light and the rim of the area which it represents is sensitive only to blue.

The color areas of the retina can be mapped out by means of the perimeter shown and described on page 587. Small paper discs colored, respectively, blue, red, yellow, and green are used instead of the white test object employed in mapping the visual fields. Normally, the blue disc will be seen at the greatest distance from the center of the visual field; at a somewhat shorter distance the red disc will appear, then the yellow, and finally the green (see PI. 14B).

Color blindness. In some persons the retina is totally insensitive to color; to such their surroundings are seen in black and white and various tones of gray. Such a condition, called achromatism, is very rare. In the common type of color blindness, the subjects are blind to red or to green On the basis of the Young-Helmholtz theory of color vision, it may be presumed that one of the three color substances present in the retinas of persons of normal vision is absent from the retinas of the red or green blind, the red substance being lacking in the former abnormality, the green substance in the latter. The subjects of these visual defects match any color with only two primary color sensations, green and violet, or red and violet, instead of the three possessed by persons of normal color vision. Their color vision is therefore called dichromatic (Gk. di, two + chroma, color) instead of trichromatic (Gk. tri, three) which is the normal. Red blindness is also called protanopia (Gk. protos, first + anopia, blindness) and deuteranopia (Gk. deuteros, second), which means that the blindness is with respect to the first (red) or to the second (green) primary color, John Dalton

(1766-1844) the celebrated chemist who was red blind and gave the first scientific description of protanopia wrote, 'Crimson has a very grave appearance Woolen yarn dyed crimson or blue is the same to me."

Anomalous trichromatism or partial color blindness is the most common defect of color perception. There is not complete blindness for any color, but the appreciation of red or of green is below the normal. If for example, a person with such a defect is asked to match a pure yellow light with a combination or mixture of red and green lights he uses more of the red light than would a normal person to make a per fect match, his perception of red is defective. If he uses more green his perception of this color is subnormal. The defective perception of red is called protanomally, that of green deuteranomally. A special in strument called an anomaloscope is used for testing these defects. It was devised by Lord Rayleigh the celebrated English physicist.

Ninety five percent of the subjects who suffer from some form of defect in color vision are males. Most cases are inherited and the in heritance is of the sex linked character being transmitted from generation to generation in a manner similar to that of hemophilia (p. 107).

A color blind person may be quite unaware of his visual abnormal ity. Though his color sensations are different from normal persons he is usually able to make some distinction in his own way between different colors usually by some slight difference in brightness. Though great care is exercised in testing the color vision of men such as sea men and locomotive drivers whose occupations require a normal color sense we rarely hear of an automobile driver disregarding a red traffic light because he thought it was green.

Color vision is tested in several ways but the oldest and most usual method is by means of colored wool yarns. The subject is shown a number of yarns dyed in colors which are most likely to be confused (confusion colors), and is risked to pick from the lot the one that matches best another single yarn dyed a rosepink. If he chooses a violet or a blue yarn he is red blind. If green red, and brown yarns are chosen, he is green blind. Anomalous trichromatism is tested, as al ready described, by means of the anomaloscope.

Contrast effects. It is well known that when black is placed against white or vice versa they "set one another off," the black looks blacker and the white a purer white than if either were placed against a colored ground It is also true that blue against a yellow ground (or )ellow against a blue ground) is more vivid than if placed against any other color. Also green is enhanced by red and red by green. These phe nomena are examples of simultaneous contrast.

The maximum effect of contrast is obtained when complementary colors are paired. Any pair of colors which when fused as lights, produce white are said to be complementary to one another. Examples of complementary color pairs are the following.

Red and greenish blue Orange and cyan blue Yellow and indigo blue <sup>1</sup> Violet and greenish yellow Purple and green

Not only those listed above but every color and shade of color has its complementary (Fig. 13.25). There are consequently, a great number of complementary colors and the contrast effect is produced




Fig 13.25 Complementary colors 1 he colors of the spectrum from violet to red are marked upon the curve (roughly transpular) Other colors would fall upon intermediate points on the curve. The base I ne of the figure (from violet to red) represents the nonspectral purple. Any two points upon the curve joined by a straight line drawn through the point marked white ind cate complementary colors. (After Hall.)

when the colors of such a pair are placed side by side Furthermore when a color is laid beside one that is not complementary to it or against white each color appears to be tinted slightly with the complementary color of the other, or the white becomes tinged with the complementary of the color For example blue placed beside green as

A distinct on must be drawn here between I ghts and paints we know that yellow and blue pant when mixed give green not white The color of a paint is due to its reflecting certain wavelengths in white I ght and absorbing the remainder. Even the bluest pant is not pure blue it absorbs all light but blue and a I tile green. These it reflects A yellow paint is not pure yellow it reflects a little green with the yellow. When the two paints are mixed blue I ght is absorbed by the yellow paint and yellow light by the blue. The green rays of which a little is reflected by each separate paint are doubly reflected when the two are me.

sumes a faintly red cast (violet), and yellow against red is tinged faintly green. Again, a lemon or other yellow object, placed in a bright light, casts a shadow which is tinged with blue—the complementary of yellow. The shadow cast by a red object would be tinged with greensh blue, and one cast by a purple object with green. These principles are applied in art. The artist makes a yellow flood of sunshine more brilliant by painting blue into the shadows which, in turn, are given depth and an appearance of reality which they would otherwise lack.

When one stares for a time at a colored surface in a strong light and then directs the gaze to a gray surface the gray appears slightly tinged with the color complementary to the one to which the eyes were first directed, and objects of the complementary color itself are made more vivid. For example, if one looks at a red surface for a time and then at a green one, the latter color is intensified—it appears greener. This phenomenon is called successive contrast.

After images. If one looks at a bright white light for a moment and then closes the eyes or turns them upon a dark surface, an image of the light slowly floats into view, becomes more distinct for a time, and then gradually fades Similarly, if the eyes are stimulated by a colored light or a brightly colored object of any sort, and then darkened, an image of the same color appears. These are called positive after images, If, instead of closing the eyes or turning them to a dark surface after looking at a white light, the retinas are stimulated a second time and diffusely by white—e g, directing the eyes to a sheet of paper—one then sees a dark image against a white ground This is called a negative after image. If the first stimulus was colored, then the after image is in the complementary color. Negative after images of colored objects are the cause of successive contrast described in the last paragraph.

Positive after images are due apparently to the chemicophysical changes in the receptors of the retuna, caused by and outlasting the stimulus Negative after images are thought to be due to a state comparable in certain respects to the refractory period of nerve. Cones which have responded to a stimulus will not for a time respond again. Ordinarily, white light stimulates all three types of cone (p. 600). The negative after image which appears upon applying a circumscribed and then a diffuse white stimulus to the retina is, therefore, a dark patch against a white background. When the object looked at is colored, and the retina is then stimulated by directing the eyes to a white surface, the image is in the complementary color, because only those cones that had not previously been stimuluated can respond. For example,

if the object looked at is red, the red-sensitive cones are not excited by a subsequent stimulus of white; those sensitive to green and to violet alone respond, giving the complementary of red —namely, a bluish green.

Optical illusions. The brain may be deceived by imitations of certain effects upon which our visual judgments of the size, shape, and distance of objects are based. Visual errors of this nature are called optical illusions or optical deceptions. Some interesting examples are shown in Figures 13.26 to 13.29.



Fig. 13.26 Zollner's lines. The long diagonal lines appear to converge; actually, they are parallel.

Movements of the eyes and eyelids. The eyeballs are turned in their sockets by small muscles, attached, on the one hand, to the walls of the orbital cavity and, on the other, to the sclerotic coat at a distance of from a third to half an inch behind the circumference of the cornea (Fig. 13.30). There are six muscles for each eye, the four recti muscles (internal, external, superior, and inferior) and the two obliques (superior and inferior). The internal rectus muscle turns the eye inward—i.e., toward the nose; the external rectus turns it outward. The superior rectus turns the eye upward and inward, the inferior rectus downward and inward. The superior oblique rotates the eyeball downward and outward, the inferior oblique upward and outward (see diagram, Fig. 13.31). The eye muscles, especially the obliques, also cause a wheel-like rotatory movement.

The superior oblique muscle is supplied by the trochlear (fourth cranial) nerve, and the external rectus by the abducent (sixth cranial)




Fig. 13.27 Illusions of size. A, the vertical lines are of the same length; B, the black circles are the same size.

Fig 13 28 (Above) Illusion of distance The distance from A to B appears to be greater than that from B to C, they are the same

Fig 13 29 (Right) Illusion of size The figure of the man is actually smaller than that of the child, but the effect of distance produced by the converging lines causes it to appear larger



nerve, all the other ocular muscles are supplied by the oculomotor (third cranial) nerve

The muscles of the two eyes act in unison For example, the eyes are turned to one or other side—conjugate deviation—by the contraction of the external rectus of one eye and the internal rectus of the other Convergence of the eyes—ie, turning both eyes inward—is brought about by the contraction of both internal recti In any movements of the eyes, the muscles which bring about the opposite move ment—ie, the antagonistic muscles—are inhibited. Thus the principle of reciprocal inhibition (p. 517) applies to the ocular muscles as well as to other muscles of the body.

The visual axis of the eye is represented by a line extending from an




Fig 13 30 Showing the eyeball in its bony socket (orbital cavity) and the ocular muscles

object in the visual field, through the cornea, a little to the nasal side of its center of curvature, to the fovea centralis When the sight is directed to a far view, the axes of the two eyes are parallel. But during accommodation for near vision, when the eyes converge, the axes meet at a point in the object at which we look. This is called the fixation point. When (as a result of unbalanced action of the ocular muscles) the visual axes do not bear the normal relationship to one another, they may be converged too acutely ("cross eyes") or not enough, or they may diverge. The condition is known as strabismus or squint.




Fig 13 31 Diagram of the actions of the muscles of one eye (From Fuchs after Marquez) OI, infernor oblique; RS, superior rectus, R ext, external rectus; R int, infernal rectus, OS, superior oblique, RI, infernor rectus. The arrows indicate the direction in which each muscle tends to move the eyeball it will be noted, however, that the actions of some muscles are antagonized by those of others. For example, the eye is turned outward by the combined actions of the external rectus, inferior

oblique, and superior oblique. The upward and rotary movement caused by the inferior oblique is neutralized by the opposing action of the superior oblique

The eyelid is raised by a muscle attached to the upper bony wall of the eye socket and inserted into the skin and dense connective tissue of the upper lid It consists of a smooth muscle and a voluntary muscle component. The former is supplied by sympathetic nerve fibers, the latter by a branch of the oculomotor (third cranial) nerve. The muscle is called the levator palpebrae superioris (elevator of the upper lid). The lids are closed by a sphincterlike muscle known as the orbicularis oculi. The fibers of this muscle are arranged in circles and sweep around the opening between the lids. It is supplied by branches of the facial nerve.

The structure and functions of the iris. The iris is a circular diaphragm or curtain lying behind the cornea, and in front of the lens. The circular opening at its center is called the pupil. The iris around its circumference is attached to the ciliary body. It is composed of two sets of smooth muscle fibers. One set of fibers is arranged circularly; this set causes narrowing of the pupil, and is therefore called the sphincter or constrictor pupillae. The fibers of the other set run radially from the pupillary margin; they dilate the pupil and constitute the dilator pupillae muscle (see Fig. 13.32).

610

to blue (see Fig 13 34) For this reason, a series of Iringes, colored from violet to red from within outward, borders the image formed by a simple cheap lens Thus, if a screen (e.g., a white card) were placed at the point designated violet in the figure a colored halo would be thrown upon it with the violet in the center and red on the outside If the screen were at the point indicated by red, the rays having crossed the colors would be in reverse order, red in the center and violet out side If the black and white circles in Figure 13 35 are observed at a distance of about 1 inch from the eye—nearer the eye than the point of clear vision—the white circles will appear in a family blush tint





Fig. 13 35 (Left) Illustrating chromatic aberration (See text)

Fig 13 36 (Center) Diagram showing the structure of an achromatic lens 1, flint glass 2 crown glass

Fig. 13.37 (Right) Showing the contour of the crystalline lens at different periods of life From left to right in infancy in early adult life and in old age.


as a result of chromatic aberration, as the defect under discussion is called It is corrected in camera and microscope lenses by cementing a biconvex lens of crown glass to a concave block of fiint glass (see Fig 13 36). Such a lens is called achromatic. The lens of the eye is not corrected for chromatic aberration. A red and a violet object at the same distance from the eye therefore, cannot both be seen sharply at the same time. If, for example, the eyes are focused upon a violet light, they must accommodate in order to maintain the focus when the light is changed to red. In ordinary vision, the colors surrounding the images on the retina are not perceived largely because we have come to ignore them.

Presbyopia This is a defect of accommodation which develops after middle age. Distant vision is unimpaired but the lens like other its sues, loses its resilience or elasticity with advancing years and cannot in consequence bring the image of a near object into focus upon the retina. The subject of presbyopia is, therefore unable to read ordinary print without the aid of convex glasses.

The shape of the lens also changes with advancing years, its curva

ture being flatter, and its refracting power correspondingly less in middle and late adult life than in the young (see Fig. 13.37). Defects due to abnormalities in the form of the eyeball. The three

Defects due to abnormalities in the form of the eyeball. The three defects about to be discussed are due to abnormalities in the conformation of the eyeball. The normal or emmetropic eye is nearly spherical,



the vertical diameter being only about 0.5 mm shorter than the anteroposterior and transverse (see Fig. 13.38 B).

Myopia or shortsightedness. In this defect of the eye, the refracting power of the lens is usually the same as in the normal eye, but the anteroposterior diameter of the globe is abnormally long. The image is therefore brought to a focus a little in front of the retina. In other words, the lens is too strong for the length of the eyeball. The rays after coming to a focus disperse again, and a blurred image is formed upon the retina, just as when the film of a camera is adjusted at too great a distance from the lens. Myopia is corrected by fitting the subject with concave lenses which, by diverging the rays before they enter

the eye and thus partly offsetting the converging power of the crystal line lens, bring them to a focus upon the retina (Fig. 13 38 A)

When his defect is not fully corrected, the subject of myopia holds any object which he wishes to see clearly very close to his eyes and relaxes his accommodation—that is, accommodates for far vision. The reason is that rays of light from the near object are more divergent,




Fig 13.39 Astigmatism Showing that light rays passing through different meridians are not brought to the same focus *Upper cut* the horizontal curva ture of the lens is the same as the vertical rays of light passing through the horizontal and vertical diameters (meridians) are therefore focused to the same point *Lower* the horizontal curvature is greater than the vertical Rays passing through the horizontal meridian are focused at F<sub>1</sub> and those passing through the vertical Rays (See text)

he thus to some extent corrects his defect. But if he accommodated his eyes for near vision the curvature of the crystalline lens would be in creased and his visual disability accentuated.


Hypermetropia or longsightedness. This defect is due to the anteroposterior diameter of the eyeball being too short. Since the refracting power of the crystalline lens is the same as in the normal eye, the rays cannot be converged acutely enough to form a clear image upon the retina. Hypermetropia is corrected by means of convex lenses which converge the rays before they enter the eye, thus aiding the crystalline lens (Fig. 13.38 C).

Astigmatism This is probably the commonest of all defects of the eye In astigmatism, as the name itself implies (Gk. a not + stigma a point), rays of light are not brought to sharp points upon

the retina, but form, instead, short lines. The defect is present in all eyes to a certain degree, and it is only when well marked that it can be considered abnormal. The stars, for example, which should appear to us as small dots of light, seem to radiate short lines of light. For the same reason, a light shining through the dark appears to emit radiating beams.

It must be remembered that rays of light pass through all meridians of a lens; in converging to a focus they, therefore, form a cone of light, not simply a flat, pennantlike beam. If, as in the normal eye, all meridians of the lens have approximately the same curvature, then rays in all planes are refracted to almost the same degree and come to a focus together. If the curvatures differ, the rays transmitted through a meridian with the greater curvature will be refracted more strongly and brought to a focus in front of rays passing through other meridians. For example, should the vertical meridian be more curved than the horizontal, then, when the rays passing through the vertical meridian are in focus, those in the horizontal will form, not a point, but a horizontal line (see Fig. 13.39). Such inequalities of curvature in the meridians of the cornea or, less commonly, of the crystalline lens, are the cause of astigmatism. The greater curvature may be in either

Fig 13 40 Chart used in testing for astigmatism In the astigmatic eye, the image of the lines in one or more meridians is blurred (not in focus).



a vertical, horizontal, or a diagonal meridian. When the subject of astignatism looks at a card on which is printed a figure like the dial of a clock (Fig. 13.40), the vertical lines may be clearly seen, while the horizontal are blurred, or vice versa. Again, the diagonal lines may be out of focus while the horizontal and vertical ones are clearly defined.

Astigmatism is corrected by the use of spectacle lenses, convex in

the meridian corresponding to that of the cornea (or crystalline lens) having the lesser curvature. Thus, if the curvature of the cornea is greater in the vertical meridian, the subject is fitted with a cylindrical lens having its convexity in the horizontal meridian.

Opacities in the vitreous body Minute opacities or semiopaque bodies are present in the vitreous body of the best of eyes. Though they cause no inconvenience and we are usually unaware of them they can be demonstrated by anyone in his own eyes. They appear when the eyes are closed and one concentrates his attention as he gazes against closed lids or without closing the eyes looks at a clear sky or some blank surface. They are called muscae volitantes (L. flying files). They are seen to move either upward or downward when the eyes are moved but when one attempts to focus upon them they appear to fly away. The visual sensations are caused by the shadows east upon the retina by epithelial cells or other debris—e.g. embryonic remnants in the vitreous body.

Contact lenses These are thin bowl shaped shells of glass made to fit over the cornea they are used in some cases of refractive errors especially those due to an abnormal shape of the cornea to astigma tism or to absence of the crystalline lens. The contact lens is made in the form of the normal cornea and before being applied to the surface of the eye is filled with saline solution which when the lens is in position hese as a thin film between it and the cornea. Now, the saline is of approximately the same refractive power as the cornea and for pur poses of refraction cornea and saline are, therefore, one. Thus the posterior surface of the glass lens being in the form of a normal cornea molds as it were the patient's cornea into normal shape. The lens may



Fig 13 41 Drawing of contact lens also be ground like any other lens so as to render it of the required curvature to correct a refractive error—

e g astigmatism myopia etc (Fig 13 41)

The evil effects of bad lighting upon the eyes Read
ing in a light of such low illumination that the letters
are not seen sharply and read easily too fine a print

or lack of contrast between the letters and the paper cause eyestrain. Our telephone directories seem to suffer from both of the two last mentioned defects. The ciliary muscle contracts to its maximum in order to increase the convexity of the lens and some persons even press upon the cornea with the lids in an effort to bulge it forward—that is to make it more convex and increase its refractive

power Headache results

Glare is also a common cause of eyestrain There are three kinds of

glare—veiling, dazzling, and blinding. Veiling glare is that which is created by a bright light which is uniformly diffused over the scene and reduces the contrast in the image of light and shade. Dazzling glare is caused by a light shining into the eyes at an angle of about 45 degrees. It does not enter into the formation of the retinal image and is scattered by the refracting media of the eye. Blinding glare is light, also not part of the retinal image, which enters the pupil more directly, as when one looks into a very bright light.

The best light for reading is one of from 10 to 20 foot candles, falling upon a dull-surfaced paper, and not causing glare in any of the ways listed above There should also be a generally diffused and indirect illumination of the room.

Blindness, partial or complete. Any loss of transparency of the refracting media of the eye—cornea, aqueous humor, lens, or vitreous body—reduces visual acuity. The parts of the eye most commonly affected in this way are the cornea and the crystalline lens. Destruction, by injury or disease, of the corneal surface is replaced by a scar, which is translucent or opaque, and, if so situated that light rays are prevented from entering the pupil in beams which can be focused, blindness of the affected eye results, in so far as recognizing objects is concerned (though light and shade may be appreciated).

Cataract Cloudiness or opacity of the crystalline lens results in a corresponding limitation of vision Lack of transparency of the lens is called cataract. It is a defect usually occurring in persons past middle age The changes in the transparency of the lens in the usual form of cataract are due to a degenerative process involving the lens proteins which starts near the center of the lens and spreads toward the surface. During this time, the cataract is said to be immature. It is mature when the process has extended to involve the superficial layers of the lens. The cure of cataract may be postponed until the mature stage has been reached The surgeon then resorts to "needling" Needling consists of piercing the anterior chamber of the eyeball, making an opening in the lens capsule, and breaking up the degenerated and softened material This permits the entrance of aqueous humor and the gradual absorption of the lens substance. Then light rays can again reach the retina. The operation for "needling" or discission for cataract goes back to ancient times, but today extraction of the lens in its entirety is the most usual operation. After destruction or removal of the lens, strongly convex glasses are prescribed to compensate for the loss of its refracting power.

4 A foot-candle is a measure of illumination, it is the light given out by a standard candle to 1 square foot of surface 1 foot distant from it.

616

The cause of the ordinary type of the condition, which is known as senile cataract, is thought to be due to the prolonged action of sunlight or other sources of ultraviolet rays upon the lens proteins. It is much more common in India, Egypt, and other tropical and sunny regions than in temperate zones.

Other types of cataract which may occur in earlier life are associated with certain metabolic diseases, especially diabetes. Another type, which occurs in glass blowers, is probably due to heat (infrared) rays.

Total blindness, in which not even a bright light can be perceived.

Total blindness, in which not even a bright light can be perceived, may result from injury or disease of the retinas, of the optic nerves, of the optic tracts, or of the visual areas situated in the cortex of the occipital lobes of the cerebrum. If one retina is affected, or one optic nerve, blindness of the corresponding eye alone results. If the optic ract (i.e., the visual nerve fibers between the optic chiasma and the occipital cortex) or the visual area of the cortex of one side is injured or diseased, partial blindness results in both eyes. Complete blindness results only if the retinas, optic nerves, optic tracts, or visual areas of both sides are involved (see also p. 588).

The pressure within the eyeball; intraocular fluid. The pressure within the eyeball is from 20 to 25 mm Hg during life, but falls to about 10 mm shortly after death The higher pressure is due to the pressure in the capillaries in the wall of the globe. The lower pressure of 10 mm which persists for a time after death is due to the pressure of the aqueous humor and the fluid in the vitreous body.

The intraocular fluids are not secreted, apparently, but are formed by the passage of fluid by dialysis (p. 18) from the blood across the membranes on the surface of the ciliary body and the posterior aspect of the iris The fluid is reabsorbed, mainly by draunage through canals at the angles of the anterior chamber formed by the iris and cornea (see Fig. 13 1). There is thus a fairly free circulation of fluid—from blood to the interior of the eyeball and back to the blood by reabsorption.

sorption.

Glaucoma. This is a very serious disease of the eye of which the initial abnormality is an increase in the intraocular pressure. The ocular hypertension leads to pathological changes within the eyeball which, if the tension persists, lead to blindness through pressure upon the delicate retinal structures. One form of the disease follows some other eye disease and is due either to increased production of intraocular fluid, or to blockage of the drainage canals at the angles of the anterior chamber. Another form of the disease is not preceded by any known ocular disease; its cause remains obscure.

## The Ear

The structure of the ear. The ear consists of three distinct parts, called the *outer*, *middle*, and *inner ears*, each of which has a special part of play in the mechanism of hearing.

The outer ear consists of an appendage of cartilage and skin situated on the side of the head and known as the auricle or punna, together with a short funnel-shaped passage called the external auditory meatus (or canal). The latter leads into the temporal bone of the skull, and is closed at its inner end by a flexible membrane called the drum or tympanic membrane. The auricle and auditory canal serve simply to col-




Fig. 13 42 Plan of the ear.

lect and direct the sound waves to the drum membrane; the latter separates the outer from the middle ear (see Fig. 13.42)

The middle ear, drum, or tympanum is a small chamber within the temporal bone. As just mentioned, the drum membrane is interposed between it and the outer ear. The middle ear contains three miniature bones (the auditory ossicles) named individually from their shapes, the mallets (hammer), the incus (anvil), and the stapes (stirrup) (see Fig. 13.43). They stretch across the cavity of the middle ear from the tympanic membrane to the inner wall of this chamber, and are joined to one another by minute joints. The handle of the malleus is attached to the tympanic membrane; its head articulates with the body of the incus A process of the incus is joined to one end of the stapes; the other end or footplate of the stirruplike bone is fitted into a small

61R

opening in the inner wall of the middle ear, called the oval window. Situated a little lower in the inner wall of the tympanum is the round window; this is closed by a thin membrane.

A narrow canal—the auditory or Eustachian tube—which connects the middle ear with the posterior part of the nose, serves for the passage of air between the middle ear and the atmosphere. Thus equality




Fig 13 43 The auditory ossicles.

of air pressures upon the two sides of the drum membrane is maintained

Two very small muscles are to be found in the middle ear. They arise from neighboring bone or cartilage and are inserted, one (the tensor tympani) into the handle of the malleus, the other (the stapedius) into the stapes. These two muscles contract reflexly and simultaneously when very forcible vibrations strike the eardrum, and by damping the movements of the ossicles protect the internal ear from injury.

The inner ear or labyrinth lies internal to the middle ear; it contains the organs of hearing and equilibrium. The auditory sense organ is held within and on the outer wall of a spiral tunnel in the temporal bone named the cochica—so-called from its resemblance to a snail's

shell (L cochlea a snails shell) The spiral canal makes two and three quarter turns around the central column of spongy bone called the modiolus (Figs 13 44 and 13 52 p 631) The larger turns of the spiral are at the base of the cochlea the smaller turns at the apex or cupola A shelf or ledge of bone which projects from the modiolus and winds around it like the thread of a screw divides the cochlea moon




Fig 13 44 Upper drawing a view of the bony cochlea divided vertically through the m ddle lower drawing enlarged drawing of one turn of the duct of the cochlea a vestibular membrane b bas lar membrane c organ of Cort d acoustic press.

pletely into two spiral channels. But in the living state the division of the cochlea is completed by a membranous septum called the basilar membrane which extends from the tip of the bony shelf mentioned above to the outer wall of the cochlea. A second more delicate septum called the vestibular membrane proceeds from the upper surface of the shelf of bone to the outer wall where it is fixed a short distance above the outer attachment of the basilar membrane. The spiral tube enclosed between these two membranes is called the membranous cochlea or duct of the cochlea? The part of the cochlea lying above

a The cochlea is here described as though placed upon its base but actually it les upon its sde with its base of cered inward toward the cranial cavity and its apex or cupola pointing outward and forward.

620

this membranous tube is named the scala vestibuli, and the part lying below, the scala tympani. The duct of the cochlea is sometimes referred to by its older name, scala media. The spiral compartments were given these names from their fancied resemblance to stairways (L. scala, a stairway). The scala vestibuli and the scala tympani are filled with a fluid called perlymph, which is identical in composition to, and continuous with, the cerebrospinal fluid. The duct of the coch lea is filled with endolymph.



Fig 13 45 The internal auditory meatus

The scala vestibuli communicates with the middle ear through the oval window which, as already stated, lodges the footplate of the stapes

The membrane of the round window is interposed between the commencement of the scala tympan; and the middle ear

Resting upon the basilar membrane is a structure consisting of specialized cells of different sizes and shapes, and known as the spiral organ or organ of Corti. It and the basilar membrane follow a spiral course throughout the entire length of the duct of the cochlea. The specialized cells are the auditory receptors, they are clongated elements from the summits of which hairlike processes project. They are therefore called hair cells. The tips of the hairs are attached to, or embedded in, a thin elastic membrane—the tectorial membrane—which extends outward from near the attachment of the vestibular areas that a special content of continuous contents.

membrane and, floating in the endolymph, overlies the organ of Corti.

The internal auditory meature is a short, bony canal, about I cm long, which runs inward from the internal ear to open into the eranial cavity It lodges the cochlear and vestibular nerves. The outer end or bottom of the canal (Fig. 13 45) is occupied by the base of the modiolus and a perforated plate of bone. From the former, central fila-

ments of the cochlear nerve enter the meatus (see below); the perforated plate gives passage to the peripheral fibers of the vestibular nerve (p. 631).

The auditory pathway. The nerve of hearing is known as the cochlear nerve. Its fibers are the processes of nerve cells in a small ganglion called the spiral ganglion of Corti, which occupies the spiral canal that tunnels through the center of the modiolus. Processes of the ganglion cells are in two sets—peripheral and central. The former pass along the spiral ledge of bone to reach the organ of Corti. They terminate as minute filaments around the bases of the hair cells—the receptors of hearing. The other processes, which arise from the opposite poles of the ganglion cells, pass centrally with the vestibular nerve




Fig. 13.46 Diagram of the auditory pathway.

in the internal auditory meatus to enter the medulla oblongata; here they connect with groups of nerve cells (cochlear nuclei). Secondary neurons ascend from here to the midbrain (medial geniculate bodies); from there a third set of neurons passes to the auditory center (see Fig. 13 46) in the cortex of the temporal lobe of the cerebrum.

Only the auditory part of the labyrinth has been described in the foregoing paragraphs. The inner ear contains, also, the semicircular canals and otolithic organs. These, which have no auditory function, will be described later.

622

The physiology of hearing. The structure of the auditory apparatus having been sketched, we are in a position to understand the manner in which the energy of sound waves is transmitted to the auditory receptors. The outer and middle ears serve merely to transmit the sound waves to the inner ear in which the receptors are contained. The waves set up in the air by a sounding body strike the ear drum and set it into vibration. In man, the pinna and the somewhat funnel-like shape of the auditory meatus have only a slight effect in concentrating the sound waves upon the drum membrane. In many animals, however, the pinna serves an important function in this regard, being a large funnel-shaped structure which can be turned toward the source of the sound.

The vibrations set up in the tympanic membrane are transmitted across the middle ear by the auditory ossicles. The head of the malleus





Fig 13 47 Diagram of the passages of the cochlea straightened out to show the manner in which vibrations are transmitted from the oval to the round window through the scala media. The wavy line represents the organ of Corti

is connected to the body of the incus in such a way that a movement inward or outward of the tympanic membrane causes a corresponding movement of the handle of the malleus, of the long process of the incus, and of the footplate of the stapes, lodged in the oval window between the middle ear and the scala vestibuli of the cochlea. Thus movements of the tympanic membrane are transmitted through the ossicles to the inner ear. The series of inward and outward movements of the footplate of the stapes, acting like a plunger in the oval window, sets up corresponding movements in the fluid contents of the scala vestibuli. These movements are transmitted freely through the vestibular membrane to the fluid in the cochlear duct, and through the basilar membrane, in turn, to the scala tympani (Fig 13 47). Thus, each time the footplate of the stapes is pressed into the oval window, the basilar membrane moves toward the scala tympani and pulls the processes of the hair cells fixed to the tectorial membrane; when

the stapes moves outward, the tension on the hair cells is relieved. Thus, by a series of mechanical stimuli, impulses are set up in the terminals of the auditory nerve. The round window, situated between the scala tympani and the middle ear, permits free movements of the contents of the inner ear. Since liquids are incompressible, no movement of the stapes could occur, of course, were there no other part of the bony wall of the cochlea which yielded to pressure.

The auditory ossicles are connected to one another in such a way as to constitute a small lever (see Fig. 13.48). The handle of the mal-

Fig. 13.48 A mechanical model illustrating the transmission of sound waves through the middle ear. The lever represents the chain of auditory ossicles.



leus, attached to the tympani membrane, forms the long arm of the lever; the long process of the incus is the short arm. The long arm is one and a half times the length of the short arm. The movement of the long process of the incus (and of the stapes with which it connects) is, therefore, only two thirds as great as that of the tympanic membrane. We should expect, on the other hand, that the force exerted by the stapes at the oval window would be one and a half times greater than that applied to the handle of the malleus. But, on account of a large proportion of the energy being lost in transmission (due to inertia of the ossicles and friction of the air surrounding them), only about half of the force exerted upon the handle of the malleus reaches the oval window. However, the area of the tympanic membrane is nearly twenty times that of the oval window; this fact alone, were there no loss of energy in transmission, would cause the force concentrated at the oval window to be increased twentyfold over that of the vibrations of the drum membrane. Now, as just mentioned, the force transmitted to the stapes is only one half of that applied to the malleus; the net magnification of the force exerted by the movements of the footplate of the stapes, brought about through the mechanism of the middle ear, is, therefore, about ten

The appreciation of pitch (discrimination of the frequency of the sound waves). Sound is transmitted from a vibrating body to the ear as a series of alternating condensations and rarefactions of the atmosphere The disturbance in the air travels as a succession of waves A complete wave—that is, from the crest or trough of one wave to the corresponding phase of the other—is referred to as a double vibration (dv) or a cycle The pitch of a tone depends upon the frequency of the waves—that is, upon the number of double vibrations or cycles reaching the ear per second For example, a tuning fork which vibrates at a frequency of 100 dv per second emits a note of higher pitch (by an

a frequency of 100 dv per second emits a note of higher pitch (by an octave) than one vibrating at 50 dv per second

Sounds ranging in frequency from 40 to 20,000 dv per second are audible to the human ear. The range of audible frequencies varies, however, between individuals and between different species of ani mals. The eat, for example, can hear the very high pitched sounds made by a mouse, which are usually quite maudible to human ears. The bat also makes a cry which of course, can be heard by its kind, and probably also by other animals, but which is of a pitch too high to make an impression upon the human ear

There has been no little discussion as to the mechanism whereby variations in the frequency of the sound waves striking the ear are converted into messages which the brain interprets as differences in pitch. The two classical theories are the resonance theory of Helmholitz. and the telephone theory of Rutherford

The resonance theory states that the basilar membrane, like the wires of a piano is a scale of resonators. If the dampers of a piano are raised and a note sung near it, a tone of the same pitch is given out by the instrument Those wires of the piano alone vibrate which, if struck, would themselves have emitted the same note This well known phe nomenon is called sympathetic resonance. The basilar membrane con tains some 24,000 fibers. These so called auditory strings increase pro gressively in length from the base to the apex of the cochlea Each fiber, or a group of fibers of a given length, vibrates in unison, it is supposed, with vibrations of a particular frequency transmitted to them from the drum membrane, the short tense fibers at the base of the cochlea vibrate to high notes, and the longer, more lax fibers at the apex to low notes. Of course, only those hart cells overlying the fibers which are set into sympathetic vibration are stimulated.

The resonance theory, therefore, claims that the analysis of sound.

is a function of the cochlea, the particular region of the latter from which the nerve impulses are discharged being the sole basis upon which the brain rests its faculty of pitch discrimination. The resonance theory is supported by a number of facts. (a) When the ears of animals (e.g., guinea pigs and rabbits) are subjected to prolonged stimulation by a loud high- or low-pitched tone, degenerative changes are produced in the basal or the apical portion, respectively, of the organ of Corti. (b) Boiler-makers or others who work among clanging noises sometimes become deaf to high tones (boiler-maker's disease), the organ of Corti in the basal turns of the cochlea is found to have degenerated. (c) Destruction of the apical part of the cochlea in animals results in deafness to tones of low frequency; injury to the basal part causes deafness to high tones. (d) The structure of the cochlear duct with its basilar membrane strongly suggests that it acts as a resonator, and recent experimental work has provided direct evidence of such a function.

When the ear is stimulated by a sound, the vibrations of the organ of Corti set up rhythmical changes in electrical potential within the cochlea (Wever and Bray effect). When amplified and recorded graphically, these are found to have exactly the same frequency as that of the sound. Indeed, if the electric currents are conducted to a loud speaker the original sound is faithfully reproduced. With low notes, the potential changes are greatest in the apical turns of the cochlea; with high notes, in the basal turns, thus demonstrating clearly the resonating function of the cochlea

But the electrical effects do not represent nerve impulses originating in the terminals of the cochlear nerve; they are due to movements with consequent pressure changes in the basilar membrane and organ of Corti set up by the sound waves. The mechanical energy (pressure) is transformed to electric energy—a piezoelectric effect (Gk. piezo, I press)

The telephone or frequency theory. In a telephone system the transmitting instrument converts the sound waves into electrical impulses of the same frequency. Upon reaching the receiver, the electric impulses set up vibrations which reproduce the original sound. The telephone theory of hearing postulates that the basilar membrane does not vibrate selectively in the manner demanded by the resonance theory, but as a whole, in unison with the vibrations transmitted to it from the drum membrane. The frequency of the impulses caused by stimulating the hair cells corresponds with that of the vibrations of the basilar membrane, and, consequently, with that of the sound waves.

The telephone theory claims, therefore, that the cochlea possesses no power of sound analysis, and that such is purely a function of the brain, the frequency of the unpulses received by the auditory center being the basis for the discrimination of pitch


The telephone theory as originally proposed is faced with the object

tion that the maximum frequency at which impulses can be transmit ted by a nerve fiber is about 1000 per second, whereas, sounds with frequencies up to 20,000 dv per second are audible. But the auditory nerve consists, of course, of a large number of fibers, and it has been proposed that the impulses are discharged over the separate fibers, not synchronously, but in a scattered volley That is, some fibers are in their refractory phase (p 574) while others are conducting Thus it is possible for a group of fibers to transmit impulses in "broken step at frequencies as high as, or even higher than 20,000 per second Nevertheless, the volley theory, as this amplification of the telephone theory is called, does not fit all the facts, it fails to explain the appreciation of tones having a vibration frequency higher than about 3000 per second For example, the action potentials set up by a sound and recorded (p 469) from the auditory tracts in the brain have exactly the same frequency as that of the sound up to 1000 dv per second Be tween 1000 and 2000 dv per second, one impulse occurs for every two sound vibrations, and one for every three between 2000 and 3000 dv per second, but at higher frequencies (between 3000 and 20,000 dv per second) there is no relationship As a general theory of pitch discrimination, the telephone or fre quency theory is unsatisfactory. The resonance or harp theory is now generally accepted To summarize The cross fibers or "strings" of the basilar membrane, according to the principle of sympathetic reso nance, vibrate at different levels (base to apex) of the cochlea as de termined by the frequency of the sound waves reaching the internal

As a general theory of pitch discrimination, the telephone or frequency theory is unsatisfactory. The resonance or harp theory is now generally accepted. To summarize The cross fibers or "strings" of the basilar membrane, according to the principle of sympathetic resonance, vibrate at different levels (base to apex) of the cochlea as determined by the frequency of the sound waves reaching the internal ear. Thus, the short tense fibers in the basal turns whrate in unison or "sympathy" with sound waves of high frequency (high pitched or treble tones), the longer fibers at the apex respond to sound waves of low frequency (low pitched or bass tones). The basilar fibers between these two extremes vibrate to tones of intermediate pitch from treble to bass. The vibrations of the fibers at any particular level of the basilar membrane stimulate the auditory receptors, and thus set up impulses in the nerve terminals in the hair cells at that place alone. The mod erri resonance theory is, therefore, also spoken of as the place theory just as we know at once the precise spot on the skin which is stimulated when we feel a touch at that spot, so we are able to discriminate

THE EAR

between impulses arising from the different levels of the basilar membrane. But the actual stimulated place in the basilar membrane is not perceived as we perceive the position of a stimulated point on the skin. The different levels of stimulation are interpreted by the brain as differences in pitch. Such localization is possible because each and every level of the cochlea is connected by nerve fibers with a particular group of nerve cells arranged in regular order in the auditory center—



that is in the cortex of the temporal lobe of the cerebrum (see Fig 13 49) In other words each point upon the basilar membrane is rep resented by a corresponding point in the cerebral cortex

Intensity and loudness of sound. The power transmitted through the atmosphere from a sounding body is referred to as the intensity of the sound Sound intensity is expressed as the quantity of energy, in microwatts, passing per second through an area of 1 quare centimeter. The greater the intensity of the sound, the greater, of course will be the amplitude of the movements of the drum membrane and generally speaking, the louder will be the sound. But the ear does not possess the same sensitivity at all frequencies. Within the range of from 1000 to 2000 dv per second the lowest intensity (intensity threshold) is required to arouse an auditory sensation, and a sound of any given intensity is loudest. The sensitivity of hearing—i.e. the loudness of a sound of a given intensity—diminishes progressively as the frequency

is increased above 2000 dv per second or reduced below 1000 dv per second (see Fig 13 50). In other words the threshold of hearing is lowest for sounds with frequencies around 2000 dv per second. It will be appreciated from these remarks that intensity and loudness are no synonymous terms. The intensity of sound is an absolute physical value whereas the loudness of sound is a matter of auditory perception.

As the intensity of the sound increases other sense organs than those of hearing are stimulated. We feel the sound and if very intense it arouses the sensation of pain. For example, the report of a rifle fired




Fig 13.50 Chart showing the thresholds of hearing and feeling of sound at different frequencies (After Wegel)

at a distance is heard but at short range is felt as well and if too close to the ear causes pain. It will be seen that the chart in Figure 13.50 that in contrast to that of hearing the threshold of feeling is highest at frequencies around 500 dv per second.

Detective hearing, deathiess. Deathiess is a rather indefinite term but is usually taken to mean defective hearing of such a degree that ordinary conversation cannot be heard distinctly and without difficulty. It varies greatly in degree from duliness of hearing the subject being unable to recognize words spoken by another in his usual voice to complete loss of hearing. As a rule the hearing defect is not equal for tones of every pitch. In some types of deathiess, bearing loss affects mainly or solely the higher sound frequencies in others the lower frequencies.

Classification of dealness. There are three main types of dealness called respectively transmission dealness perceptive or nerve deal ness and central dealness. Any condition of the external or middle ear which prevents the sound waves from being transmitted without interference is called transmission dealness. Thus inflammation with de struction of the structures of the middle ear plugging of the outer ear

with wax, or closure of the auditory tube, will cause some degree of deafness. Fixation of the footplate of the stapes in the oval window by hardened tissue blocks the transmission of the sound waves to the cochlea and is also a frequent cause of deafness. When the auditory tube is blocked, the middle ear becomes a hermetically sealed cavity. The air within it soon becomes absorbed and, as a consequence, the pressure on the inner side of the drum membrane is much less than that of the atmosphere. The membrane is pushed inward against the inner bony wall of the middle ear and, therefore, cannot vibrate freely. Temporary blockage of the auditory tube often occurs with a cold, as a result of swelling of the lining of its mucous membrane. Inequality of pressure on the two sides of the drum membrane is brought about also, in a somewhat different way, in airplane descents. The middle ear is filled with air at the lower pressure of the atmosphere at the high altitude; when the plane reaches the ground, the atmospheric pressure at this level, being much higher, pushes the drum membrane inward. This event is usually guarded against by chewing gum during the descent which, since it stimulates salivary secretion, induces swallowing. Swallowing, in turn, opens the mouths of the auditory tubes in the pharynx and permits air to enter the middle ear from time to time, and thus equal pressures are maintained on the two sides of the drum membrane.

If descent from a high altitude is made too rapidly, and swallowing fails to allow air to enter the auditory tubes, rupture of one or of both eardrums may occur; the higher outside (atmospheric) pressure would be unable to force air upward into the middle ears through the closed pharyngeal ends of the tubes, and thus to equalize the pressures on the inner and outer sides of the tympanic membranes.

In rapid airplane ascents, on the other hand, the higher pressure within the middle ear is reduced from time to time by the air forcing open the sphincter muscle around the opening of the auditory tube. In other words, the sphincter can be forced from within but not from without.

Perception deafness is due to disease or injury of the cochlear structures or of the cochlear nerve itself.

Central deafness is caused by a lesion—e.g., a tumor, abscess, or injury of the auditory tracts or of the auditory cortical centers.

Testing defective hearing. Hearing is tested (in a room as nearly silent as possible) by means of a series of tuning forks of graded vibration frequencies held at varying distances from the subject's ear. Or an instrument known as an audiometer is used, by means of which ለገበ

tones are produced electrically which can be very precisely varied as to pitch and intensity. The results are plotted on graph paper similar to that shown in Figure 13.51. Such a record is called an *audiogram*. In this instance, hearing is defective only over a limited range of sound frequencies

The nonauditory part of the labyrinth—the organ of equilibrium. The inner ear contains another organ besides the cochlea, consisting of three semicircular canals and two small sacs, called, respectively,




Fig 13 51 An audiogram The shaded area indicates the area over which hearing is defective

the utricle and the saccule. The semicircular canals and the utricle have no auditory function, being concerned solely with the maintenance of the equilibrium of the body (see Fig. 13.52).

The semicircular canals are membranous tubes filled with fluid known as the endolymph. They are contained within semi-circular tunnels in the temporal bone—the osseous canals—but

separated from the bone by perlymph. The three canals lie approximately at right angles to one another, one in each of the three dimensions of space. Two canals are vertical, one horizontal. The two vertical canals (superior and posterior) are placed diagonally in relation to the anteroposterior diameter of the skull; the convexity of one is directed outward and backward, of the other outward and forward (see Figs. 13.52 and 13.53), and that of the horizontal (lateral) canal, outward One end of each canal is dilated into a fusiform swelling called the ampulla. A small elevation, called the crista, is situated in each of the three ampullae. This is composed of cells with bristlelike processes—the hair cells—surmounted by a cap of gelatinous material called the cupula. The cristae are the sense organs (proprioceptors) of the canals. Both ends of each canal open into the utricle.

The utricle and saccule are oval membranous sacs measuring (in man) about \(^1\) inch in their longest diameters. They are contained within an oval cavity in the bone, called the vestibule, situated between the cochlea and the semicircular canals. The utricle, as just mentioned, communicates with the semicircular canals, the saccule connects through a narrow canal with the duet of the cochlea. The utricle and saccule are indirectly connected through a narrow Y-shaped tube

(Fig 13 52) The entire membranous labyrinth, auditory and non auditory, thus forms a continuous system of communicating passages and chambers The utricle and saccule each contains a sense organ called the macula, this is a plaque of hair cells covered by a layer of a gelatinous substance Adherent to the latter are a number of crystals of




Fig 13 52 The membranous labyrinth

carbonate of lume, known as otoluths (ear stones) or otoconua (ear dust)

The pathway for impulses from the semicircular canals Impulses from this part of the labyrinth are carried by fibers of the vestibular nerve—the other division of the acoustic nerve. Its fibers, like those of the cochlear nerve, are in two sets, peripheral and central, and are processes of cells in the vestibular (Scarpa's) ganglion which forms a swelling on the nerve at the bottom of the internal auditory meatus. The peripheral set of fibers ends around the bases of the sensitive hair cells of the semicircular canals, utricle, and saccule. The central processes (fibers) of the ganglion cells connect with groups of cells (ves-





Fig. 13.53 Showing the positions of the semicircular canals in relation to the vertical planes (anteroposterior and transverse) of the skull. 1, horizontal canals; 2, posterior vertical canals; 3, anterior vertical canals

tibular nuclei) in the medulla oblongata. Some of these secondary neurons carry the impulses to the cerebellum; other neurons convey the impulses to the anterior horn cells of the spinal cord (see Fig 13.54). The axons of the latter constitute the vestibulospinal tract (p. 528).

The functions of the semicircular canals. If a frog is rotated rapidly on a turntable, the limbs on the side toward which the movement is made are extended, the limbs of the opposite side are flexed, and the head turned toward this side (see Fig. 13.55). Thus displacement of the body is prevented If the table is tulted forward, the animal avoids being thrown upon its face by extention of its forelimbs; a backward tilt causes extension of the hind limbs; and tilting laterally is followed by extension of the limbs on the corresponding side, accompanied by CORTEX OF




flexion of the opposite pair of limbs. These reflex reactions, whereby the animal resists being upset from its "normal" position by some unusual movement, are the result of stimulation of the end organs of the semicircular canals. Impulses initiated in the cristae are transmitted to the vestibular nuclei in the medulla oblongata, and thence relayed to the cerebellum and, by the vestibulospinal tracts, to the cord. The reactions resulting from rotation in the horizontal plane are due to stimulation of the horizontal pair of canals: those caused by titing forward or less than the stimulation of the horizontal pair of canals: those caused by titing forward or less than the stimulation of the horizontal pair of canals: those caused by titing forward or less than the stimulation of the horizontal pair of canals: those caused by titing forward or less than the stimulation of the horizontal pair of canals those caused by titing forward or less than the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the horizontal pair of canals that the stimulation of the stimulation of the stimulation of the horizontal pair of canals that the stimulation of the s



Fig 13.55 Position taken up by a frog during rotation to the left. (After Ewald.)

stimulation of the horizontal pair of cato the left. (After Ewald.)
rals; those caused by tilting forward or backward, or laterally (i.e.,
by a rotary movement in the plane, approximately, of one or other
pair of vertical canals), are due to stimulation of the cristae of the
corresponding canals. The movement causes a mechanical stimulus to
be applied to the hair cells of the cristae. When the head is rotated
in the plane of one or other pair of canals, the endolymph, owing to
its inertia, exerts a momentary pressure which bends the hair cells or
the canal on the side away from which the rotation is made. For the
same reason, the pressure upon the hair cells of the canal of the opposite ear is reduced. The effects upon the two ears, though opposite ir
nature, act conjointly in initiating the reflex muscular movement.

Fig. 13.56 Showing endolymph movement during and after rotation. Upper drawing, during rotation, large arrows indicate the direction of rotation; small arrows, the endolymph movement. Lower drawing, after rotation. (After Best and Taylor, The Physiological Baits of Medical Practice.)



Stimulation of the semicircular canals also causes movements of the eyes When, for example, a person is rotated rapidly with the head erect—i.e., approximately in the plane of the horizontal canals—the eyes are observed to make a series of rhythmical side to side movements after the rotation has ceased These to and fro movements, which are termed nystagmus, are due to the momentum of the endo-lymph That is, the endolymph movement continues for an instant after the rotation of the head has ceased, and thus exerts pressure upon the crista of the canal on one side (see Fig 13 56); the pressure on the hair cells of the canal of the opposite ear is diminished. It is clear that the effects following rotation will be the reverse of those occurring at the commencement of rotation If the head is bent forward, backward, or onto one shoulder during the rotation, the vertical canals are stimulated The nystagmus which follows rotation in these positions of the head is not horizontal, the eyes move either up and down (vertical nystagmus) or show a rotary movement (rotary nystagmus), according to which particular pair of vertical canals was in the plane of the rotation

Other effects follow stimulation of the semicircular canals by a rapid rotary motion Everyone is familiar with the sensation of dizziness, or vertigo, and the staggering gait which result from spinning (stimula tion of horizontal canals) A more pronounced disturbance of the equi librium sense follows stimulation of the vertical canals, this may be demonstrated simply upon oneself by bending over, so as to bring the head and neck into the horizontal plane, and then circling a few times around a mark on the ground Impulses set up in the canals may travel over efferents of the autonomic nervous system causing such reflex effects as nausea and vomiting, dilatation of the pupil, sweating, pallor. and a fall in blood pressure

The hair cells, since they are stimulated by rotary motion, are called rotary receptors. It should be emphasized that the cristae are stimulated only as a result of the inertia or the momentum of the endolymph—that is at the commencement or the termination of the rotation. The hair cells are not bent after the rotary movement is well started, for then the mertia of the fluid has been overcome and moves with the wall of the canal Means of detecting a continuous rotary motion have not been evolved, for we are not, except perhaps rarely and under very unusual circumstances, ever exposed to it.


The functions of the utricle. Whereas the receptors of the semicir cular canals respond to movements of the head, the position of the

We must, of course, exclude from this statement the rotation of the earth

head is the factor responsible for excitation of the hair cells of the utricular maculae. The canals are organs of kinetic sense, the utricles of static sense. Two types of equilibrium reactions—the righting and attitudinal reflexes—are dependent upon the utricles.

The righting reflexes. When an animal is placed upon its back, it immediately rights itself. The head is first brought into its "normal" position in space by a contraction of the neck muscles, the body follows,

Fig. 13 57 The positions of an animal's head, each marked with the angle which the mouth cleft makes with the horizontal plane. (Drawn from photographs after Magnus.)



being brought into its normal relationship with the head by contractions of the trunk and limb muscles. The movement of the neck muscles, whereby the head is righted, is dependent upon the sense organ
of the utricle; it is, therefore, called the labyrinthine righting reflex;
the center for this reflex is in the midbrain. When the head is in the
wrong position, mouth looking upward, the otoliths adhering to the
maculae hang downward; the tension which their weight exerts upon
the hair cells acts as a stimulus to the sensitive hair cells. The movement of the neck stimulates, in turn, proprioceptors in the neck muscles, through which the reflex righting movements of the trunk and
limbs are initiated. These movements of the trunk and limb muscles,
caused by stimulation of proprioceptors in the neck muscles, are termed
neck reflexes. Their center is situated in the upper cervical segments
of the spinal cord.

The importance of the righting reflexes in man is evident if one con-

siders how a swimmer, after diving into deep water, orients himself without difficulty and rises to the surface. The other senses (e.g., vision or touch) can give him little or no information of his position in space. He relies upon his utricles to bring his head "right side up." Deaf mutes, whose labyrinths as a whole are not developed, run a grave risk, even though that he code divingers. If placed into dear water was

though they be good swimmers, if plunged into deep water.

Attitudinal reflexes. When a decerebrate animal, with its neck immobilized in a plaster cast (in order to exclude the neck reflexes) is placed in different positions, variations in the tone of the limb muscles will be observed. When the animal is placed back down with the cleft of the mouth at an angle of 45 degrees above the horizontal plane, the tone of the muscles of all four limbs is maximal; in the opposite position (feet down with mouth cleft at 45 degrees below the horizontal plane), the tone of the muscles is minimal. In positions of the head between these two extremes, gradations in muscle tone result, in accordance with the position of the head in relation to the horizontal plane (Fig. 13.57). These tonus reactions are the result of variations in the activity of the utricular proprioceptors (maculae). The sense organs are stimulated most strongly when the otoliths are hanging from the maculae—i.e. pulling upon the hairs, as when the animal is on its back or on its feet with snout pointing upward (neck extended); when the otoliths are resting upon the maculae—i.e., with snout pointing downward (neck flexed)—the hair cells are not stimulated.

The tone of the lumb muscles is also influenced reflexly by movements of the neck muscles. When the neck is extended, the tone of the extensor muscles of the forelimbs is increased, while the extensor tone of the hind limbs is reduced Flexion of the neck causes the reverse effects—namely, decreased tone of the forelimb extensors together with increased tone of the extensors of the hind limbs. The neck reflexes are studied best after the vestibular nerves have been severed and the utricular reflexes thus abolished

Though the attitudinal reflexes are most readily demonstrated in the decerebrate preparation, they constitute an important mechanism controlling the posture of the normal animal. Otolithic and neck reflexes cooperate to facilitate various postural adjustments. When a cat, for example, turns its head to look upward at a piece of food on a shell, the cololithic organ is stimulated with a consequent tendency for the extensor tone of all four limbs to increase. The neck reflex initiated by the extension of the neck reinforces the otolithic influence on the fore-limbs; but the effect of the neck movement upon the hind limbs is the reverse of that of the utricle. The influence of the neck movement upon




Fig 13.58 Drawings of a normal cat, showing the animal's posture (above) when its attention is attracted by an object placed above it. Photograph of the same animal (below) when its attention is drawn to an object below it. The difference between the two positions of the forelimbs is very marked, because in them the neck and labyrinthine reflexes reinforce one another, the difference in the position of the hind limbs is not so great, since the two sets of reflexes oppose one another. (Drawn from photographs after Magnus)

the hind limbs, however, predominates so that the animal assumes a sitting posture with forelimbs extended and hind limbs flexed. On the other hand, when the animal lowers its head to look beneath a cupboard, the otolithic influence reduces the extensor tone in all four limbs; but the neck movement, though it also reduces the extensor tone of the forelimbs and thus acts in conjunction with the labyrinthine effect, antagonizes and predominates over the labyrinthine influence upon the hind limbs. The animal, therefore, assumes an attitude with flexed forelimbs and extended hind limbs-an appropriate posture of the body for that particular position of the head (see Fig. 13 58)

The hair cells of the utricular maculae since they are stimulated by the weight and consequently the pull of the otoliths are called gravity receptors. Seasickness and other forms of motion sickness are due to the intermittent and erratic stimulation of these receptors by the sudden changes in the positions of the head

### Taste

Smell and taste are chemical senses-that is the receptors (chemo receptors) respond to chemical stimuli. In order for a substance to arouse the sensation of taste it must first be dissolved-either taken in solution or dissolved in the saliva, a solid placed in a perfectly dry mouth cannot be tasted

The organs and nerves of taste. The anterior two thirds of the upper surface (dorsum) of the tongue is beset with minute projections of the mucous membrane called papillae The papillae at the edges tip and more anterior part of the dorsum of the tongue are very small conical cylindrical or mushroom shaped structures. They give a velvety char acter to this part of the lingual mucosa. In accordance with their form they are called filitorm or functiorm napillae. The more posterior




Fig 13 59 A taste bud

part of the tongue's surface is rougher owing to the presence of much larger papillae These are of peculiar construction each is surrounded by a groove or trench the whole structure resembling a squat little tower within a moat They are there fore called vallate papillae (L vallum a rampart) (see Pl 15A)


Embedded in the covering

of the papillae (both large and small types) are groups of slender cells provided with hairlike processes packed lengthwise into bundles. The cells are the receptors of taste the bundles which they compose are called taste buds (Figs 13 59 and 13 60) Each cell receives a fila ment from one of the nerves of taste The taste bud opens upon the surface of the papilla through a small pore The ends of the cells con verge toward this point, where their processes become massed together

Substances in solution enter the pores and act as chemical stimuli. A few scattered taste buds are present on the extreme posterior part of the tongue and even in the mucosa of the epiglottis. These later receive fibers from the vagus nerve.

The fundamental sensations of taste are four in number—sweet, bitter, sour, and salty; two others—namely, alkaline and metallic tastes—are sometimes included with these. The various other tastes which we experience are due (a) to the blending of some one or other of the fundamental sensations, or (b) to the combination of the latter with sensations caused by the stimulation of the ordinary nerves of the

Fig 13.60 Section of a vallate papilla of the tongue. A taste bud is shown enlarged in Figure 13.59.



mouth. For example, ginger is recognized not only by its actual taste (i.e., by the stimulation of the taste buds), but from the burning sensation which results from excitation of the ordinary sensory nerves of the mouth. Oils are unpleasant to take, largely because of their "feel." Acetic and many other acids, as well as having a sour taste, give rise to an astringent or burning sensation which is confused in consciousness with the sense of taste and blended with it.

Many of the finer flavors are in reality sensations of smell (p. 640), and smell enters largely into the many sensations which we attribute generally to taste. For this reason, when the nose is held or the nasal mucous membrane is inflamed, as by an ordinary cold, the sense of taste is blunted. On the other hand, certain substances which we think we detect by smell are actually recognized by the sense of taste. The sweetish smell of chloroform is an example, the vapor reaching the taste buds in the inspired air.

The four fundamental taste sensations are not aroused with equal intensity over all parts of the surface of the tongue. Each type of taste sensation is served by its own kind of taste bud. Taste receptors sensitive to sweetness and to saltiness are most numerous at the tip and forepart of the tongue, whereas those responding to sourness are found along the edges (Pl. 15A). The taste buds causing a bitter sensation

are scattered over the back of the tongue and epiglottis. A bitter sweet substance (such as sodium salicylate), when first taken into the mouth tastes sweet the bitter element being noticeable only after the substance has passed over the posterior part of the tongue. Little or no sensation of taste can be aroused from the central portion of the tongue surface.




Fig 13 61 Pathway for the transmission of taste impulses CT chorda tympaninerve GN glossopharyngeal nerve IS superior salivatory nucleus LN lingual nerve SS superior salivatory nucleus efferent (secretory fibers) interrupted lines affectent (taste) fibers continuous lines

The sense of taste is much less sensitive than the sense of smell Sweetness for example is detected in a dilution of 1 part in 200 salti ness (common salt) 1 part in 400, sourness (hydrochloric acid) 1 part in 130 000 and bitterness (quinine) 1 part in 2,000 000

The chief nerves of taste are the chorda tympani branch of the facial nerve and the glossopharynged nerve. The chorda tympani supplies taste fibers to the anterior two thirds of the tongue, the glossopharyngeal to the posterior third (Fig. 13.61). The fibers of the chorda tympani nerve are conveyed to the tongue in the trunk of the lingual nerve (a branch of the mandibular division of the trigeminal nerve). The center for taste lies at the lower end of the somesthetic area of the cerebral cortex.

#### Smell

Smell is very closely allied to taste and has been aptly described as "taste from a distance" In many animals the sense of smell is almost

SMELL incredibly acute, a large proportion of the brain being concerned with this sense. In such species the sense of smell is of paramount importance, warning the animal of the approach of its enemies, and guiding it in the quest for food. Even in man, in whom the sense of smell is comparatively rudimentary, certain substances, such as mercaptan, can be detected in a dilution of 1 part in 30 billion or more parts of air. An odorous material continually emits particles of molecular size

which are carried in the air to the olfactory receptors. Substances which pass readily into the gaseous state such as turpentine, gasoline, and the




Fig. 13.62 Vertical transverse section through the nasal cavities to show position of the olfactory area.

essential oils have, in general, strong odors; whereas nonvolatile materials-e.g., the heavy metals-are relatively inodorous. A substance in order to be smelled must reach the nose in gaseous form,

The mucous membrane on each side of the nose is raised into three ridges by three spurs of bone (the superior, middle, and inferior turbinates or conchae) which spring from the outer nasal wall. The interior of the nose is thus divided incompletely on each side into four compartments or regions placed one above the other (see Fig. 13.62 and Fig. 6.3, p. 227). The lower three of these serve as air passages; they communicate with the outside through the nostrils and behind with the pharynx. The uppermost compartment is a narrow cleft lying immediately beneath the anterior part of the floor of the skull. The olfactory receptors are embedded in a small patch of mucous membrane situated on each wall of this narrow space, which is a blind pocket from which the main air currents are excluded. Air containing the odorous particles must therefore be carried to the olfactory mucous membrane 642

either by diffusion or by convection currents set up when the inspired air meets the warmer air within the nose When, for example, we wish to smell some particular scent, we make a quick short inspiration or to smell some particular seem, we make a quick short inspiration or "smill" This sharp indrawing of the cooler outside air creates ascend ing (convection) currents which convey the scent to the sensitive area. The material does not act directly upon the olfactory receptors but is first dissolved in the layer of fluid covering the mucous membrane—a fact which emphasizes the similarity between the senses of taste and smell

There are an almost infinite variety of odors, and it is very difficult to make a satisfactory classification, nevertheless, an attempt has been made to group them under the following eight headings


- 1 Ethereal odors-e g, of fruits
- 2 Aromatic or resinous odors-e g, of camphor, bitter almonds
- 3 Fragrant or balsamic odors-e g, of flowers, extracted or artificial perfumes
- 4 Ambrosial odors-e g, of musk
- 5 Garlie odors-e g, of garlie, onions, and of sulfur and selenium com pounds
- 6 Burning odors-e g, of burning feathers, tobacco, roasted coffee, and
- 7 Goat odors—e g, of caproic acid sweat and ripe cheese 8 Foul odors—e g of excrement decaying meat, and vegetable matter

It is not possible to correlate the chemical nature of substances with the odors which they emit, for materials quite different in their chemical constitution may have similar odors, while others closely similar chemically may have quite different smells. Certain sensations which we usually class as olfactory, such as those aroused by pungent and acrid substances are in fact due to the stimulation of the common nerves of the nasal mucous membrane

The olfactory epithelium is composed of spindle shaped nerve cells distributed evenly among other elongated cells which are purely sup porting in function Both types of cell he perpendicular to the epithelial surface (see Fig. 13 63) The nerve cells, or ollactory receptors, give rise to two types of process-an axon and a modified dendrite. The axons spring from the deep aspects of the cells and, combining in groups, form a number of slender bundles called the fila olfactoria The dendrites are thick cylindrical processes, they penetrate, individually, to the surface through small gaps between the free ends of the supporting cells, except for these gaps, the supporting cells are joined

together to form a continuous covering. Each dendrite, after emerging from between the supporting cells, divides into a tuft of some six or eight straight filaments which project a short distance beyond the epithelial surface. The fila olfactoria pass up through perforations in the floor of the skull and enter the olfactory bulb (Fig. 12.32), the primary olfactory center. Here, the axons of the olfactory receptors syn-

Fig. 13 63 Connections of the olfactory bibs. Left, 1, receptors in olfactory bulb. Left, 1, receptors in olfactory mucous membrane, 2, supporting cells Right, 3, olfactory receptors, 4, olfactory reves energing the cranial cavity through perforations in the bono, 5, connections with mitral cells, 6, mitral cells.



apse with other neurons (mitral cells) which convey the impulses to the cortical center for smell situated in the hippocampal gyrus.

The olfactory receptors adapt rapidly—that is, they soon cease to respond to some particular stimulus. It is common experience that an odor, though strong when first smelled, becomes imperceptible after a short time. This phenomenon of adaptation—a property which the olfactory sense organs exhibit in common with several other types of receptors—should not be confused with fatigue. That it is not simply a matter of fatigue of the olfactory mechanism is evident from the fact that, when some particular odor is no longer smelled, another odor is readily perceived. Some persons are unable to smell certain odors at all, though there is no general impairment of the olfactory sense. Hydrocyanic gas, for example, a powerful poison used in the extermination of vermin, has a strong odor of bitter almonds, but is quite inodorous to a few people.

### Cutaneous Sensations

The sensations which can be aroused by stimulating the skin are five in number—namely, touch, pressure, pain, heat, and cold Touch may be defined as the sensation elicited by lightly brushing the skin with a wisp of cotton wood, or by pressing a stiff hair vertically upon the skin until it bends. The latter method is used to test the degree of sensitivity to touch of a very small cutaneous area. The apparatus consists of a series of hairs graded in thickness and, therefore, in stiff ness (von Frey's hairs). The one that causes a sensation, when pressed upon the skin with just enough pressure to bend it, gives a measure of the tactile sensibility of the underlying skin (see Fig 13 64)

If a more rigid object such as a match stick is pressed against the skin, the sensation aroused is one of pressure Pressure, though usually classed as a cutaneous sensation and often confused with touch, is, in reality, due to the stimulation of receptors (Pacinian corpuscles) situ ated in the subcutaneous tissues. These sense organs are also found in other parts—e.g., in the surface membranes of bones (periosteum), beneath tendons, and in the mesentery If the rigid object is pressed still more firmly into the skin or if the skin is pricked with a sharp pointed instrument such as a pin, pain is experienced

The sensations of touch, pressure, heat, and cold are each dependent

upon a special type of sense organ in which the nerve fiber terminates after losing its neurilemma and myelin sheath. The sensation of pain, on the other hand is transmitted by fibers which terminate as bare axis

Fig 13 64 A von Frey hair (or esthesiometer) for measuring the sensitivity of the skin to touch

cylinders That is, the nerve fiber mediating pain loses its neurilemma and myelin sheath, but does not end in a structure of special design. The receptors of touch, pressure, and tem perature each respond to one type of stimulus, but the nerve endings giving rise to pain respond also to any other type of stimulation—mechanical, thermal, electrical, or chemical-pro-

vided it is intense enough. Thus the sensation of pain is protective in

function, serving to signal a threat of injury to the body. The several types of cutaneous receptor are shown in Figure 13 65.

The different types of cutaneous sense organ are separated from one another by measurable distances. By applying the appropriate stimulus to points upon the skin, the positions of the receptors can be determined. The small cutaneous areas mapped out in this way are referred.

to as "spots." Thus, when the sensitivity of the skin to touch is investigated with a von Frey hair, the sensation is elicited only from certain
points; these are called touch "spots," while those which respond to
heat or cold or to pain are called, respectively, hot, cold, and pain
"spots" (see Fig. 13.66). The touch spots are most numerous over the
tips of the fingers and in the mucous membrane of the tip of the tongue.




Fig. 13 65 Cutaneous receptors. A, touch (Messarer's corpuscle); B, cold (Krause's end bulb), C, warmth (Ruffini's end organ); D, deep pressure (Pacinian corpuscle), E, pain (bare nerve endings in cornea).



Fig. 13.66 Map of cold spots, I, and hot spots, 2, within an area on the palm The sensation in each case was most intense in the black areas, less so in the lined and mildest in the dotted areas. In the blank parts, no definite sensation was aroused (After Goldscheider.)

In cutaneous regions covered with hair—that is, almost the entire skin surface except the palms of the hands and the soles of the feet—the touch spots lie on the "windward" side of the hairs (the hairs slant in one or other direction like grass bent by a breeze). For this reason, light contact with the hair tips causes a sensation of touch The hair, when moved, acts as a tiny lever, transmitting the movement to the skin at its base; thus the touch end organ is stimulated. Pain fibers form a rich network in the skin; they also ramify within the hair sockets around the hair roots.

The localization of cutaneous sensations is effected with remarkable

accuracy. The localization of touch is tested by bringing a wisp of cotton wool or a von Frey hair in contact with the skin while the subject's eyes are closed, and then asking him to place a finger upon the spot touched. Pain and pressure are also very accurately localized, but the sensations aroused by heat and cold are more diffuse. Cutaneous localization is an acquired faculty, being developed through an association previously established in the brain, between the point upon the skin and the muscular movement required to touch it. In other words, it is based upon memories of muscular movements resulting from conditioned response to extraors the missing the state of the tioned responses to cutaneous stimuli

Aristotle's experiment illustrates our dependence upon experience for the interpretation of tactile stimuli. A small object, such as a pea, held between the adjacent sides of the first and second fingers is felt as one, because a touch stimulus when applied simultaneously to both of these surfaces has always in our experience been associated with but a single object. But when the small object is held between the fingers, a single object. But when the small object is held between the fingers, crossed as in Figure 13 67, the skin is stimulated at two points which experience tells us may be touched by two separate small objects but not by one alone. The single pea is, therefore, felt as two. Nerve cells in the sensory area of the cerebral cortex which receive impulses from the opposing sides of the fingers in their normal positions are probably very closely connected, whereas those receiving impulses from the other sides of the fingers are not in such intimate association. The

dual nature of pain is described on page 475

Spatial discrimination. This faculty is closely allied to the foregoing, it is the recognition of the separateness of two simultaneous stimuli. For example, if the points of a pair of compasses an inch apart are applied to the skin of the forearm, a sin gle sensation is felt. If the distance between the compass points is increased to one and a half Fig. 13 67 Arustotles experiment toolies experiment toolies experiment toolies and applied as before, two distinct toolies experiment toolies expe



shape of various objects, it is indispensable. This faculty varies considerably in different regions, being most highly developed in the coverings of the more mobile parts of the body—that is parts such as the fingers hips, and tip of the tongue, which have received the most practice in investigating the immediate environment In Table 13 1 several different regions are compared with regard to the minimal distance at which two simultaneous stimuli must be sepa rated in order to arouse a double sensation

Table 13-1

Different regions compared as to the minimal distance by which two

stimuli must be separated in order to arouse a double sensation	
Region	Minimal distance in millimeters
Tip of tongue	11
Palm side of finger tip	2 3
Red part of lips	4 4
Tip of nose	66
Palm of hand	113
Heel	22 0
Back of hand	31 6
Forearm	39 6
Middle of back, upper arm and thigh	67 0

# The Physiology of Reproduction

The cells that have been set apart from the general body mass for the perpetuation of the race are called the sex cells, germ cells, or gametes (Gk. gametēs, a husband, gametē, a wife). The organs that contain them, and in which they are fully or incompletely developed, are known as the gonads.

# The Female Reproductive Organs

The gonads of the female are called ovaries, and the cells that they produce are known as ova or egg cells; the latter receive different and more specific names at progressive stages of their development (see p. 661). The human ovaries are two bodies about the size and shape of olives, though somewhat flattened from front to back (3 cm long, 1.5 cm broad, and 10 mm thick). They lie one on each side of the pelvis. The ova are developed from columnar epithelial cells-the germinal epithelium-covering the surface of the ovary. Columns of these cells penetrate deeply into the connective tissue or stroma of the ovary. Small groups of cells separate off from the columns and become arranged with a large one in the center and others in a single layer around it. These structures are found in great numbers in the fetal ovary and in the ovaries of children. They are called primary, primitive, or primordial Graafian follicles (Fig. 14.1). The central somewhat larger cell is a primordial ovum or oocyte. In the sexually mature female—that is, from the time of puberty to the menopause—

1 The sex hormones are dealt with in Chapter 10

some of the primitive follicles, under the influence of the folliclestimulating hormone (FSH) of the hypophysis, become, from time to time, mature. Two more layers of cells derived from the ovarian stroma develop around the original one encircling the ovum. The outer of these is fibrous and known as the theca externa; the inner layer, called the theca interna is more vascular and cellular in character. The cells of the original layer surrounding the primitive ovum multiply to form a mass several strata deep. This cell mass is soon separated into two parts by the collection of fluid—the liquor folliculi—near its center.




Fig. 14.1 A section of the ovary.

A heaped-up hillock of cells remains at one side of the follicle in which the ovum lies embedded, though separated from the surrounding cells by a clear, rather thick membrane called the zona pellucida. The heaped-up mass of cells just mentioned is known as the discus proligerus (L. proles, offspring + gero, I bear) or the cumulus oophorus (L. cumulus, a heap + oophorus, egg-bearing). A single row of regularly arranged cells immediately encircling the zona pellucida is named the corona radiata. The other mass of cells, separated off by the liquor folliculi, is pressed against the wall of follicle which they line. They constitute the membrana or zona granulosa. As the follicle matures or ripens, it becomes distended by the accumulation of liquor folliculi and moves outward again to the surface of the ovary. It projects from the ovarian surface as a small cystlike oval swelling which eventually bursts and discharges the ovum. In women the discharge of an ovum, or ovulation, as it is termed, occurs at regular intervals of about 28 days (see p. 447). The cavity of the ruptured follicle becomes filled with a clot of blood which is soon replaced by a mass of cells filled with a vellow fatlike material called lutein. The cells are derived from the membrana granulosa and theca interna of the follicle. The yellow mass filling the follicle is called the corpus luteum (yellow body), it elaborates a hormone named progestin when referring to it in pure crystalline form, it is called progesterone (Chap 10).

The own itself, when discharged from the ovary, is immature that the transfer of the transfer




Fig 14.2 Drawing of a Graafian follicle approaching maturity

along the duct by the movement of the cilia in its mucosa, as well as by peristaltic contractions of its wall. The uterine tubes open below into the upper part of the uteris (womb). Conjugation of the ovum with the male germ cell (spermatazoon), or jertilization of the ovum, as this event is usually termed, is thought to take place in the uterine tube. In birds, the oviduets transmit the fertilized ovum to the exterior, but in the human body, and in the bodies of most other mammals, it is delivered into the uterus.

The uterus is a hollow pear shaped organ though somewhat flattened from front to back. Its walls are formed of smooth muscle and lined by

mucous membrane; the latter is called the *endometrium*. The small narrower end, the neck or *cervix* of the uterus, is directed downward, opening by its mouth or *os* into the upper end of the *vagina*. The latter is a narrow sheathlike canal of muscle and mucous membrane which




Fig 14.3 Above, diagram of the human female organs of reproduction Below, a section through the female pelvis to show the relations of the reproductive organs to neighboring parts. The bladder is distended.

receives the male copulatory organ or penis. The fertilized ovum, upon reaching the cavity of the uterus, establishes connections with the uterine mucosa and develops into the fetus (p. 669). The reactions of the endometrium will be described in greater detail later.

The growth of the uterus in the virgin state and during pregnancy. The virgin uterus reaches its full development at puberty as a result of the maturation of the Graafian follucles and the liberation of the follicular hormone, estradiol. At this time it weighs only 50 grams or less, but




Fig. 14.4 D agram of a segment of a mature mammary gland A section of skin and subcutaneous tissue has been removed *Lower* drawing shows the microscopical appearance of the glandular tissue during secretion note that disintegration of the follicular cells occurs as part of the secretory process.

during pregnancy it shows phenomenal growth reaching a weight exclusive of its contents of some 1000 grams—a twenty fold increase Almost immediately after fertilization of the ovum, before it has be come attached to the endometrium, and even while it is still in the uterine tube the uterine muscle shows very active hyperplasia, a great increase in the number of fibers occurring through mitosis Under microscopic examination thousands of mitotic figures are to be seen in a small sample of the uterine muscle But this hyperplastic phase soon comes to an end. The growth of the uterus from this time on is confined entirely to thickening and lengthening of the muscle fibers. The in

crease in the number of fibers (mitosis) is due mainly to the action of progestin. The enlargement of the individual fibers, and the consequent great increase in the size of the uterus, is the result of the distension caused by the progressive increase in bulk of the fetus, a stretch stimulus being thus applied to the uterine muscle. The effect of this stretch stimulus upon uterine growth is modified after a somewhat complicated fashion by both estradiol and progestin. The growth of the uterine muscle ceases toward the end of the suxth month of pregnancy. But the fetus shows rapid growth from this time on, accompanied by a commensurate distension of the uterus. The stretch stimulus applied to the muscle fibers, though it elicits no growth response, induces other effects which lead eventually to the termination of pregnancy (p. 674).

The structure and development of the mammary glands. The composition of milk. The mammary glands of women, commonly known as the breasts, are composed of a mass of cells arranged to form follicles or alveoli which are drained by a branching system of ducts. The mammary tissue, capable of secreting milk under appropriate stimulation, is separated by means of partitions or septa of connective tissue into some twenty lobes in each breast (Fig. 14.4). Each lobe is divided again into a number of lobules. The breast is surmounted at its center by a small conical eminence composed of erectile tissue and covered by darkly pigmented skin; it is known as the nipple, and is surrounded by a circular area of similarly pigmented skin called the areola. The whole mass of glandular tissue is embedded and covered by a variable, but often a very abundant, amount of fatty (adipose) tissue. In the active gland, the milk is secreted into the cavities of the follicles and drained away by the duct system. The milk is carried from each lobe by a single duct (lactiferous duct) to the nipple, from the apex of which it reaches the exterior through a tiny pore.

The development of the mammary glands to the adult size and form commences at puberty. Their enlargement at this time is one of the chief secondary sex characteristics of the mammalian female. Before the age of puberty, the gland of the female shows little difference from that of the male either in the amount or in the histological appearance of the mammary tissue. In both sexes at this time, as well as in the adult male, the glands consist mainly of conective tissue with a limited number of rudimentary collapsed follicles and a few scattered sparsely branched ducts. With the onset of puberty, signs of cellular activity (hyperplasia) appear in the female gland. The duct system becomes more extensively branched, though the follicles remain rudimentary.

Not unless pregnancy supervenes do the latter become fully developed and capable, under ordinary circumstances, of secreting milk. There are, therefore, two periods in the growth of the mammary glands, the first at puberty and confined to the duct system. This is brought about by the action of the follicular hormone (estradiol, p. 450). The very active growth of the follicles which is characteristic of the pregnant state, is caused by progestin Due to the high concentration of estradiol in the blood during pregnancy, the duct system also receives further stimulation at this time. Though milk is formed within the gland toward the end of pregnancy, none appears externally, as a rule, until the child or the young animal has been born. The mechanism underlying the secretion of milk (lactation) has been described on page 454.

Milk provides all the materials necessary for the growth of the infant up to the age of about 6 months. Human milk contains about 6 7 percent of sugar (lactose or milk sugar), 1.5 percent protein (mainly lactalbumin and cassein), and about 4 percent of fat. Its main mineral is calcium (approximately 1 gram per quart of cow's milk). But it also contains adequate amounts of phosphorus, sodium, potassium, and magnesium. The chief difference between human and cow's milk is the higher concentration of protein (3.5 percent) in the latter. Both kinds of milk are poor in iron; it is for this reason that the infant after the age of about 6 months is likely to become anemic if fed upon milk alone. The baby comes into the world with a good supply of iron, but the stores of this mineral (derived from the hemoglobin freed by the disintegration of the excess of erythrocytes during the few days after birth) become exhausted after the first half year or so. The milk of healthy mothers usually contains the required amounts of vitamins A, B, and C, but tends to be low in vitamin D This is also often true of cow's milk, and for this reason the diet of a child after 6 months of age-a time when rickets may develop-should be supplemented with some

preparation of vitamin D, especially in the winter months.

The various constituents of the milk are derived from the blood flowing through the gland. In a secreting gland the fat can be seen as microscopic globules within the cells of the alveoli As the globules collect and coalesce, the cells swell, and finally bursting, they discharge their load of fat into the center of the alveolus (Fig 14.4). This type of secretion, known as apocrune, is less common than that in which the cells extrude their product-saliva, for example-and remain intact.

The menstrual cycle is described on page 447, the actions of the ovarian hormones on page 449, and of the gonadotrophic hormones of the anterior lobe of the pituitary on page 452.

## The Male Reproductive Organs

The testes. These are the gonads or sex organs of the male and, herefore, correspond to the ovaries of the female. They are two ovoid bodies which, in such animals as fish, frogs, repules, and birds, lie within the abdominal cavity. Even in the mammalian fetus they occupy the position. Shortly before, or soon after birth, the testes of mammalis descend and, leaving the abdomen, become enclosed in a small cutaneous pouch, suspended from the pubic and perineal regions, and called the scrotum. In the human fetus the testes enter the scrotum two months before birth.

The interior of the testis is a compact mass of narrow and very tortious tubules—the convoluted seminiferous tubules. Fibrous partitions divide the testicular substance into a great number of wedgeshaped lobes, each of which consists of from one to three convoluted tubules. The tubules of neighboring lobes unite to form a series of

larger straight ducts which, after a short course, unite in a plexiform manner. The plexus-known as the rete testis-leads again into a number of ducts, small and straight at first, but which, after a short course, become enlarged and tortuous (Fig. 14.5), and ultimately unite into a single large convoluted duct. This is called the epididymis; it is applied to the posterior aspect of the testis, its upper part or head being considerably larger




Fig 14.5 Diagram showing the structure of the interior of the testis.

than the lower part or tail. From the tail, a straight tube—the vas deferens—ascends along the posterior border of the testis to enter the abdomen, wherein it joins the duct of the seminal vesicle of the corresponding side (Fig. 14.6).

The seminal vesicles are two coiled tubes with sacculated walls, situated between the lower part of the bladder and the rectum. The ejaculatory ducis are short tubes formed one on each side by the union of the duct of the seminal vesicle with the vas deferens; they open into the urethra (the canal of the penis) near the outlet from the bladder.

The seminiferous tubules are lined by several layers of cells. Those of the outermost layer—that is, the layer lying upon the basement membrane—are of two basic types, (a) spermatogenic cells and (b) columnar cells. The former give rise through a series of divisions (p. 660) to the mature male sex cells (spermatozoa). The columnar cells are supporting cells and are named the cells of Sertoli: they extend




Fig 14.6 Showing the relation of the vas deferens, V, seminal vesicles, S, and prostate, P, to the bladder, B R, urethra, U, ureter. (Redrawn from Cunningham.)

inward (i.e., toward the center of the tubule) through all the other layers. The inner cell layers forming the wall of the tubule consist of spermatogenic cells showing various stages in the maturation process, those most advanced in development lying nearer the lumen of the tubule. The cells, as they mature into free-swimming spermatozoa, become detached from the tubule wall (Fig. 14.7). The spermatozoon is about 0.1 mm long; it has an oval flattened head and a long tail-like process (Fig. 148) by

which it propels itself. The head is the essential part of the cell, consisting of a large nucleus surrounded by a narrow rim of protoplasm.

The connective tissue lying between the convoluted tubules contains scattered cells with yellow granules in their cytoplasm. These are called interstitual cells or the cells of Leydig. They are believed to furnish the male hormone, testosterone (p. 452).

The spermatozoa are conveyed from the convoluted seminaterous tubules along the complex system of canals, just described, to the epididymis. The spermatozoa show no spontaneous movements in the convoluted tubules, but become actively motile in the epididymis. As coitus nears completion and the climax or orgasm of the act occurs, contractions of the epididymus and vasa deferentia propel the spermatozoa through the epaculatory duets into the urethra. At the same time the seminal vesicles contract and expell a viscous secretion. The semen, which this fluid with its suspension of spermatozoa is now called, is ejected from the urethra with considerable force by the contractions of the urethral muscle and of striated muscle in the perineum. The ejection of the semen and the movements which bring it about constitute the act of ejeculation. It is a reflex act of which sensory nerves in the

penis are its afferent limb, and sympathetic nerves its efferent limb. A thin secretion from the prostate gland and secretions of small urethral glands are added to the semen in the urethra. The secretion of the seminal vesicles appears to be essential for maintaining the life and motility of the spermatozoa, but the prostatic secretion and the secretions of the urethral glands are not of importance in this respect; they probably serve mainly as a lubricant.

The spermatozoa, deposited in the upper part of the vagina during coitus, propel themselves upward by lashing movements of their tails




Fig. 14.7 Cross section of seminiferous tubule showing spermatogenesis (semidiagrammatic).



Fig. 148 A human spermatozoon. A, front view, B, profile view.

at the rate of about 6 inches per hour and, passing through the uterus, enter the uterine tube where fertilization normally occurs. It is probable that contractions of the uterus during coitus may draw the semen into the uterine cavity.

The head of the spermatozoon penetrates the ovum; in a short time its tail disappears and the changes described on page 663 follow. The semen contains an enzyme (hyaluronidase, p. 51) which effects the removal of the mass of cells (discus proligerus, p. 649) in which the ovum is embedded, thus facilitating its penetration.

The life of the spermatozoon after it has been deposited in the vagina is from four to five days. The ovum, on the other hand, if unfertilized, survives for only about seven hours; after this, it starts to degenerate.

Nature, in her provisions for the conveyance of the ova to their destina tion, has taken a rather hazardous chance. Whereas the spermatozoa follow a completely closed course from the testes to their exit from the penis the trumpet shaped end of the uterine tube and the surface of the ovary are not structurally continuous Sometimes the ovum fails to enter the tube but escapes into the pelvic or abdominal cavity where it may meet a sper




Fig 14.9 Upper drawing a section through the male pelvis to show the rela tionship of the reproductive organs Below a cross section of the penis The bladder has been artificially distended

matozoon Or after the ovum has entered the tube and been penetrated normally by a spermatozoon it may be prevented from continuing its journey to the uterine cavity Or again, the ovum and the spermatozoon may meet upon the surface of the ovary. In the event of one of these sen ous mishaps occurring pregnancy takes place in the general abdominal cavity, on or within the ovary, or in one or other uterine tube. This is called extrauterine pregnancy or ectopic gestation, or, more specifically, abdominal tubal or ovarian pregnancy according to the site. The embryo may develop to maturity in the abnormal situation, but more usually severe hemorrhage within the abdominal cavity occurs with death of the embryo and grave danger to the mother

Pregnancy occurring primarily in the abdomen is rare Abdominal pregnancies are usually secondary to rupture of a tube in which the pregnancy originally occurred Ovarian pregnancy is also very rare.

The pens. The copulatory organ of the male, the pens, is com posed of erectile tissue arranged as three longitudinal columns bound together by fibrous tissue, and covered with skin Two of the columnar masses, called the corpora cavernosa pens, lie side by side on the up per or anterior aspect of the organ. The third called the corpus caver nosum urethrae or corpus spongiosum penis, lies beneath the other two and is canalized by the urethra. The urinary bladder is evacuated through the urethra which also transmits the semen during coitus. The extremity of the corpus cavernosum urethrae is expanded into a pyram dal structure called the glans penis, which is molded over the ends of the corpora cavernosa penis (Fig. 14.9)

Erectile tissue possesses a spongelike structure, showing a mesh work of wide blood spaces (cavernous spaces) These are fed by capillaries and arterioles and drained by small veins. Smooth muscle fibers run in the walls of the blood spaces and surround their venous outlets. To this construction of erectile tissue is due its peculiar property, namely, the ability to alter in volume and consistency. Erection of the penis is brought about in the following way. The arterioles feeding the blood spaces dilate, and the muscle fibers in the walls of the latter relax. The muscle guarding the venous outlets contracts thus tending to impede the outflow of blood. The spaces of the erectile tissue are expanded as the blood under high pressure is driven through them. The organ, thus becoming turgid with blood, is rendered tense, hard, and erect.

The nerve fibers governing this mechanism are derived from the pel vic (parasympathetic) nerves. The sympathetic sends fibers which exert the reverse effect (contraction of the smooth muscle of the arteriol of that surrounding the venous outlets) with consequent relaxation of the penis.

The erectile organ of the female corresponding to the penis is called the clitoris. It is situated above and just outside the entrance to the vagina. Erectile tissue is also present beneath the mucosa of the vagina just within the vaginal orifice.

The prostate. This is a body about the size of a chestnut and some what conical in shape. Its base is directed upward and lies in contact with the lowest part of the bladder. It embraces the first 1½ inches of the urethra. The ejaculatory ducts pierce its upper and posterior part (see Fig. 14.9). The prostate is composed of muscular and glandular tissues. It secretes a thin fluid into the urethra.

Prostatic enlargement, with consequent interference with the pas

sage of urine, not uncommonly occurs in men past middle age. There has been much speculation as to the cause of enlargement of the prostate, but so far the problem has not been solved. Some hormonal imbalance is thought to be the fundamental cause.

# The Processes of Reproduction

Maturation of the sex cells. The sex cells in the course of their development to their mature form—spermatozoa or ova—pass through a series of divisions which will now be described.

The maturation of the male gametes—spermatogenesis. The youngest male gametes produced by the multiplication of the primary sex




Fig. 14.10 Spermatogenesis The black bodies represent the chromosomes, X and Y are the sex chromosomes (see p. 665). Only one autosome is shown in the spermatozoa, in man the total number of chromosomes is 46.

cells (primordial germ cells) are called spermatogonia. They are large plump cells lying adjacent to the basement membrane of the seminiferous tubules

The spermatogonia multiply by mitosis (p. 25) through several generations. The cells produced in the final mitotic division pass into a growth period. The large cells that result are called primary spermatocytes (Fig. 14.10). At the end of the growth period, the latter again divide, but the division with respect to the arrangement of the chromo-

somes is fundamentally different from that occurring in the mitotic process. The chromosomes come into apposition in pairs but do not, as in mitosis, split lengthwise. As a result of the mere contact, or synapsis, of the chromosomes and their failure to divide, the diploid number is maintained and not temporarily doubled, as in mitosis. When, therefore, the cell divides and two daughter cells, now called secondary spermatocytes, are formed, each new cell receives only half the number (haploid number) of chromosomes characteristic of the body cells of the species. This type of cell multiplication is called re




Fig 14 11 Oogenesis Oval bodies represent the chromosomes

duction division or meiosis and is peculiar to sex cells, occurring in no other type of cell. The secondary spermatocytes, having received their haploid quota of chromosomes, now divide by mitosis. The small cells so produced are called spermatids which, owing to the splitting of the chromosomes, also receive the reduced number of chromosomes. The spermatids, without further division are gradually transformed into mature spermatozoa.

Maturation of the female gametes—oogenesis The maturation of the primitive ovum or oogonium follows essentially the same course as that described for the spermatogonia, but there are certain differences. The earliest stage in the maturation process is, as in spermatogenesis, the production by mitosis of daughter cells with the diploid number of chromosomes

The cells resulting from the final mitotic divisions enter a period of growth and are transformed into primary oocytes (Fig. 14 11) Production of the latter is the last stage of maturation which occurs in the

<sup>.</sup> See glossary for definitions of diploid and haploid.

ovary ' (Graahan follicle) After rupture of the follicle and the escape from the ovary of the primary occyte the latter enters the uterine tube (or oviduct) If there met and penetrated by a spermatozoon it under goes further ripening otherwise it degenerates. The resumed matura tion consists of the meiotic division of the primary occyte and the production of cells containing the haploid number of chromosomes. But unlike the corresponding stage of spermatogenesis the division of the




Fig 14 12 Fert I zation of the ovum

cytoplasm is very unequal one of the cells—secondary oocyte—is large and functional the other is diminutive and functionless. It is named the first polar body and soon degenerates. The secondary oocyte divides by mitosis but again unequally to produce a mature or urn and a second polar body which like the first is a miniature and functionless cell which degenerates. The first polar body also as a rule divides mitotically at this time so that from each primary oocyte one mature egg and three small functionless cells are produced. Each of the four cells contruss the haplerd number of chromosomes. The ovum is now ready for penetration by the spermatozoon.

According to some a horites the own of most vertebrates progress in the own to a size further than that of the prin ary occyte and in the human subject. If fully matured in the own;

663

Conjugation of the egg and sperm cells. The unfertilized ovum appears, in some unknown way, to exert an attractive force upon the spermatozoon which, by means of the lashing movements of its tail, reaches the female gamete. The head of the sperm penetrates the ovum. A change in the boundary wall of the ovum then occurs, which, as a rule, serves to prevent the entrance of other spermatozoa. The tail of the spermatozoon soon disappears, leaving the head containing the nucleus within the cytoplasm of the egg (Fig. 14.12). The result is the formation of a single cell, which has a power for growth not possessed by either ovum or spermatozoon alone. The single cell almost at once commences to divide and redivide. In this way are formed great numbers of cells (see segmentation, p. 666), which do not separate from one another but are held together. Gradually, as they multiply and develop in various directions, tissues and organs of different structure and functions appear, until finally a new and complete individual is created almost identical with others of its species. Whether plant or worm, fish, bird, or man, nearly all forms of life other than the very primitive have developed from a single cell resulting from the union of sperm and egg (see parthogenesis, p 664).

The conjugation of the male and female gametes is termed the fertilization or impregnation of the ovum. The cell resulting from the union is then spoken of as the fertilized ovum, oosperm, or zygote.

Changes in the nuclei of the conjoined cells. Almost immediately after the penetration of the ovum by the spermatozoon, the two nuclei move toward each other, come together, and finally fuse near the center of the ovum. Fusion is followed by the complicated changes already described as characteristic of mutosis and which leads to masses of cells being produced. In the process of mutosis, the chromatin of each nucleus is not broken up into sections at random or in any hap-tazard fashion, but is always divided into precisely the same number of chromosomes characteristic of the body cells or primitive germ cells of the species, but double the number in a mature germ cell. In other words, the original (diploid) number which was reduced during meiosis is restored by the fusion of the nuclei. The fused nucleus must therefore contain an equal number of maternal and paternal chromosomes. These are important and fundamental facts in the mechanism of reproduction.

<sup>&</sup>lt;sup>8</sup> In some insects, such as the fruit fly, the number of chromosomes in the body cells is 8, which are reduced to 4 in the sex cells. In certain species of worms, there are 12 in the body cells and 6 in the ova and sperms The mouse, the trout, and the lily have 24 in their body cells and 12 in their sex cells. Other forms of animals and plants have each a characteristic number, which may be as low as 4 or as high as 168. There are 23 chromosomes in the gametes of the human race and 46 in the body cells

Chromatin, of which the chromosomes are composed, is believed to be the essential procreative substance. Through it hereditary characters are transmitted, and upon its existence the perpetuation of the species depends. Yet it has been stated that the total quantity of chromatin contained in all the ova and sperms from which have been created the two-billion-odd persons inhabiting the globe could have been no greater in size than a match head.

Parthenogenesis—virgin birth. In some animal forms, especially certain insects, the females, though themselves produced from a fertilized egg, may for some generations lay eggs which are not fertilized. These spermless eggs, nevertheless, develop into the young of the species. In other words, the young of these generations have a mother but no father 'To this mode of reproduction parthenogenesis, a term derived from the Greek word parthenos, a virgin, is applied. The drones (males) of the honeybee, for instance, are developed from untertilized eggs, which have been laid about the same time as the unfertilized. A frog's egg, though under ordinary circumstances it must receive a spermatozoon before segmentation will ensue, may be induced to develop into a tadpole by artificial means. Pricking the wall of the own with a needle will start segmentation, which ultimately results in the production of a fatherless frog.'

Twinning. There are two ways in which twins may be conceived:

(1) by the fertilization of two separate ova, or (2) by the fertilization of a single ovum. When two ova are fertilized—and this is the commoner way—the offspring, known as fraternal twins, may be of the same or of different sexes and usually do not resemble each other more than might any two members of a family. When two individuals are formed from a single zygote, the twins are always of the same sex and are remarkably alke. They are called identical twins and are formed by the division of the cell mass within the zygote into two distinct parts, each of which develops into a separate child. Identical twins are contained within the same amniotic cavity and are on very rare occasions joined together, as in the case of the Siamese twins. Most animals produce their litters by the fertilization of several ova, but in some, such as the armadillo—a little mammal possessing a hard covering resembling a coat of mail—only one ovum is fertilized, and

The unfertulized egg gives off only one polar body (see p. 661).
 As a very rare event parthenogenesis may occur is mammals, and the possibility of its occurrence in the human species has been seriously discussed.

the mass of cells within becomes divided into several smaller groups, each of which develops into a fetus.

The determination of sex. It would be an inestimable boon to animal breeders if the sex of the offspring could be determined by artificial means, or even if it could be predicted. Man's ingenuity in this direction seems, however, to be forever checked. Had human interference been possible in the past, the history of nations, no doubt, would have run a different course. The sex of the newly created organism is fixed unalterably at the moment that the spermatozoon fertilizes the ovum. A pair of chromosomes (sex chromosomes) in the nuclei of the immature sex cells determines whether the zygote shall develop into a male or a female. None of the other chromosomes (autosomes) exercises this function. In the male, the sex chromosomes are dissimilar, one, called the Y chromosome (Fig. 14 10), being usually smaller than the other. The large one is called the X chromosome. In the immature female sex cells, the sex chromosomes are identical and like the male X chromosomes. Now when the male sex cells develop into mature spermatozoa and reduction division occurs-i.e., when the pairs separate—half of the sperm cells receive Y chromosomes and half receive X chromosomes. If a spermatozoon possessing a Y chromosome conjugates with the egg, a male (XY) will be produced, if the sperm cell contains only X chromosomes, a female (XX) results Obviously, it is all a matter of chance what the sex of the offspring will be. When the gonads of the embryo develop, the influence of the testicular or the ovarian hormone encourages development along male and female lines, respectively.

Hermaphroditism. The hormonal influence upon sexual development is sometimes seen in a very striking way in cattle when, as a result of a developmental anomaly in the circulations of twin embryos of opposite sexes, the blood of the two fetuses is permitted to mix. The female embryo then comes under the influence of the male hormone and abnormalities in its generative organs are induced. The uterus remains undeveloped and masculinization of other reproductive structures occurs. The masculinized female is called a freemartin; it is sterile. Feminization of the male fetus does not occur.

Some lower animal forms—certain worms, mollusks, crustacea, etc.—like most plants, contain both male and female reproductive tissue in the same body. Sometimes both testes and ovaries are functional, producing spermatozoa and ova; sometimes only the testes or ovaries produce sex cells. This combination of the sex organs (gonads) is called hermaphroditism, and the individual an hermaphrodute (Gk.

god and goddess, Hermes and Aphrodite). An hermaphrodite may fertilize (p. 663) its own eggs or those of another.

In man, hermaphroditism occurs as a rather rare developmental ahnormality. The two types of tissue, testicular or ovarian, may be present as separate organs (testes and ovaries) or fused to form ovatestes. The external sex organs—penis, clitoris, etc.—are indeterminate between male and female as a result of the double hormonal influence. Such a condition is called true hermaphroditism. It is of genetic origin. On the other hand, only one type of sex gland, testes or ovaries, may exist, though the external genital organs show both male and female features. This is called pseudohermaphroditism. It most fre-




Fig 14.13 First stages of segmentation of a mammalian ovum and the formation of the morula (semidiagrammatic). (Redrawn and modified from Allen Thomson)

quently has an endocrine basis Occasionally certain regions of the body, even an entire half, has male characteristics while other regions or the opposite half are female. Such persons are called gnandro-morphs (Gk. gynē, a woman + anēr (and-), a man + morphē, form). It is often difficult to decide in which sex a pseudohermaphrodite should be placed and, if a child, whether dress, education, training, etc., should be along male or female lines. Recently a means has been discovered whereby the dominant sex can be determined. There is present in the body cells of females, but not in those of males, a dot of chromatin within the nucleus in contact with, or close to, the nuclear membrane. It is called the sex chromatin. By excising a small piece of skin and examining it microscopically for the presence or absence of this body, the real sex of the individual can be ascertained.

Segmentation of the fertilized ovum and the development of the embryo. The zygote immediately divides into two Each daughter-cell again divides, and through a succession of divisions groups of 8, 16, 32, 64, 128 cells, and so on appear. So, by a process of division and

redivision, large masses of cells are formed, which ultimately produce a new individual. This process whereby the fertilized ovum undergoes repeated divisions is called segmentation or cleavage. With each cell division, the chromatin breaks up into chromosomes; and, as described above for mitosis, each chromosome splits into half, and an equal number of halves go to each new nucleus. All the cells resulting from the long series of divisions which occur in the development of the offspring must therefore contain the same number of chromosomes as

Fig 14.14 The blastocyst. On left, diagram showing the segregation of cells of the morula into an outer and an inner group. (After Simon.) On right, formation of amnion, A, and yolk sac, Y. (After Bryce)





did the original fused nucleus of the fertilized ovum, and each receives chromatin material from both parents.

In the earlier stages of embryonic development of the various mammalian species, cell multiplication follows a common pattern. At first a rounded, mulberrylike mass of cells is formed, called the morula (L. diminutive of morus, a mulberry) The morula soon differentiates into an outer and an inner group of cells (Figs. 14.13, 14.14, and 14.15); fluid collects within it. The fertilized ovum at this stage is called the blastocyst. The outer group of cells now pressed excentracally to form an annular wall is called the trophoblast (Gk. trophe, nourishment + blastos, germ or primitive form) because through its agency the blastocyst receives nutriment from the mother. The trophoblast later develops numerous fringelike processes which gives the ovum a shaggy appearance. These processes grow larger and branched and are now known as chorionic villi; the outer covering of the ovum from which they arise is called the chorion. The more centrally situated mass of cells becomes so disposed as to form the walls of two sacsthe amnion and the yolk sac (see Plates 15C and 16A, and Fig. 14.14). The cells of the area over which the amnion and the yolk sac come into contact multiply to form a plaquelike elevation called the embryonic shield or disc, from which the body of the embryo is developed. The

cavity of the amnion enlarges and comes to enclose the embryo, the sac expanding until it comes into contact with the inner surface of the chorion. The yolk sac shrinks and almost disappears.

When the embryonic shield is no more than 2 mm or so in diameter, its cells become arranged in two layers, separated by a narrow space. The outer of these layers is called the *ectoderm*, and the inner one the



Fig. 14.15 Photograph of an 8-day-old blastocyst of a monkey attached to the mucous membrane of the uterus. The lighter area within is the embryo itself. (After Heuser and Streeter, Contributions to Embryology, 479, Carnegie Institute of Washington, 1937.)

entoderm. A little later, a third layer appears in the space between the other two; it is known as the mesoderm. From these three germ layers, to give them their collective name, all the structures of the body are ultimately formed (Fig. 14.16). The nervous system, the greater part of the eye, the salivary glands, the skin, the epithelial lining of the nost and part of the mouth, and the dental enamel originate from the ectoderm. The mesoderm gives rise to the skeleton and the muscles (both striated and nonstriated), the heart, blood vessels, and blood, the lymphatic system, the kidneys, the urinary bladder (with the exception of its epithelial lining), and the connective tissues. From the entoderm are developed the epithelial linings of the alimentary and respiratory tracts exclusive of those parts derived from the ectoderm, the epithe-

lium lining the urinary bladder, and the secreting (epithelial) cells of the liver and the pancreas, and of the parathyroid and thyroid glands.

The earlier of those changes just described-up to the stage of the blastocyst and the development of the trophoblast-take place in the Fallopian tube while the blastocyst is being propelled toward the uterus by ciliary action and peristaltic movements of the tubal walls. Between the third and the fifth days after fertilization of the ovum the blastocyst enters the cavity of the uterus. Through the destructive effect that the trophoblast exerts upon the uterine mucosa and the reaction set up in the latter, the ovum is enabled to embed itself and, like a parasite, obtain nourishment from the maternal tissue. The activity shown by the uterine mucosa at this time consists of multiplication of the surface epithelium and elongation and hyperplasia of its glands, together with dilatation and twisting of its vessels. A special type of uterine tissue is thus produced called the decidua because it is shed after the birth of the offspring (Pl 16A) These changes in the endometrium commence before the blastocyst reaches the uterus, or after ovulation, even though fertilization has not occurred (see p. 450) They are dependent upon the action of progestin (progesterone)



Fig. 14.16 The germ layers.

This hormone also diminishes the irritability of the muscle of the uterus; any contractions of the muscle which might occur to disturb the secure implantation of the blastocyst are suppressed. Hyperplasia of the uterine muscle, as mentioned on page 450, also occurs at this time.

In Figure 14.17 a fetus at the age of 8 weeks is shown completely enclosed within the amniotic cavity. Most of the chorion has been cut away. The chorionic villi have disappeared except where the chorion is attached to the uterine wall, in which situation they have increased greatly in size and complexity. They are surrounded by spongy masses of decidual tissue consisting largely of blood spaces; thus the embryonic and the maternal tissues are intimately interlocked. The chorionic villi, which are supplied through a rich vascular system with fetal blood, are bathed by the mother's blood. The structure formed by the

union of the maternal (decidua) and embryonic tissues is called the placenta. When fully grown, the placenta is a disc shaped mass which occupies a third or so of the uterine wall (Fig. 14.18). Though the fetal and maternal circulations come into the most intimate association in the placenta, the blood in the two sets of vessels does not mix, the delicate capillary walls of the chorionic villi are interposed. Oxygen,



Fig. 14.17 Showing the fetus at age of about 8 weeks enclosed in the amnion magnified a little over 2 diam eters (After Thomson from Gray's Anatomy)

food materials etc, are transferred across the walls of the chorionic vessels from the mother to the fetus and carbon dioxide and other waste materials from the fetal to the material circulation. Thus the placenta serves the respiratory, nutritional and excretory functions of the fetus.

The law of recapitulation or biogeness: The vertebrate embryo from the earliest stages of its development tends to recall or recapitulate in a more or less vague way the forms of its ancestors from a remote period in its evolutionary history. This is known as the law of recapitulation or biogenesis. For example, the embryo of the human species or of any vertebrate commences its existence as two cells (ovum and

sperm) fused into a single one. Some unicellular forms reproduce by a similar preliminary fusion of their bodies. The segmentation of the fertilized ovum with the formation of the morula is identical with that seen in the development of many of the most primitive forms of life.




Fig 14 18 Drawing of a human embryo A, at the age of 5‡ weeks (about 8 mm long), and of a monkey embryo B, of about the same age C, enlarged drawing of the five branchial arches and clefts The inner aspect of the throat region is exposed to show the pharyngeal pouches

Again, in the early vertebrate embryo, including the human, a series of five arched ridges appear in the entoderm of the primitive pharyux, corresponding elevations are to be seen on the surface (ectoderm) of the neck. The arched ridges have a core of mesoderm, and are known as the branchial or visceral arches, the depressions or furrows between them on the outer aspect of the neck are called the branchial clefts, the furrows (entodermal) in the pharyux between the arches are re-

ferred to as the pharyngeal pouches (Fig. 1418). The branchial structures are found in all vertebrate embryos. From them the respira tory apparatus of fish is developed, the tissue between the arches (bran chial clefts and pharyngeal pouches) disappearing to leave slits which establish communications between the pharynx and the exterior In the fish the arches (with the exception of the most headward one which goes to form the jaw) develop rich capillary plexuses which, with the supporting tissue, constitute the gills Water taken into the mouth and pharynx is ejected through the slits and passing swiftly over the gills, gives up oxygen to the blood in the capillary vessels and re ceives carbon dioxide in return. Obviously a respiratory apparatus of this nature would be quite useless to an air breathing animal. But the branchial arches of mammalian embryos, which seemingly at first are designed for the development of gills, are devoted to other purposes. They enter into the formation of the lower jaw, the muscles and bones of the face, the structures of the neck, the ear, mouth, pharynx, and the thyroid and laryngeal cartilages. The branchial clefts disappear almost entirely, but the entoderm of the pharyngeal pouches gives rise to the thymus, the parathyroid glands and part of the thyroid gland

Other examples to illustrate the law of recapitulation, such as the persistence in the human adult of the useless ear and tail muscles and persistence in the human adult of the useriess ear and tail influences the outmoded vermiform appendix, may be mentioned Again, the heart of the early embryo is a simple tubular structure as in fish, but later, it becomes bent upon itself and develops separate chambers. For a short time it closely resembles the heart of an amphibian such as the frog, possessing a sinus venosus and other features of the cold blooded.

heart

As development proceeds, the human embryo ascends, as it were, the evolutionary ladder passing rapidly over the lower rungs, more slowly over the higher For some time it resembles its mammalian kind, not until about 6 weeks after fertilization of the ovum does it show distinctly human characteristics. The human embryo possesses a tail which only commences to disappear in the seventh week

The fetal circulation. The fetal heart drives the blood which has a low content of oxygen and a high content of carbon dioxide, to the placenta through two vessels-the umbilical arteries The oxygenated blood is returned from the placenta to the fetus by a single vessel—the umbilical vein (see Pl 16A) The umbilical vessels run together, coiled or twisted with one another and covered by a soft jellylike substance,

a A child is sometimes born with a tail but it is a developmental anomaly of extreme rarity

to constitute the umbilical cord. This enters the body of the fetus about the middle of the abdomen; its point of entrance is marked in after life by a circular, depressed, and puckered area of skin called the navel or umbilitus.

The oxygenated blood of the umbilical vein passes, in part, through the liver to enter the *inferior vena cava* of the fetus, but the greater proportion pours directly into the inferior vena cava, wherein it mixes with blood returning from the lower limbs and abdomen. This mixed blood empties into the right auricle; most of it is then directed through an opening in the interauricular septum, called the *foramen ovale*, to the left side of the heart, but a much smaller part passes into the right ventricle. The blood which passes through the foramen ovale, upon reaching the left ventricle, is discharged into the aorta; a part is distributed to various parts of the body; the remander is carried by the umbilical arteries to the placenta, where it is reoxygenated from the mother's blood.

The blood returned from the upper part of the body (head and neck, upper limbs and thoracic walls) enters the right auricle through the superior vena cava. It then passes into the right ventricle (without mixing, apparently, with the stream from the inferior vena cava passing to the left auricle through the foramen ovale). Upon reaching the right ventricle, it is pumped into the pulmonary artery, but only a small proportion is distributed to the lungs—an amount sufficient only for the nourishment of the pulmonary tissue. The greater part is short circuited, through a vessel known as the ductus arteriosus, into the aorta. Thus the great bulk of the blood delivered to the right side of the fetal heart is "shunted" to the arterial side by the short cuts provided by the foramen ovale and the ductus arteriosus.

With the first few respirations, the lungs of the newborn child are expanded, and the course of the circulation becomes altered to meet the requirements of an air-breathing organism leading an independent existence. The foramen ovale closes and the channel afforded by the ductus arteriosus becomes obliterated. All the blood reaching the right auricle is now directed through the pulmonary circuit. The umbilical vessels shrink and are converted to solid cords. One or other or both of the circulatory adjustments which direct the blood through the lungs at birth may fail to occur. If the ductus arteriosus remains pervious part of the arterial blood, owing to its higher pressure, is driven from the aorta into the pulmonary artery; also, if the foramen ovale does not close, blood may pass from the left to the right auricle. These abnormalities cause, as a rule, little or no cyanosis. But in other types

of circulatory defect, such as a large gap in the interventricular septum combined with narrowing of the pulmonary artery, the subjects —so\_called\_"blue\_babies".—show.intense\_cyanosis.

If the reader has followed the rather intricate course of the fetal circulation, he will have observed that the fetal tissues are never furnished




Fig 14 19 A comparison of the dissociation curves of fetal and maternal hemoglobins.

with fully arterialized blood; the blood is always less saturated with oxygen than the arterial blood after birth Yet strange to say, the blood returning from the placenta-that is, the blood in the umbilical vein -has a higher oxygen saturation than the maternal blood in the placenta-the only source from which the fetal blood can obtain oxygen. This physiological paradox has been explained by the discovery that the oxygen dissociation curve of fetal hemoglobin is different from that of animals in postnatal life (p. 250). The oxygen dissociation curve of the blood of the


fetus is steeper than that of the mother—that is, it lies more to the left. This means that at a given oxygen pressure (tension) the fetal blood absorbs a greater volume of oxygen than can the mother's blood, or than the baby's blood will be able to absorb after birth (see Fig. 14.19) Through such a provision the fetus is more adequately furnished with oxygen than would otherwise be possible.

Pregnancy and parturition. Pregnancy or gestation is the term applied to the period in the reproductive cycle occupied by the growth and development of the new organism within the body of the mother—that is, from the fertilization of the ovum (conception) to the birth of the young animal (parturition). Most of the structural and physiological adaptations of the pregnant state, such as the fertilization and implantation of the ovum, and the enlargement of the uterus and the growth and development of the mammary glands, have been touched

upon elsewhere in this chapter. In a book of this scope little more remains to be written.

Pregnancy in the human race has a duration of about 280 days, or from 9 to 91 months. From the fertilization and implantation of the ovum to the eighth week the product of conception is called the embryo; from this time until birth it is referred to as the fetus From about the middle of pregnancy onward, movements of the fetus occur of which the mother is aware, they are referred to as quickening. Suck-

Fig 14 20 Showing the usual position of the fetus just before birth. The umbilical vein (black) carries oxygenated blood from the placenta to the fetus, the umbilical arteries (white) carry reduced blood from the fetus to the placenta



ing and spasmodic movements of the chest resembling those of respira tion have been observed in fetal animals. The unborn child lies within the sac formed by the amnion and chorionic membrane, submerged in a fairly large body of fluid (liquor amnu) which serves to protect it from sudden jars or injuries from the outside world. This membranous sac fills the uterus, and is considerably larger than the fetus itself which is thus permitted a certain freedom of movement.

Toward the latter part of pregnancy, the unborn child usually takes up a position with its head directed downward and, most frequently, its posterior part (occiput) pointing forward and to the left (Fig. 14.20). It is fitted in this position into the cavity of the pelvis. The descent of the fetus is aided very materially by previous softening of the ligaments and capsules about the pelvic joints which are loosened thereby and rendered more pliable. Indeed, this is a process essential for normal childbirth. The birth of the child is brought about by strong contractions of the uterine muscle, aided in the later stages by voluntary contractions of the abdominal muscles. The contractions of the walls of the uterus, weak at first and of short duration, become stronger and more prolonged in an hour or two; they cause molding of the infant's head and gradual dilatation of the outlet of the uterus with drawing up and disappearance of its neck or cervix; at the same time the vaginal walls become more pulsable and distensible (Fig. 14.2.1). Later,




Fig. 14 21 Showing the thinning of the lower segment of the uterus and the dilatation of the out-tet during the first stage of labor. Left, before the commencement of labor, right, toward the end of the first stage

the membranous sac enveloping the fetus bulges into the vagina, and after a time ruptures, a part of the amniotic fluid then escaping. When the canal from the uterus to the exterior has enlarged sufficiently to allow the passage of the baby's head, the latter is expelled by powerful contractions of the uterus, accompanied usually by contractions of the abdominal muscles. The rest of the child's body follows almost immediately.

For a time the newborn babe still remains attached to the interior of the uterus through the umbilical cord and placenta (Fig. 14 22). The physician ties the cord with tape close to the baby's body and divides it on the mother's side of the tape Not until 15 or 30 minutes later is the placenta, and the attached sac composed of the chorionic and amniotic membranes expelled. The placenta and membranes are commonly referred to as the afterbirth. Labor or parturition is the term given to the series of events bringing about the emptying of the uterus and the termination of pregnancy.

A consideration of the factors determining the onset of labor. The

termination of pregnancy at the usual time is not due to a single factor, but to several. A sufficient number of these factors are known to enable us to draw in outline a picture of the mechanism leading to the precipitation of labor. The picture must of necessity be incomplete and many details left blank to be filled in when more knowledge has been gained.

It has been stated (p. 653) that the growth of the uterus ceases between the sixth and the seventh months of pregnancy, this is attributed




Fig 14 22 The uterus and child immediately after birth Note the separation of the placenta and the commencement of the peeling off of the membranes

to the rise, from this time onward, of the concentration in the blood of estradiol which has an inhibitory effect upon the growth response of the uterus to distension. But the fetus is growing rapidly at this time, and the distension of the uterus now takes place in the long axis of the organ. The tension set up in the uterine tissues tends gradually to compress, narrow, and finally obliterate many vessels supplying blood to the placenta. Degenerative changes proceed in the latter, and

the nutrition of the fetus suffers, it must be released and take on a new mode of existence if it is to survive. At the same time the output of placental hormones is reduced and the corpus luteum undergoes de generation Thus the depressing effect of progestin upon the irritability of the uterine muscle-an effect very necessary in early pregnancyis withdrawn The high concentration in the blood of estradiol further increases uterine irritability and thus enhances the tendency of the uterus to contract under slight provocation. Estradiol also raises the sensitivity (lowers the threshold) of the uterus to the action of the oxytocic principle of the posterior lobe of the hypophysis A further word should be said about this latter principle. Its liberation is appar

## INCREASE IN ESTADIOL CAUSING

1 CESSATION OF LITERINE GROWTH




Fig. 14.23 Diagram summarizing the factors leading to the onset of labor

ently under nervous control for in animals stimulation of the vagina or uterine cervix causes contractions of the uterus, a response which does not occur if the stalk of the hypophysis, along which nerve fibers reach the posterior lobe, has been first divided. An oxytocic substance is found in the blood and urine of women in labor, though there is a strong suspicion that this substance is derived from the hypophysis, such has not been definitely established. For this reason, it is not pos sible to estimate accurately the importance of the oxytocic principle of the pituitary body in the mechanism which brings on labor. Fur thermore, though labor is sometimes prolonged and difficult in animals after hypophysectomy, in other cases the young are born easily and

within the usual time after the pituitary has been removed. It can therefore be said, at any rate, that the pituitary principle is not absolutely essential to the labor mechanism (see Fig. 14.23).

A few facts in the physiology of the fetus and the newborn child. The beating of the fetal heart can be heard in the later part of pregnancy by means of a stethoscope placed on the mother's abdominal wall over the uterus. The heart rate runs from 140 to 200 per minute. In the new born it is around the former figure. The respirations of the new born are from 30 to 35 per minute, as compared with the adult rate of from 18 to 20. The liver as well as the kidneys of the new born are functionally immature. The infant's urine, like that of certain lower orders of animals (e g, the frog) is hypotonic, indicating that the ability possessed by the adult kidney to concentrate the glomerular filtrate has not yet been acquired (see Chap 7) The high erythrocyte count with which the baby comes into the world and the difference between the hemoglobin of the fetus and that of the new born or of the adult have been mentioned elsewhere. The destruction of the excess of red cells shortly after birth, combined with the functional immaturity of the liver, accounts for the slight jaundiced tint of the skin sometimes seen in the newborn infant. The intestine of the new born contains a dark brown or black tarry material known as meconium; it is quite sterile, and for this reason, the baby, if it does not receive vitamin K from the mother in sufficient amounts before birth, may suffer from hemorrhages which are sometimes fatal (see p 397) After the first day or so of the infant's life, the danger passes, for by this time bacteria capable of synthesizing the vitamin have reached the intestine in the food, or have been carried from the mouth in the saliva; the hemorrhagic tendency is thus corrected

Most unconditioned reflexes are well developed at birth. The accommodation reflex of the eye is, however, usually in abeyance; and, since the corticospinal tracts of the brain and spinal cord are not fully developed, the plantar response (p. 519) is of the extensor type (Babinski response).

The puerperium or puerperal period. This is the period following the third or final stage of labor during which the mother is recuperating, and the uterus with the other structures of the birth mechanism are being restored to the nonpregnant state. Immediately following the expulsion of the placenta and fetal membranes (afterbirth), the uterus contracts down to a firm rounded body which can be plainly felt by a hand pressed upon the mother's abdomen This strong gripping contraction compresses the open mouths of blood vessels and vascular

## 085 THE PHYSIOLOGY OF REPRODUCTION

puerperal period.

volution.

infection.

spaces in the uterus, and is essential to prevent excessive bleeding. Intermittent uterine contractions at this time may be distressingly painful ("after pains"). In the days which follow, the uterine muscle shrinks.

the redundant tissue being absorbed, and the uterus returns nearly to its virgin size. This process, which is called the involution of the uterus, is accompanied by a gradually lessening vaginal discharge consisting chiefly of blood and endometrial debris. The completion of the process of involution, which takes from 6 to 10 weeks, marks the end of the

The secretion of milk becomes well established usually by the end of the first day or so, and if the baby is put to the breast, the act of suckling stimulates contractions of the uterus and encourages its in-

The two chief dangers of the puerperal period are hemorrhage and

## Glossary

Pronunciation. The vowel sound of an accented syllable is indicated by the position of the accent mark, if the vowel is long the mark immediately follows it, if short, the mark follows the consonant Thus, in ab'scess, the a is short, whereas in adeno'ma the o is long Vowel sounds in all unaccented syllables unless otherwise indicated are short Further information on pronunciation is given by the mark - or . above the vowel letter, or in round brackets following the word

Plurals are given after the word separated from it by a dash

Derivations are within square brackets, the following abbreviations are used F, French G, Greek Ger, German L, Latin ME, middle English, Med, medieval OE., old English fr , from, prec , preceding (item), foll , following

Only those words are listed that have not been described adequately in the

text, or have been mentioned before they have been fully explained

A, an [privative, Greek negative, meaning the absence or removal of something, fr L privatio, deprive Used as a prefix (a before a vowel and an before a consonant), in words derived from the Greek, to indicate negation.

Ab- [L from] A prefix meaning away from, off Ab'oral [L ab away from + os (or-), mouth] In a direction away from the

mouth

Ab'scess [L abscessus, going away] A circumscribed area of tissue, usually accompanied by the signs of acute inflammation, and containing pus Cold abscess is one not showing the ordinary signs of inflammation, it is usually tuberculous

Accessory food factors, Old term for vitamins

Acidophil, acidophile [acid + G philos, fond] Having an affinity for acid, oxyphil

Ad [L.] A prefix meaning to toward, or a suffix indicating in a direction toward a part as designated by the main part of the word, such as caudad Adapta'tion. The failure to respond to a stimulus which has been applied re-

toward the tail, or cephalad toward the head

peatedly for some time. Thus, the ticking of a clock is after a time unperceived, and the pressure and rubbing of the clothes against the skin is dis regarded This is a property of receptors but not of nerve fibers

Adeno'ma [G aden, a gland + oma, indicating a tumor] A tumor of glandular

Adrenalec'tomy [adrenal + G ektome, excision] Removal of the adrenal glands

Adrenolytic [adreno (adrenal) + G lysis (lyt-), solution] Referring to the action of certain agents which annul the effects of adrenaline

Adsorp'tion [L ad, to + G sorbo, to suck up] The attraction and adhesion of

minute particles to the surface of a solid Agoph'ony [G aix (aig), a goat + phone, voice] The sound like a gost blest

ing heard in auscultation of the chest in certain diseases of the lungs Aero'bic [G aer, air + bios, life, manner of living] Referring to respiration and

other processes carried on in the presence of oxygen Ag'minated [L agmen (agmin), a multitude] Aggregated, arranged in clusters Agno'sia [G ignorance, fr a, privative + gnosis, knowledge] A condition allied to appears due to a cerebral lesion, in which the patient does not know the

meaning of the sounds that he hears (auditory agnosia), or recognize the

nature and use of ordinary things that he sees (visual agnosia), though his hearing and sight may be perfect Agran'ulocyto'sis [G a, privative + granulocyte + osis, a state or condition] A disease, often fatal, characterized by an almost complete absence of granulocytes from the circulation. Since these cells are of such importance in the defense of the body against the inroads of bacteria, severe uncontrollable infections may result. The disease is often caused by certain synthetic drugs or

antibiotics which poison the bone marrow and suppress its function Agraph'ia [G a, privative + grapho, to write] A condition allied to aphasia in which the patient cannot express himself in writing, or write from dictation.

Aldos'terone. A steroid hormone obtained from the adrenal cortex which has a more pronounced effect than has desoxycorticosterone upon the metabolism of water and salts (sodium and potassium).

All'ergy [G allos, other + ergon, work] Hypersensitivity to certain agents, e.g. pollen, in hay fever

Alve'olus of tooth [L diminutive of alveus, a trough] A tooth socket. Amne'sia [G a, privative + mnēmē, memory] Loss of memory due to cerebral

lesion

Am'phi-, ampho- [G on both sides, around, both] A prefix in words of Greek origin meaning on both sides, around, surrounding, both

Amphoter'le [G ampho, both] Referring to compounds, such as the amino acids that can act either as an acid or a base, depending upon the pH of the medium in which they are dissolved

Ampulla [G a double handled bottle] A saclike dilatation of a tube or canal,

e g, a semicircular canal of the inner ear

Ana- [G up] A prefix in words of Greek origin meaning up Anabolism (G anabole, a building up, fr ana, up + bole, to throw) The meta-

bolic process es involved in the synthesis of compounds, as in growth. Anaero'ble [G an privative + aerobic] Referring to respiration or other process

carried on in the absence of oxygen

An'alogue [fr G ana logus, analogous] An organ, or part, of one species which though differing more or less in structure from that of another species is similar in function, e g, the hoof of a horse and the fore part of the human fool

Anastomo'sis [G to furnish with a mouth (stoma)] The junction between the branches of two nerves or arteries either produced naturally in development or by surgical operation or the surgical union of two hollow organs eg. intestine to stomach, intestine to intestine, ureter to colon, etc.

An'drogen [G aner (and), a man + root gen, to produce] Any compound that exerts an effect like that of the male hormone, testosterone

An'eurysm [G aneurysma, an aneurysm, fr. eurys, wide] A localized, usually

saccular or fusiform, dilatation of an artery with weakening of its wall Arteriovenous aneurysm, communication between a large vein and an artery. with a local dilatation of the former

Angi'na (L. a sore throat] Sore throat or severe pain from any cause

Angina pectoris [L. pectus (pector), the chest] Severe, often agonizing pain in the chest radiating down the left arm, due to disease of the coronary arteries resulting in a deficiency in the oxygen supply to the heart muscle

An'nular [L annulus, a ring] Ring shaped

Anorex'ia [G an, privative + orexia appetite] Loss of appetite

An'te- [L before] A prefix in words of Latin derivation meaning before, in Antellex'ion [L. ante, before, forward + flexio, bend] A bending forward, ante-

flexion of the uterus, bending forward of the uterus, upon itself at the junction of the cervix with the body

Anteversion [L. ante, before, forward + versio, to turn] Falling forward of the uterus as a whole without bending upon itself

Anti- [G against] A prefix meaning against, opposite, antagonistic

Antiketogenic IL, anti, against + keto (ne) + root, gen produce] Indicating an action of certain foods, especially glucose, in antagonizing the tendency to

ketogenesis Antisep'sis [G anti, against + sepsis] The destruction of bacteria usually by

the use of chemicals Aphasia [G a, privative + phasis speech] A speech disorder due, not to any paralysis of the muscles normally involved in speech but to a lesion of the part of the cerebral cortex concerned with the memory of words or of the motor mechanism of articulation. There are several types motor, visual au ditory, etc. In motor aphasia, due to a lesion of the posterior and inferior part of the frontal lobe, the patient, though he may understand spoken and writ ten words, cannot speak some or any words. In visual aphasia, the ability to recognize printed or written words is lost, and in auditory aphasia the patient does not understand what is said to him

Ap'o [G from] A prefix in words of Greek origin meaning from

Ap'ocrine [G apo, from + crino, I separate] Indicating a type of secretion in which the gland cells discharge a part of their protoplasm

Apraxia [G a, privative + pratio, to do] The mability to perform a given act at will or when commanded to do so, this is motor apraxia. In sensory apraxia, the patient does not comprehend the use of an object, when given a pencil for example, and asked to use it, he may attempt to brush his teeth with it, or to smoke it like a cigarette

Arbonza'tion [L. arbor, a tree], Fibers arranged in a pattern like the branchings

of a tree Archen'teron (k) [G archē, beginning + enteron, intestine] The cavity of the

Arrect'or muscles (L arrector, to raise) Smooth muscles in the skin attached to

the shafts of the hairs

Arthro'dia [G a gliding joint] A joint that permits a sliding or gliding movement only

Arth'us phenom'enon. A localized anaphylactic reaction, when a sensitized rabbit is injected with the antigen, necrosis occurs and a sterile abscess forms at the site of the injection

Artic'ular [L., articulo articulate] Relating to a joint

Articulation [prec ] (1) A joint (2) Speech

Asci'tes [L. fr G askos a bag] An abnormal and excessive collection of serous

fluid in the peritoneal cavity, as may occur in hepatic, renal, or cardiac dis ease

Asep'sis [G a, privative, not + L sepsis, putrefaction] The state of being free from bacteria, or the method of bringing this about, usually by means of a high temperature

Ataxia [G a, privative + taxia order, arrangement] Muscular incoordination. Atresia [G a, privative + tress, a hole] Congenital absence of an opening e.g.,

the anus or pathological closure of a normal orifice

At rophy [G a privative + trophe, nourishment] Reduction in size of any organ or tissue due to shrinkage or wasting of its parenchyma which is largely, or at least in part, replaced by connective tissue

Anricle [diminutive of L auris, ear] The pinna of the ear or the older term

for an atrium of the heart Auscultation [L. ausculto (atus), listen to] A method of examination in which the examiner listens, usually with a stethoscope, for sounds within the chest

or any other part of the body Autocoid [G autos (o) + akos a remedy] An active principle secreted by one

of the endocrine organs. It may be either a hormone or a chalone Aut'o- [G autos, self] A prefix in words of Greek origin meaning self

Autotransplanta'tion. The transplantation of an organ or the grafting of a ussue into another part of the same body. The transplanted structure has a good chance of surviving

Avitamino'sis [G a privative + vitamin + osis, a state or condition] Defi ciency of one or other of the several vitamins, thus, one may speak of avita minosis A, avitaminosis B., etc

Az'ygos [G a, privative + zygon, a yoke] Single, unpaired

Babln'ski response. Extension of the great toe upon gently stroking the inner side of the sole with the finger or a pencil It is seen in certain nervous dis eases, e.g., interruption of the corticospinal tracts. The normal response to

such a stimulus is plantar flexion of the great toe Bacteriol'ysin [foll] An agent which causes bacteriolysis, some formed in the body act specifically upon the type of bacteria which causes a certain disease

Bacteriol'ysis [bacteria + G lysis, solution] Destruction of bacteria by breaking

up or dissolving the cell. Bacteriostat'ie [bacteria + G statikos, causing to stand] Referring to agents e g, sulphonamide drugs and antibiotics which do not destroy bacteria or do so as a secondary effect, but interfere with some normal and essential process in the bacterial cell, and thus arrest or inhibit their multiplication

bel [after Alex Graham Bell, inventor of the telephone] A unit of loudness

Bi-, bis- [L twice] A prefix meaning twice, double

Bl'ceps [L bi, double + caput, head] Two heads designation for a muscle or other structure with two origins or heads-e g, biceps of arm or of the thigh

Bicip'ital [G biceps (cipitis), two-headed] Relating to any structure with two heads e g, the biceps brachii or biceps femoris

Bleus'pld teeth. Premolar teeth

Biliver'din [L. bilu, bile + Fr verd (L. viridu), green] A green pigment in bile

derived from the oxidation of bilirubin.

Bi'otin [G bios life] A factor of the vitamin B complex which neutralizes the effects of a poisonous protein in egg white, called avidin Animals fed a diet with a high percentage of egg white, and deprived of biotin develop a severe skin condition and other symptoms Biotin deficiency in man apparently does not occur

Rias'tula. Blastocyst.

Blee'ding time. The interval, measured in minutes, elapsing between the moment that blood appears from a prick of the skin, and when it clots, as determined by dabbing the wound with a scrap of filter paper.

Bohr effect. The action of carbon dioxide in causing a shift to the right of the oxygen dissociation curve of blood; that is, in reducing the affinity of hemo-

globin for oxygen.

Bo'lus [L. a throw, a choice morsel]. A masticated mass of food in the mouth ready for swallowing, or in the esophagus or intestine that is moved along the tube by peristalsis.

Bradycar'dia [G. bradys, slow + kardia, the heart]. Unusually slow action of the heart.

Brain case. The part of the skull that encloses the brain.

Bronchiolitis (k). Inflammation of the bronchioles.

Bronchi'tis (k). Inflammation of the bronchi.

Cal'cify [L. calx, lime + facio, make] To harden a tissue, such as the arterial wall in arteriosclerosis, by the pathological deposition of insoluble salts of calcium, or the incorporation of calcium into cartilage or membrane in the normal formation of bone.

Calc'ulus [L a pebble]. A stone formed in any organ (usually hollow), such as

the gall-bladder, ureter, urinary bladder, salivary duct, etc

Canalic'ulus [Diminutive of L. canalis, a canal] A minute canal Can'thus [G. kanthus]. The junction of the cyclids at the outer and inner angles of the eve.

Car'dia [G. kardia]. The cardiac orifice.

Caryokinesis. See Karyokinesis.

Caryoplasm. See Karyoplasm

Castra'tion [L. castro (-atus), to deprive of genital organs]. Excision of the gonads, either male or female.

Cara, Kara [G. kata, down]. A prefix in words of Greek origin meaning down,

as opposed to ana, up.

Catab olism [G. cata, down + bolē, throw]. The metabolic processes involved

in the breakdown of compounds to simpler ones, see Anabolism.

Catal ysis [G. cata, down + lysts, solution, dissolution]. The process whereby

certain agents (e.g., enzymes), while not entering intimately into a chemical reaction or forming part of the final product, increase the speed of the reaction.

Caud'a equi'na [L. cauda, a tail + equinus, relating to equus, a horse] The bundle of nerve roots arising from the lower part (lumbar region) of the spinal cord resembling a horse's tail and which descends for some distance in the spinal canal.

Celiac (se'liak) [G. koilia, belly]. Relating to the abdomen.

Cellulitis (seluli'us) [L. cellula, a cell + itis, meaning inflammation] Inflammation of connective tissue.

Ceph'nlad [G. kephalē, head + L. ad, to]. In a direction toward the head, or

Cephal'le [prec.]. Relating to the head, or brain.

Cerebroside (ser'ebrosid). A lipid containing the sugar galactose.

Chalone (kalôn) [G. chalāo, to relax]. An internal secretion with an inhibitory action; e.g., enterogastrone.

Chem'orecep'tor. A receptor, as in the carotid body, that responds to chemical stimuli.

Cheyne-Stokes respiration (chân stôks) An abnormal type of breathing occurring in grave disease in which the respirations increase progressively in deph and vigor to a maximum, then diminish again and finally cease (apuea) for a time. The return of the breathing marks the beginning of another cycle

Chlor ophyll [G chloros, green + phyllon, a leaf] The green pigment of plant life which functions in photosynthesis, it is allied to hemoglobin

Choana—ae (ko'an ah) [G choane, a funnel] A funnel-like passage, such as

the posterior apertures of the nasal cavities Choline (ko'len) A factor of the vitamin B complex deficiency which results in

fatty liver Deficiency, however, does not appear to occur in man

Chorda tympani nerve (kor'da) [L. chorda a chord or string + tympani of the

tympanum (fr G rympanum, a drum)] A branch of the facial nerve that passes through the tympanum of the ear and supplies the submaxillary gland and the anterior part of the tongue (taste fibers)

Chromatin (hor)mafin) theory (he sales) + 1 dins (fr. ad + finit, boundary,

Chromaffin (kro'mafin) [chrom (-ic salts) + L affints (fr. ad + fints, boundary, end), meaning attraction] Having an affinity for the salts of chromic acid c g, the cells of the adrenal medulla and of the carotid body which are stained yellowish brown

Chro'matid One of the parts resulting from the longitudinal splitting of the chromosomes in the prophase of mitosis

Chro'moproteins [G chroma, color + protein] Proteins whose molecules con

tain a pigment group, example hemoglobin
Chronaxía, chronaxíe (krôn ak'se a, krôn ak'se) [G chronos, time + axia,
value] The length of time, measured in thousandths of seconds, required for
an electrical stimulus to act in order to excite a tissue, such as nerve or muscle, when the strength of the stimulus is twice that of the rheobase

Cleatrix, cicatrices (sik a tricks, sik-a tri sez) [L sear] A sear

Circum- [L. around] A prefix meaning around, surrounding, circular

Clearance, See Plasma clearance

Cosgulation time. The time elapsing between the instant that a sample of blood is collected and the appearance of the first signs of clotting. There are a number of methods for determining this, but whatever method is used the conditions are standardized so that all results by that method are strictly comparable. The congulation time of human blood is normally between 3 and

4 minutes Coelom (selom) [G koilôma a hollow] The cavity of the embryo formed between the layers of the mesoderm, and from which the pericardial, pleural, and peritoneal cavities are developed This is the intra-embryonic coelom it communicates with the cavity of the chorion which is called the extra-embry-

onic coclors Cof'or In'dex. An index employed in distinguishing between different types of anemia that expresses the relative quantity of hemoglobin in each red cell. It

is a quotient derived from the following fraction

Hemoglobin in percent of normal

number of red cells per millimeter of blood in percent of normal.

The index is 10 in normal blood, above this in hyperchronic anemias, and below in hyporchronic anemias, and below in hyporchronic anemias.

Commissure [L. commissura, a joining, a seam] Any band of fibrous, nervous or other tissue that joins two parts of (usually) similar structure

Complemental air. The volume of air that can be inspired above that of a normal ordinary inspiration

Congen'ital [L. congenitus, born with] Relating to any disease, condition, struc-

ture, etc. present at birth.

Con jugated pro tems [L con jugo ( jugatus) join together] Any protein con taining in its molecule a pigment carbohydrate lipid nucleic acid or other nonprotein group e g hemoglobin glycoprotein nucleoprotein etc

Con tra [L against] A prefix meaning against opposite Convection [L. con veho ( vectus) carry or bring together] The change in the positions of particles in a gas (e.g. air) or liquid caused by temperature the heated particles rising while the cooler particles being heavier fall Thus

convection currents are set up

Corpus-corpora [L body] A body Cortex [L. bark] The outer part of a structure that is d stinguishable from the

more centrally placed tissue-called the medulla

Crepita tions [L fr crepo to rattle] Fine grating rattling or crackling sounds characteristic of certain rales or the gritty grinding or clicking sensation g ven to an examiner s fingers laid upon a fractured bone. They are caused by the fragments rubbing upon one another and can usually be felt by the pa

tient himself Crib riform [L cribrum a sieve + forma form] Perforated sieve like

Crossing over The interchange of chromatids between the members of a chro

mosome pair which occurs in the prophase of mitosis Cum, com, co- [L with] A prefix meaning with combined jo ned

Cune form [L cuneus a wedge + forma form] Wedge shaped

Cuta neous [L cutis skin] Relating to the skin

Cyst [G kystis a bladder] Any sac like structure

Cysti tis [G kystis bladder + itis meaning inflammation] Inflammation of the

Cytol ogy [G kytos cell (hollow) + logos word d scourse treatise] The study

Cytol ysis [G kytos cell (a hollow) + lysis breaking up solution dissolution] Disintegration solution of cells

Deamina tion, deaminization [L de from + am (fr ammonia)] The splitting

off of an amino group as from an amino acid Decibel (des i bel) [L deci (abbrev decimus) ten + bel] One tenth of a bel Deciduous (de-cid u us) [L decido to fall down] Pertaining to anything that

falls off or is discarded e g the first set of teeth or the part of the endome trium that is shed with the placenta or during menstruation

Decussation [L decusso make in the form of an X] A crossing as of nerve

fibers with their fellows of the opposite side Dehydra ton [L de from + G hydor water] The removal of water from the

body or from any structure or substance Del toid Shaped like the Greek capital letter delta(A)

Desquima tion [L de from + squama a scale] The shedding of the superficial layers from the surface of the skin or of any part covered by squamous epi

Deutoplasm deuteroplasm (du to plazm) [G deutero second + plasma a something formed] The nutritive yolk of an egg. It is very small in the ova of mammals but relatively enormous in the eggs of birds. The rest of the yolk from which the embryo is developed is called the cytoplasm or the

Di, dis- [L apart sundering into two parts two] A prefix meaning two or

separating into two Dia [G through] A prefix meaning through in different directions or com pletely

Diapede sis [G dia through + pedesis a leaping] The passage of leucocytes through the capillary wall

Disth esis [G an arrangement] A constitution predisposed to some disease or abnormal state

Dichotomy [G dicho in two + tome a cutting] A branching into two hence dichotomous in referring to such a division

Dierot le pulse [G dierotor fr dt two + erotor a beating a pulse] A type of pulse occurring in typhoid fever and certain other cond tions associated with low blood pressure and lax arterial walls in which the dierotic wave (ordinarily not palpable) can be felt as a distinct tap following the main pulse wave

Dig'it (L. digitus) A finger or a toe

Digitation [L. digitus a finger or toe] One of a series of fingerlike projections especially of a muscle

Dip loe [G feminine of diplous double] The cancellous layer of bone lying be tween the two layers of compact bone of the cranium

Dip loid [G diplous double + eidos resembling] The double number of chromosomes characteristic of the immature sex cells the zygotes and the body cells generally see Haploid

Dist'al (L. dist (ant) + all In a position farther from the center or midline of the body or from the origin of a limb or structure such as an artery or nerve thus the elbow is distal to the shoulder

Dominant IL for dominus a master [1]. Relating to an hereditary character which appears in the offspring and masks another character called recessive though the gene carrying the latter has also been transmitted [2]. Relating to the cerebral hemisphere which determines right or left handedness. Thus the left hemisphere is dom nait in right handed persons and the right hemisphere those who are left handed.

Dors'ad [prec ] In a d rect on toward the back.

Dor'sal [L. dorsum the back] Relating to or in a position toward the back

Drop sy [G hydrops an abnormal collection of fluid in a tissue or cavity] An old term for edema or ascites and still in popular use any abnormal and excessive collection of fluid in the issues or a body cavity.

Dine [G dynamis force] A unit of force defined as that which acting on a mass of 1 gram for 1 second causes an acceleration of 1 cm per second.

Dys. [G bad difficult] A prefix in words of Greek derivation meaning difficult bad abnormal or distressing

Dysarthr'in [G dys bad + arthrão articulate] Difficulty in speaking due to weakness or paralysis of the muscles of articulation or to stuttering or stammering

Dyspha gia (G dys bad + phageun to eat). D fliculty in evallowing owing to pain or any other cause

Ec-, ex [G ektos without out of] A prefix meaning out of outside removal of Ectop ie [G ek out + topos a place] Misplaced not in the normal situation e g ectopic gestation pregnancy occurring in a uterine tube or in the peri toneal cavity

Edema [G orderna a swelling] An abnormal increase in tissue fluid especially in the subcutaneous tissues causing a puffy swelling of the part which when pressed with a finger or thumb shows a hollow imprint due to the temporary dispersal of the fluid

Elec trocor'tin [electro (abbrev electrolyte) + cortin] Aldosterone

- Electrophore is [Electro + G phoresu a carrying] The movement of particles such as protein molecules in an electric field to the anode or cath ode
- Embolus [G a wedge or stopper] A clot mass of agglulmated cells a piece of fibrin etc carried in the blood stream and blocking a vessel
- Em'bryo [G embryon fr en in + bryo be full to swell] The developing or gains from the fertilization of the ovum to emergence from the egg as in birds in man the product of conception from the second to the eighth week of pregnancy
- Embryol ogy [G embryon (prec) + logos treatise discussion word] The study of the development of the embryo and fetus
- Empirical [G eperikos experience] Not based upon any known scientific fact but purely upon experience
- Empye ma [G empyema suppuration fr en in + pyon pus] Pus in any cavity but especially in the pleural cavity
- En [L. in] A prefix meaning within
- Endem ic [G endemos a native fr en in + demos the people] Relating to a disease of which one or more cases are always present in the particular district under consideration
- Endo- [G endo within] A prefix meaning in within
- Endog enous (1) [G endon within + root gen beget produce] Formed within the body
- En terogastro'ne [G enteron intestine + gaster belly] A principle (chalone) formed in the mucosa of the duodenum which under the influence of fatty materials in the chyme passes into the blood stream and inhibits the secretion of gastric juice and the movements of the stomach
  - En to [G entos in] A prefix meaning within
  - Epi [G upon] A prefix meaning upon above on top of the upper side
- Erg [G ergon work] A un t of work, defined as that performed in moving a mass against a force of 1 dyne
- Estrogen [G oistros mad desire + root gen produce] Any substance that in duces effects similar to those of estrad of
- Eu [G well good] A prefix in words of Greek origin meaning good well or normal
- normal Evagina tion [L. e out + vagina a sheath] The protrusion of the wall or other
- part of any structure

  Exfolia tron [L ex out + folium a leaf] The shedding of superficial parts of
- any structure especially in layers or sheets a form of desquamation Exocrine [G exo outside + krino to separate (secrete)] Relating to an exter
- nal secretion e.g. saliva gastric juice etc. that is secreted on to the surface of the body or into one of the body cavities.

  Exogenous () [G exo outside gen produce] Formed or originating outside
- Exophthal mos [G ex out + ophthalmos the eye] Protrus on of the eyeballs
- beyond their normal positions

  Ex'tero [L exterus outside] Same as preceding
- Ex'terolec'tive [L exterus' outs de + facto do] Professor Cannon's term for the cerebrospinal nervous system as opposed to the interofective (or autonomic) nervous system
- Ex'tra [L. outside] A prefix meaning on the outside
- Ex'udate [L. ex out + sudare to sweat] A fluid or semifluid material eg serum or pus which flows or oozes from any part

Facet' [F a very small face] A small smooth area on a bone (usually for ar ticulation) or other hard structure

Falciform (făl si form) [L. falx (falc ) a sickle + forma form] Shaped like a

Pascia (fash 1a) [L a band] A thin sheet of fibrous tissue covering a muscle or other structure or separating one structure from another

Fecunda tion (k) [L fecundo (atus) make fruitful] Fertilization

Fenes tra [L a window] A window e g fenestra ovale of the internal ear

Fe tus, foe tus IL offspring! The unborn offspring from the eighth week to the termination of pregnancy

Fibril la-ae [L fibrilla diminutive of fibre a fiber] A minute fiber

Fibrillation [prec ] Fine contractions or twitchings of muscle due to independ ent activity of the separate fibers. Such occurs in degenerating skeletal muscle in cardiac muscle in electrocution as a result of poisoning by certain drugs

and just preceding death from almost any cause Fil ament [fr L filum a thread] An extremely fine fiber or thread like struc ture

Fi lum termina le [L. terminal thread] A siender filament that descends from the end of the spinal cord to the termination of the vertebral canal

Fim briated [L fimbria a fringe] Fringed or like fringe

Fis'tula-fis'tulae [L, a p pe tube] An abnormal passage lead ng from an abscess cavity or any hollow organ or part to the exterior

Fora men-foram ins [L an aperture] A small opening or hole in a bone (usu ally) or other structure

For eign pro tein A protein not naturally present in the body e.g. a vegetable protein or one from another species it acts as an antigen

For'sa [L. a trench or d tch] A depression usually in a bone and long tudinal in shape Funiciplus (k) Ideminutive of L. funis a cord! A cord or cord like structure

such as the umb lical or spermatic cord Galvanom eter [Lu gi Galvanni + G metron a measure] An instrument for

indicating and measuring a galvanic electric current Gang'lion-gang'lia [G a small subcutaneous tumor] A circumscribed collection of nerve cells in the brain on a sensory nerve or of the autonomic

nervous system (2) A small cystlike swelling on a tendon usually at the wrist Gasse rian gang'ilon Semilunar ganglion trigeminal ganglion Gas'trula [fr G gaster belly] The embryo in the stage of development when it consists of a sac with a double layered wall the inner layer is the endoderm

the outer the ectoderm Gene [G root gen beget] One of the theoretical particles of a chromosome which carries an hered tary character

Genera tive [L generare beget] Capable of reproduction of procreation

Genet'le [G genesis (-etic) generation] Relating to heredity Genetics [prec ] The study of the laws and processes of hered ty

Ge notype [G genos birth + typos an imprint] The genetic constitut on (com bination of genes) of an organism

Genus [L. birth descent] In the classification of animals or plants the d vision next above the species 1 e below the Family or the Tribe

Glanco ma [G glaukos bluish green] Raised pressure within the eyeball due to an excess of fluid the result in turn of blockage of the drainage system. It is a progressive disease leading to failure of vision

Gliss'on's cap'sule. The capsule of the liver

Glycogen'esis [G glykys sweet (sugar) + root gen, produce] The formation

of glucose from glycogen or of glycogen from glucose Glycogenol'ysis [G glyk)s, sweet (sugar) + lysis, dissolution] The breakdown of glycogen to glucose

Glycol ysis [G glykys sweet (sugar) + lysis dissolution] The breakdown of plucose to lactic acid or to carbon dioxide and water

Gly'coneogen'esis, glu'coneogen'esis [G glyk)s, sweet (sugar) + neos new genesis origin! The formation of glucose from noncarbohydrate material, as from protein or fat

Gly'coprotein, glu'coprotein [G glykys, sweet (sugar) + protein] A conjugated

protein containing a sugar group, it is present in mucin Granula'tion(s) [L granulum diminutive of granum a grain] Masses of small

grain like bodies Granula'tion tissue [prec ] The vascular granular tissue that covers a healing open wound and becomes converted to scar tissue

Hap'loid [G haplous single] Refers to the reduced number of chromosomes in

the gametes see Diploid Heart block Disease of the atrioventricular bundle that slows or blocks conduction of the impulse from the atria to the ventricles. In complete blockage, the

atria beat at the normal rate (70 per minute), but the ventricles neat at their own rate of about 40 per minute. The arterial pulse is of course, at the ventricular rate

He'matin [G haima (haimat ), blood] The pigment (porphyin) of hemoglobin

Hemi- [G half] A prefix meaning a half

structure and composition throughout

Hemol'vsis [G haima, blood + lysis a setting free dissolution] The liberation of hemoglobin from the red cells

Hen'le's loop The U shaped part of a renal tubule

Hep'ar [G the liver]

Hepat'ic [G hepar (hepat ), the liver] Relating to the liver

Hetero- [G heteros other, different] A prefix in words derived from the Greek

meaning different from something else Heterol'ogous [G heteros different other + logos relation] Indicating a tissue or part derived from another species or of a tissue or cells in a part not nor-

mal to them Het'erotran'splant [G heteros different, other + transplant] The grafting of a tissue or organ taken from an animal of one species to the body of an animal

of another species Except for certain tissues and under very special circumstances the operation is not permanently successful, i.e., the transplanted tis sue dies Hia'tus [L. an opening, fr hio (hiatus), yawn] A gap or opening aperture or

fissure

Histol'ogy [G histos a web (tissue) + logos treatise, discussion word] The microscopic study of tissues Hol'ocrine [G holos all + krino separate (secrete)] Referring to a type of

glandular secretion which consists of the cells of the gland itself Ho'mo- [G homos same] A prefix in words of Greek derivation meaning of

the same kind or class alike Homoge'neous (1) [G homos same + genos family kind] Having uniform

- Homogéenous (j) [prec ] Having the same consistency throughout and usually the same composition
- Homologous [foll] Referring to a homologue having the characters of a homologue
- Hom ologue [G homor same + logor relation treatise word] Any organ or part of the body that resembles in structure or origin another organ or part elsewhere in the body or an organ or part in one species that corresponds in some way with an organ or part in another species such as the wing of a bird and the arm of a ma.
- ord and the arm of a man Ho motransplanta ton [G homos same + transplant] The grafting of tissue or of an organ from one person to another or from one an mal to another of the
- same species

  Höst [G hospes a host] An organism that harbors and gives sustenance to an other (usually pathogenic) organ sm
- Hu mor [L hum or (or umor) a fluid] (1) One of the body fluids such as the aqueous humor of the eye which fills the space between the cornea and the lens (2) A substance such as a secretogogue (extractive) in the food which is absorbed into the blood and stimulates a digestive secretion (3) A substance instructed in the remainder of the first of the firs
- upon which the ancient theories of disease were based (humoral doctrine)
  Hydlod [G hydlos glass + eidos a resemblance] Glass like in appearance
  Hydrocephalus (hi-drò-kef al us) [G hydor water + kephale head] Excessive
  amount of fluid in the ventricles of the brain (internal hydrocephalus) or in
- the subarachnoid space (external hydrocephalus)

  Hydrol ysis [G hydro water + lysis solution dissolution] The taking up of water by a compound which then spl is into smaller molecules. It is brought about by certain enzymes and by the action of heat and strong chemicals—
- acids or alkalis

  Hydrostat'is pressure [G hydror water + statos standing] The pressure exerted by the weight of water or other liquid
- Hyper [G hyper above over] A prefix indicating excess above the normal Hyperpla sta [G hyper above + plasts a molding] Active multiplication of the cells of any tissue
- Hyperpnea [G hyper above + pnoise the breath] Exaggerated respirations with an increase usually of the volume of air breathed per minute
- Hypertonic [G hyper above + lonus tone] Having a relatively high tone
- (e.g. muscle) or a high osmot c pressure (e.g. a solution of salt)

  Hyper trophy (G hyper above + trophe nourishment) Increase in bulk of a
- tissue or organ caused by an increase in the individual cells

  Hy'po- [G hypo under] A prefix meaning under beneath less than normal
- Hypoderm le [G hypo under] A prenx meaning under beneath less than normal Hypoderm le [G hypo under + derma skin]. Beneath the skin
- Hypoton ic [G hypo under + tonos tone] Having a relatively low tone (e.g. muscle) or a low osmotic pressure (e.g. a solution of salt)
- Impreginate [L in (im) + praegnans pregnant] (1) To introduce particles of one substance into any other substance to saturate (2) To fertilize (the penetration of the ovum by the spermatozoon) to fecundate
- Incon tinence of urine [fr L in negative + contineo hold together] Involve tary passage or dribbling of urine
- Inertia (in ursh ea) [L in negative + iners (fr in negative + ars art) slug g shness unskilled] In physics the "resistance" which a body offers to a force tending to move it from a position of rest Inertia of the uterus sluggishness or absence of contractions of the uterus during labor

- In'guinal [L. inguen (inguinis), the groin] Relating to the groin
- Inhibition (in hib-ish'un) [L. hibeo (hibitus) keep back, fr habeo. to havel . Depression or arrest of function of any organ, e g, slowing or stoppage of the
- heart by stimulation of the vagus nerve, or the lessening of any process Inorgan'ic [L in, not + G organikos, from organon, a tool, organ] Not organic, manimate, in chemistry, indicating compounds that do not contain
- carbon Inos'culate [L in + osculum diminutive of os, mouth] Anastomose
- Insemination [L insemino, sow or plant, fr semen (semin), seed] The deposition of semen in the vagina either during coitus or artificially
- In'sula IL an island! The small part of the cerebral cortex hidden in the Sylvian fissure It is disclosed by raising the adjacent overlapping parts of the frontal and parietal lobes
- Integ'ument [L. integumentum a covering] The skin
- In'ter [L. between] A prefix in words of Latin origin meaning between
- Interofective nervous system [L. inter, between + facio, do] Professor Cannon's term for the autonomic nervous system
- Interstitial (in'tur-stish'al) [L. interstitium, fr inter, between + sisto, stand] Relating to spaces between fibers, cells, etc., especially in the tissues generally, as interstitial fluid
- In'tra- [L. within] A prefix in words of Latin derivation meaning within
- In'tro- [L. into] A prefix meaning into Invag'ina'tion (1) [L in, in + vagina, a sheath] The pushing of one part of a hollow structure inwards towards its center, to ensheath
- Inversion (in vur'shun) [L. inverto (inversus), a turning about] (1) a turning into a reversed position, as inversion of the uterus in which the interior of the uterus is turned outward and the organ displaced downward (2) A turning inward, as inversion of the foot (3) The conversion of a disaccharide to a monosaccharide, e.g., of cane sugar to glucose and fructose
- I'soaggin'tinin [foll ] An antibody present in the serum of a member of one of the blood groups which agglutinate the red cells of the blood of another group, that is, causing agglutination of cells in blood of the same species
- I'soagglutin'ogen [G 1505, same, equal + L agglutino, glue together] A substance (antigen) associated with the erythrocytes that reacts with an agglutinin (antibody) in the serum of a member of the same species and causes clumping together of the cells, see blood groups
- Kar'yokine'sis [G karyon, a nut (nucleus) + kinesis, movement] Mitosis Kar'yoplasm [G karyon, a nut (nucleus) + plasma, a thing formed] The nonchromatin substance of the cell nucleus
- Kata. See Cata Ker'atin [G keras (keras), horn] A protein in the outer layers of skin, and in
- horn, nails, etc
- Kerati'tis [G keras (kerat), horn (cornea) + us, meaning inflammation] Inflammation of the cornea of the eye
- Kētogen'ie [keto (ketone) + root, gen, produce] Referring to certain foods. especially fats, which tend to cause the production of ketone bodies
- Knee-jerk. A quick extensor movement of the knee, a kick, caused by tapping the ligamentum patellae while the leg is hanging loosely in the flexed position It is absent in certain nervous diseases and abnormally active in others
- Kup'fler cells. Large star-shaped shells attached to the walls of the sinusoids of the liver They belong to the reticulo-endothelial system and are phagocytic.

Lecta'tion [L lacto (atus), to suckle (fr. lac, milk)] The secretion of milk

Lactation [L lacto (alus), to suckle (fr. lac, milk)] The secretion of milk Lacu'na, -ae [L diminutive of lacus, a hollow or a lake] A hollow space or gap

in any tissue or organ

Lamella, -ae [L. diminutive of lamina a layer or plate] A thin layer, plate, or stratum

Lanu'go [L. down, wool] The very fine hair covering the body of the embryo

or the surface of the adult body, except the palms of the hands and the soles of the feet and where the harr is long as on the scalp Larya—laryae | L. a mask. | The wormlike stage of an insect as hatched from

the egg grub, maggot, or caterpillar, or the young of any species that differs essentially from its parents in form

Larval, Larvate [prec] (1) Relating to larvae (2) Masked, applied to a disease in which the symptoms are not typical or are absent

Lesion (le'zhūn) [L' laedo (laessus), lo injure] Any injury or any structural sbnormality in a lissue, organ, or part due to disease, e.g., a tubercle of a lung in tuberculosis, or the pustule of the skin in smallpox

Leuce'mfa, leuke'mfa [G leukos, white + hanna, blood] A disease of the blood in which the white cells are greatly increased in number, there being many abnormfal and primitive forms in the circulation. The disease may be acute or chronic, and according to which type of cell predominates in the blood lymphocyte, myelocyte, or myeloblast, or monocyte. It is called lymphatic myelogenous, or monocyte leukemia.

leucope'nia [leuco (leucocyte fr G leukos, white + kytor cell) + penia poverty] A pronounced reduction of leucocytes, scarcity of leucocytes.

Lip'opro'tein [G lipos fat + protein] A conjugated protein containing a lipid group in its molecule
Lob'die It. lobulus, diminutive of lobus! A small lobe or a subdivision of a

lobe
Lumen—lumina [L a window, a light] (1) The cavity of a tubelike struc-

ture, such as an artery or the intestine (2) A unit of light Lymphangi'tis [L. lympha clear water + G angeion, a vessel + lits, meaning

inflammation] Inflammation of lymph vessels

Lysin [G Iyrr disintegration, dissolution] Any agent that destroys cells by

breaking them up or dissolving them One that destroys blood cells is a hemolysin bacteria, a bacteriolysin, and cells in general, a cytolysin Lyrsis [G disintegration, dissolution] Disintegration of any structure, red cells bacteria, fibrin etc. Also used as a suffix in words of Greek origin meaning.

breakdown or disintegration, e.g., cytolysu

Macera'tion [L macero (atus), soften by soaking] Soften or break up by soaking in water or other liquid

Macroscopic (k) [G macros large + skopeo, to view] Capable of being seen by the naked eye

Mag'enstras'se [Ger magen stomach + strasse, road] The part of the interior of the stomach extending along the lesser curvature from the cardia to the region of the pylorus that is first traversed by the food after it has entered the stomach

Maras'mus [G murasmor a withering! A state characterized by dehydration wasting, and weakness, seen especially in infants as a result of infection, or faulty feeding

Medul la [L fr medius middle] Any very soft structure or the interior of an organ e.g. the adrenal medulla as opposed to the outer part or cortex mar

row spinal cord medulla oblongata Meibomian gland (mi bo me an) One of the glands at the margin of an evel d

Meiosis (mi-o sis) [G a lessening] Reduction division Mem'brane [L a skin or membrane] A thin sheet of tissue that serves as a lining for various hollow structures (e.g. mucous membranes) covers or separates organs (e.g. fibrous membranes) or lines such cavities as the thorax and abdomen (e.g. serous membranes)

Mende lian laws The laws of heredity discovered by Gregor Mendel an Augus tinian monk of the 19th century living in Brunn Austria Studying inherit ance in pea plants he found that characters were transmitted to the offspring as independent factors or genes. When two varieties of plant, each pure for a different character (shortness and tallness) are crossed their progens (F generation) will be all tall but they received a gene for shortness from their short parent as well as a gene for taliness from the tall parent, the character tallness which appears is called dominant the hidden character shortness is termed recessive These hybrid individuals produce three types of gametes 25 percent of which contain only genes for tallness (dominant) 50 percent genes for both tallness and shortness and 25 percent genes for shortness (re cessive) only When these are interbred the offspring (F2 generation) will be 75 percent tall and 25 percent short that is only those which had not re ceived a gene for tallness from either parent will be short

Mer'ocrine [G meros a part + krino to separate (secrete)] A secretion such as the gastric or pancreatic juice which is merely a product of the gland cells

and not part of their substance see Holocrine and Apocrine

Meta [G after between over beyond] A prefix meaning after over beyond or indicating a transformation from something else that has gone before Metamorphosis [G meta prec + morphe form] A change usually radical in

form transformation as that of the tadpole into the frog

Metazoon a [G meta + zoon animal] A member of any class of animal above the Protozoa that is one composed of many cells

Mic'turate [L. micturio pass urine] Urinate

Micturation (micturish un) [prec ] Urination

Min smal air The air that remains in the lungs after their removal from the body and that gives pulmonary tissue its buoyancy

Momentum [L motion] The quantity of motion the product of mass and velocity impetus the tendency of a moving body to cont nue in its course

Mon ocyte [G monos single + kytos a hollow (cell)] A white blood cell with a single unlobed nucleus. It resembles a large lymphocyte, but its nucleus is more deeply indented being kidney or horseshoe shaped Monocytes consti tute from 2 to 8 percent of the leucocytes. They are elements of the reticulo endothelial system

Mononu'clear [G monos single + nuclear (nucleus)] Referring to a cell with

but one nucleus

Morbil li [L. morbillus diminutive of morbus a disease] Measles Motor end plate. The structure on the muscle fiber in which the motor nerve

terminates Multicellular [L. multi many] Consisting of many cells

My'eloblast [G myelos marrow + blastos a germ] A granulocyte in a very early stage of its development (in the bone marrow)

My'elocyte [G myelos marrow + kytos hollow (cell)] A granulocyte in a late stage of its development (in the bone marrow)

- Myogen'ic [G mys, muscle + root, gen produce] Originating in muscle or
- Myoneur'al [G mys muscle + neuron nerve] Relating to both muscle and nerve Myoneural junction the connection between a motor nerve ending and the muscle fiber It has the properties of a synapse
- Navic'ular (k) [L. diminutive of navis, a boat] Shaped like a boat, scaphoid Ne'oplasm [G neos, new + plasma, something formed] A new growth, a tumor composed of masses of newly formed cells

Neurogen'ic [G neuron nerve + root gen, origin] Of nervous origin

Neurohumanal [G neuron, nerve + none, qv] Relating to the theory that nervous effects are transmitted by chemical substances (humors) liberated at the nerve endings, e.g., acetylcholine

No tochord [G notos, back + chorde, cord] A flexible bar in the early embryo

around which the vertebral column is developed

Nuclease [nucleus + ase, meaning an enzyme] An enzyme that splits nucleic

acids Nu'cleopro'tein. A conjugated protein containing nucleic acid. It is an essential

constituent of chromatin

- Nu'cleus—m'clei [L. a small nut or kernel] The body containing chromatin and surrounded by the cytoplasm that is present in nearly all types of cell (2) A circumscribed mass of gray matter (nerve cells) in the central nervous system (3) a focus, such as a speck of foreign material, around which other material collects
- Material concerns

  Nystag'mus [G nystagmos, a nodding] A rhythmical oscillation of the eyebalis
  occurring in certain diseases, and normally after rapid rotation of the body
  The movements are most commonly from side to side, but may be up and
  down or rotary
- Ob- [L against, toward] A prefix in words of Latin origin meaning against or toward
- Odont'old [G odous (odont), a tooth + eidos a resemblance] Shaped like a
  - -old [G eidos, a resemblance] A suffix in words of Greek derivation meaning a likeness to resembling

Oligu'ria [G oligos, few, little. + ouresis urination] Scanty urine

-oma A suffix added to words of Greek derivation to indicate a swelling or tumor of any kind e.g., carcinoma, sarcoma

Ontogenesis, ontogeny [G on being + genesis, origin] The developmental his tory of an individual as opposed to the history of a race See Phylogeny Ontology [G on (ontor), being + logor, discussion, treatise] The study of the

Ontology [G on (ontos), being + logos, discussion, treatise] The study of the development of an individual embryology

Onychia [G onyx (onych-), nail] Inflammation of the root of a finger- or toenail with suppuration and usually shedding of the nail

O'ocyte [G con an egg + kytos, a hollow (cell)] The primitive ovum in the Graafian follicle.

Op'sonins [G opson, sauce] Substances in blood that stimulate the leucocytes (neutrophils) to attack bacteria—fancifully speaking they make the latter more appetizing

Oral [L. os (oris), mouth] Relating to the mouth

Order. A division in the classification of animals or plants between the Class or Subclass (above) and the Family (below)

- Organ'ıc [G organikos relating to an organ, from organon a tool, organ] (1) In chemistry, relating to compounds containing carbon (2) Relating to a disease associated with a structural change in a tissue or organ as opposed to functional (3) Pertaining to animal or vegetable life, as opposed to the morganic world
- Or'ganism. Any living thing, animal, or plant Orthope'dic [G orthos, straight, right, true + pais (paid), a child] Relating to
- the study and treatment of chronic diseases of bones and joints and the correction of deformities
- Orthostat'ic [G orthos, straight, right, true + statos standing] Relating to the upright (standing) position
- Os-ora [L mouth] A mouth
- Os-ossa [L a bone] A bone
- Osteogen'esis [G osteon bone + genesis production] Bone formation
- Ovariec'tomy [L ovarium, ovary + G ektome, excision] Excision of the ovaries, castration of the female
- Ox'yphil [G oxys, sharp, acid + philos, fond] Acidophil, eosinophil Oxytocin (oks 1 to'sin) [G oxys, swift + tokos, childbirth] Pitocin
- Palpa'tion [L. palpatio, fr palpo, to touch] Determining, usually with a finger or the fingers or palm of the hand, by touching
- Para- [G alongside, near, beside] A prefix meaning beside, on the side of, not
- quite normal Para-aminobenzoic acid. A factor of the vitamin B complex essential for the growth of certain microorganisms. Some sulfonamides exert their antibacterial action by rendering this vitamin unavailable for the infecting bacteria
- Paren'chyma [G anything poured in beside] The part of an organ or tissue composed of cells that perform the specific function of the organ or tissue, such as the contractile fibers of muscle or the secreting cells of a gland, as opposed to the nonspecific supporting tissue or stroma of connective tissue
- Paren'teral [G para beside, contrary to + enteron, intestine] Referring to a route of administering a drug or agent of any kind other than by the gastro intestinal tract, that is, subcutaneously, intravenously, etc.
- Path'ogen [G pathos, suffering, disease + root gen produce] Any organism which causes disease, hence, pathogenic
- Pectoril'oquy [L. pectus (pector-), breast, chest + loquor, to speak] In auscultation of the chest, the transmission of the sounds of the patient's voice to the examiner's ear
- Ped'icle [L a little foot, diminutive of pes (ped-)] A stalk
- Pepsin'ogen [pepsin + G root gen produce] The inactive enzyme secreted by the gastric glands and converted to active pepsin by acid
- Per- [L. through] A prefix denoting through, around, (in chemistry) the highest compound in a series
- Percussion (pur kush'un) [L. percussio, a beating] A method of physical examination, especially of the chest, in which the nature of underlying structure is
- determined from the sound caused by tapping with an instrument or finger Per'i- [G around beyond, surrounding] A prefix meaning around, surround ing, exceedingly
- Peri'pheral [G peri, around + phero, I carry] In a position away from the center of the body or of any organ or part, nearer the circumference Pe'trous [G petra a rock] Hard, like stone
- Phe'notype [G phaino, display + typos, model] The appearance, structure, coloring and other characteristics of an organism

- Phlebi'us [G phleps (phleb) a vein + itis meaning inflammation] Inflammation of a vein
- Phlebot'omy [G phleps (phleb) a vein + tonie a cutting] Incision into a vein, as for withdrawing blood
- Pho'bia [G phohor fear] An unreasonable fear of some particular thing or condition For example claustrophoba is a morbid fear of any closed space agoraphoba fear of open spaces The term is also used to indicate hypersen situity to some form of stimulation e.g. photophoba pain in the eyes and spasm of the lids to a light which normally would eause no discomfort.
- Phos phagen [phosphorus (phos light + phoros, bearing) + root gen] Crea tine phosphate phosphocreatine
- Pho'topho'has [G phos (phot) + phobia] Hypersensitivity of the eyes to light Pho'tosyn'thesis [G phos (phot) + synthesis] The formation of carbohydrate from carbon dioxide and water, and the liberation of oxygen by sumbirth
  - ing upon chlorophyll

    Phren'ic [G phren diaphragm] Relating to the diaphragm
- Phrem'tis [G phren diaphragm seat of the emotions (an ancient belief) + itis inflammation] Delirium inflammation of the brain
- Phylogen'esis, phylog'eny [G phyle, phylon a tribe + genesis origin] The de velopment (evolution) of a race of animals or plants see Ontogenesis
- Phy'lum [L fr G phylon a tribe race] In the classification of animals or plants a large division below the Subkingdom and above the Subphylum or the Class
- Pl'siform [L. pisum pea + forma form] Resembling a pea in shape and size
- Plank'ton [G planklos wandering] Minute free swimming or floating marine organisms either of the animal or vegetable kingdom Plant at [L planta the sole of the foot] Relating to the sole of the foot
- Plas'ma clea'rance A term for the volume of plasma from which a substance
- (such as urea) is completely removed by the kidney and excreted in the urine It is determined by dividing the figure for the total quantity of the substance excreted in the urine per minute by its concentration in the plasma
  - Thus  $\frac{UV}{P}$  = plasma clearance U is the concentration of the substance (e.g. urea), V is the volume of the urine secreted per minute and P is the
- (eg urea), V is the volume of the urine secreted per minute and V is the concentration of the substance in the plasma

  Pleth'ora [G plethore fullness] A fullness an excess especially of blood con
- gestion
  Piex'us [L a braid] An interlacement or network of nerves (nerve plexus) or
  of years (venous plexus)
- Pli ca, ne circularis es [fr L plica a fold + circularis in the form of a small circle] One of the transversely circular folds of mucous membrane in the small intestine
- Polyu'ria [G pol) many, much + ouron urine] Excessive secretion of urine Post [L after] A prefix meaning after, behind
- Prae See Pre
  - Pre [L prae before] A prefix meaning before in front of
  - Privative [L. privatio, to deprive negation] The letter a used as a prefix in words of Greek origin to indicate negation, not as atypical not typical
- Pro- [G before] A prefix denoting before forward Pro-cess [L processus an advance or progress] (1) A physiological or pathological mode of action (2) In anatomy, a projection from the main part of

any structure-e g bone nerve cell (axons and dendrites), etc

- Prolapse (prō-laps) [L. prolapsis a fall ng] The descent of an organ from its normal position e.g. prolapse of the uterus
  Proliferation [L. proles offspring + fro to bear] Multiplication of the cells of
- Proliferation [L. proles offspring + fero to bear] Multiplication of the cells of any tissue usually in the sense of excessive
- Protozo'on, -a (G protos first + zoon an animal) An animal composed of a single cell unicellular animal
- Prox Imal [L. proximum next nearest] Nearer to the center of the body or to the origin of any structure—e.g. artery nerve limb etc. as opposed to distal
- Pterygoid (ter'īgoyd) [G pteryx (pteryg-) a wing + eidos a resemblance]
  Resembling a wing
- Pto sis [G a fall ng] A falling down prolapse of any structure but especially drooping of one or both evelids
- Pulse pressure The pressure of the arterial system in excess of the diastolic pressure. For example, if the diastolic pressure amounts to 80 mm of mercury and the systolic pressure to 120 mm, the pulse pressure will be 40 mm of
- mercury

  Purpura (pur pū rah) [L fr G porphyra purple) A hemorrhagic state in which spontaneous bleeding occurs into the subcutaneous tissues, deeper lay ers of the skin into or from the surface of mucous membranes or into joints
- and other structures It is often due to a deficiency of platelets
  Progenic [G pyon pus + root gen to produce] Relating to any agent that
  causes the formation of pus especially certain bacteria (e.g. staphylococci)
- Pyram idal Shaped like a pyramid Pyr'iform [L. purum a pear + forma form] Pear shaped
- Pyrogen ic [G pyr fire + root gen to produce] Heat producing causing fever
- Quad neeps [L quadri four + caput a head] Four heads term applied to the large muscle on the front of the thigh (quadriceps extensor)
- Quadrigem Inal [L. quadr. four + gentilus twin] Referring to any structure having four similar parts but especially to the four bodies on the dorsal as pect of the m dbrain (quadrigeminal bodies or colliculi)
- Quin sy [ME corruption of Med] L. quinancia fr G. cyanel e sore throat dog throttling dog collar fr kynon dog + anclein to throttle] Inflammation of the tonsits with supportation (suppurative (snishlits)
- Rale (rahl) [Fr rattle] A crackling clicking bubbling rubbing or whistling sound heard upon auscultating the chest in certain d seases of the lungs
- sound heard upon auscultating the chest in certain d seases of the lungs Ramus, i[L a branch] A branch as of an artery or nerve or a bar like part of a bone such as the ramus of the mandible
- Reac'tion [re + action] (1) A term indicating the acidic neutral or base na ture of any material—e.g. acid react on neutral react on basic reaction (2) The response of the body or of any tissue especially muscle and nerve to a stimulus (3) In chemistry the interaction occurring between two or
- more chemicals in which an exchange of molecules results in new compounds being formed Recessive [L. recedo (recessus) draw back]. An hereditary character that is
- latent or masked by another character called dominant
  Regurgitation (regurj it a shun) [L. re back + gurgito (atus) flood] The
  movement of material in a d rection the reverse of normal as the passage
  of the stomach contents into the esophagus (usually without vomiting) of
  duodenal contents into the stomach or of blood from the norta during das

tole into the left ventricle

Rels'sner's mem'brane. The membrane of the internal ear that separates the scala vestibuli from the cochlear duct

Re'nal clea'rance, See Plasma clearance

Retic'ulo-endothe'lial sys'tem [L reticulum, a network + endothelial, relating to the endothelial tissues] A system of cells in or associated with various reticular and endothelial structures, and also present in the general connective tissues, they have diverse functions There are several types of these cells, some of those in the connective tissues are wandering, others are fixed. Those in the reticulum of the spleen are very large and phagocytic (macrophages), and those in the blood sinuses of the liver (Kupfler cells) as well as some in the general connective tissues convert hemoglobin into bile pigment Others are found in the reticulum of the bone marrow. The monocytes of the blood and certain small cells in neuroglia (microglia) belong to this system

Retic'ulum IL, diminutive of rete, a netl. A fine network

Re'tro- [L. back, backwards] A prefix meaning backwards, behind, posterior to Rhe'obase [G rheos, a stream (current) + basis, base] The minimal strength of an electric current that, acting for an indefinite but fairly long time, causes excitation of a tissue e g, nerve or muscle

Rhom'hold [G rhombos + eidos, a resemblance] Resembling a rhombus, which is a figure (exclusive of the square) with four equal and parallel sides

Santori'ni's duct An accessory pancreatic duct

Sciatic forumen (sì at'ik for-a'men) [corruption of G ischiadikos fr ischion the hip! One of the two foramina, greater and lesser, formed by a ligament closing the gap of the corresponding sciatic notch of the os coxae

Sciat'ic notch [prec ] One of the two notches greater and lesser, of the os coxae The greater sciatic notch is the large indentation on the posterior border of the ilium below the posterior inferior spine, the lesser sciatic notch is situated on the posterior border of the ischium just above the tuberosity. Both notches

are converted into foramina in the living subject by ligaments Sciatica (si at ika) Inflammation or pain in the sciatic nerve

Sciero'sis [G sklerosis hardness] Induration, hardening of a structure, such as the walls of arteries (arteriosclerosis) due to the deposition of calcium salts or of any tissue, such as nervous or hepatic, by the replacement of its parenchyma by fibrous tissue

Semi- [L semis, half] Prefix meaning half, equivalent to Greek hemi

Se'rous [foli ] (1) Relating to serum (2) Qualifying certain membranes (e membranes) that are covered by a layer of mesothelium-e g, peritoneum, pieura, etc

Se'rum [L whey] Blood fluid from which the fibrinogen has separated our through clotting of plasma

Sig'mold [G sigma letter S + eidos, a resemblance] S shaped, e.g., the sigmoid colon (that and pelvic colons) which together form an S shaped curve

Spay'ing [G spadon a cunuch] Ovariectomy, castration of a female Species (spe'shez) [L kind, form, appearance] In the classification of animals

and plants, the division pext below the Genus, its members interbreed Specific gravity (abbreviated Sp g) The weight of any given substance as compared with an equal weight of distilled water Thus, water as having value 1 000, the specific gravity of urine is 1 002-1 030 (depending upon the concentration), blood, 1 055-1 060; plasma, 1 022-1.26, and cerebrospinal fluid,

1 006-1 08 Sphyg'mogram [G sphygmos, pulse + gramma, a record] A pulse tracing

- Sphyg'momanom'eter [G. sphygmos, pulse + manos, pressed (pressure) + metron, measure] An instrument for measuring the blood pressure
- Spi'na bif'ida [L spina, a spine, thorn, the back + bifida, cleft into two parts]
  A congenital anomaly in which the arches of the vertebrae of a portion of the spinal column are absent or defective so that the spinal membranes protrude, or even the cord itself, and come to lie just beneath the soft tissues of the back. It occurs most commonly in the lumbar region
- Sta'ble, sta'bile [L stabilis, steady] Firm, steady, not susceptible to change, applied to a chemical compound or to temperament, behavior, mentality, etc Stei'late [L stella, a star] Star-shaped
- Steno'sis [G stenos, narrow] A narrowing, usually applied to an orifice or tube, e.g., aortic stenosis

Sten'son's duct. The duct of the parotid gland

- Stokes-Adams disease. Heart block, causing a slow pulse, with arteriosclerosis, the combination of the two conditions making for a low disstolic pressure and consequent interference with the blood supply to the brain Attacks of syncope and convulsions result
- syncope and convenient seems states, a mouth] Any mouth like opening Stomati'tis [G stoma (-aus), a mouth + us inflammation] Inflammation of

the mouth Stri'a [L a furrow or line] A fine line, linear groove, or elevation

Stri'ation [prec] The appearance given by numerous fine lines

- Stro'ma—stromata [G a bed, mattress] The framework of any tissue or organ, it is usually composed of connective tissue
- Sty, stye (sti) [fr OE stigan, to rise] Inflammation of one of the small glands at the margin of an eyelid, hordeolum

Sub- [L below] A prefix indicating below, less than

- Sul'cus [L a furrow, a ditch] A narrow, relatively deep, depression in any structure, but especially in the cerebral cortex
- Su'per-, su'pra- [L above, beyond] A prefix indicating above, beyond, greater than, on the upper side
- Suppura'tion [L'suppura (-atus), to form pus (pur)] Pus formation Su'ture [L sutura, a seam] (1) A surgical stitching (2) One of the joints be-
- tween the bones of the skull Sym'patholyt'ic [sympatho, fr G syn, with + pathos, feeling + lysis (lyt-),
- freeing from, dissolution] Referring to the action of certain agents in temporarily paralyzing sympathetic nerve fibers
- Sym'pathonimet'ic [sympatho (prec) + G numikos, unitating] Referring to any drug or other agent, e.g., adrenaline that imitates the action of sympathetic nerve fibers
- Syn-, sym- [G together with] A prefix meaning with, together, joined, equivalent to Latin con-
- lent to Latin con-Synarthro'sis [G. syn, together + arthrosis, articulation] An immovable junction between two bones without an intervening joint
- Syncy'tium [G syn, together, with + kytor, a hollow (cell)] A mass of multi-nucleated protoplasm, such as the cardiac muscle, in which the cells are not distinct and separate units, but are continuous with one another without an intervening cell membrane
- Syndesmo'sis [G syndesmos, ligament, fr syn, together + deō, bind] A synarthrosis in which the opposed bones are held together by fibrous tissue, as at the lower ends of the tibia and fibula
- Syndrome (sin'dröm or sin'drō-mē) [G syn, together + dromos, a running]
  A group of symptoms which (running) together form the picture of a disease

- Syn thesis [G syn together + thesis a putting an arrangement] A building up of smaller parts into a larger whole the putting together of smaller chemical groups to form larger and more complex compounds
- To bes dorsa'lls [L. taber a wasting away + dorsails of the back] A nervous disease affecting the ganglia of the posterior spinal nerve roots and posterior columns of the spinal cord. It is characterized by ataxia loss of the knee jerks severe shooting pains and often atrophy of the optic nerve. It is usu ally the result of syothis.

Tach year dia (k) [G tachys rapid + kardia heart] Rapid action of the heart Tach yphylax'is (k) [G tachy; rapid + phylaxis protection] Loss of or reduction in sensitivity to any agent tolerance following repeated use

Tax'in, tax'is [G order arrangement] [1] Suffix in words of Greek origin.

(2) The word is also used alone and then means to correct a dislocation or to return a herma by gentle pressure to its normal position (3) The tendency for certain organisms to move toward or away from a particular thing or state or to arrange themselves in a position relating to that particular thing or state or to arrange themselves in a position relating to that particular thing or state or to arrange themselves in a position relating to that particular thing or state or to arrange themselves in a position relating to that particular thing or state or to arrange themselves in a position relating to that particular thing or state or to arrange themselves.

motaxis etc

Te lophase [G telos end fulfillment + pt asis appearance] The final stage of
milosis

Ther mola bile [G therme heat + L labilis perishable] Susceptible to change or destruction by heat

Ther mosta ble, ther mosta bile [G therme heat + L stabilis stable] Not altered by moderate heat

Thrombo'sis [G a curding] Clott ng of blood within the vascular system Throm bus [G thrombos a clot curd] A clot within a blood vessel

Thymus [G thymos an excrescence sweetbread] A soft structure occupying the upper part of the mediastimum and lower extremity of the neck. It is relatively large in infants but shrinks in later childhood and is a mere vestige in adults. It consists of a reticulum the spaces of which are packed with lymphocytes. Its outer part is called the cortex, its central part the modults. Scattered throughout the latter are peculiar bodies known as Hassell's corpuscles and composed of concentric layers of flat ep thel al cells surround ng a large round granular cell. Though it has been suggested that the thymus is an endocrine organ there is no definite evidence that it should be classed as such its function is obseure.

Thy rotoxico sis [thyro (thyroid) + G toxicon po son + our state or condition] The condition caused by an excess of thyroid hormone—e g exoph

thalmic gotter

Ti dal air The air breathed during ordinary respiration it amounts to about
500 cc in man

Tone [O ronos tone or a tone] (1) Continuous action as sa d of a nerve (such as the vagus) or of the slight continuous contraction of healthy muscle (2) The pitch of a sound (3) A qualt y of color depth brill ance intensity Trans- IL across through beyond! A prefix meaning across through beyond Traum a - Traum at [G a wound] A wound or injury usually one caused by

violence or some mechanical agency

Tri be [L. tribus] In the classification of animals and plants a subdiv s on of the

Family frequently synonymous with Subfamily
Tri ceps [1. 17] three + caput a head] Applied to any structure but especially
a muscle that hat three heads or origins

Troch lea (k) [L a pulley] A pulley

Troch'old joint (k) [G trochos wheel + eidos, a resemblance] A pivot joint, such as that between the odontoid process of the axis vertebra and the atlas Tro'pism [G tropē, a turning] The tendency for certain organisms to turn toward something as growing sprouts of plants turn toward the light, photo tropism and its roots into the ground, geotropism.

Tu'bercle [L. fr tuberculum diminutive of tuber, a swelling] (1) a small circumscribed elevation on a bone (2) The lesion in tuberculosis

Tuberos'ity [L tuberositas an elevation, or swelling] An elevation on a bone larger than a tubercle

Tympam'tes [fr G tympanon a drum] Swelling of the abdomen due to gas in the intestine or, rarely, in the peritoneal cavity

Ul'tra [L beyond] A prefix meaning beyond excessive

Ul'trafiltra'tion [prec + filtration] The separation of salts (crytalloids) from colloids by filtration

U'vea [L. uvea a grape] The middle coat of the eyeball, consisting of the choroid ciliary muscle and iris

U'veal tract Uvea

Va'gusstot'fe [vagus + Ger stoffe substance] The humor formed at the terminals of the vagus nerve acetylcholine

Vallate [L. vallo (are) surround with fr vallum a rampart] Referring to certain structures e.g., papillae of the tongue, that are surrounded by a circular mostlike depression bounded in turn by a slightly elevated rim

moatiske depression bounded in turn by a slightly elevated rim Vallec'ula [L. diminutive of vallus a valley] The depression on either side of the tongue or of the vermis of the cerebellum

Varicel'la [L diminutive of variola] Chickenpox

Vario'la [L diminutive of various spotted] Smallpox

Vas [L vasa (genitive, plural vasorum), a vessel dish] A vessel or tube, e g vas deferens

Va'sa vasor'um [prec ] The small blood vessels supplying the walls of the larger arteries

Vas'cular [L vasculum diminutive of vas a vessel] Referring to any tube or system of tubes containing fluid but most commonly to the blood vessels or lymphatics

Va'somo'tor. Referring to changes in the calibers of the blood vessels

Ven'trad [L venter, belly] Toward the ventral aspect, opposed to dorsad

Ven'tral [prec] Indicating the front aspect of the human body or the under

side of a quadruped, opposed to dorsal

Vermiform [L vermus a worm + forma form] Wormlike, e.g., vermiform appendix

Ver'nix caseo'sa [L vernix varnish + caseus cheese] The fatty substance on the skin of the newborn

Vestib'ular mem'brane. Reissner's membrane

Vestib'ular or'gan The utricle and saccule

Vi'ral Relating to viruses

VI'rus—strusse [L. poison] One of a group of infecting agents of uncertain nature but probably a large and complex nucleoprotein molecule near the border line between the living and the nonliving Unlike bacteria they can multiply only within cells. They are much smaller than bacteria they can multiply only within cells. They are much smaller than bacteria to poss through a filter with much finer pores than those required to hold back bacteria hence the term filtrable viruses. Though viruses cannot be seen through an ordinary type of microscope, however powerful, the electron microscope is not proposed to the property of the proposed of the proposed of the property of the proposed of the proposed of the property of the proposed of the property of the proposed of the property of the property of the property of the proposed of the property of the pro

## 704 GLOSSARY

croscope reveals minute objects called elementary bodies that are probably the viruses themselves

Viscosity (L. viscosius viscid viscous). Stickiness internal friction of a liquid or semiliquid as measured by the rate at which the material flows along a narrow

tube The value is expressed in relation to the viscosity of water taken as 1 Viscous [prec ] Sticky glutmous

Vis'cus-viscera [L. an internal part] Any internal organ but especially of the

thorax abdomen or pelvis Vo lar [L vola palm of the hand or sole of the foot] Relating to the palm of the hand or the sole of the foot or to any similarly directed part-e g the front of the arm or the back of the leg It is synonymous with ventral and palmar in the former instance and with dorsal and plantar in the latter

Whar'ton's duct The duct of the submaxillary gland


Whit'low fetymology dub ous perhaps corruption fr OE quick (whick) or white + flawl Inflammation of the soft tissue in the neighborhood of a nail paronychia

Wirsung's duct The pancreatic duct

Wis'dom tooth One of the four molar teeth that crupt very late-from the eight centh to the twenty third year it is the third molar on each side of the upper or the lower saw

Zoology [G zoon an animal + logor treatise discussion] The study of ani mals their classification structure and phys ology etc Zy'mogen [G zyme leaven (en-yme) + root gen produce] The substance

from which an enzyme is formed proenzyme



# Appendix

## Weights and Measures

#### THE METRIC SYSTEM

#### Linear measures

The unit of the metric system is the meter which is the one ten milionth part of the meridian quadrant of the earth. In the nomenclature of the system, multiples of the meter are indicated by prefixes derived from the Greek, as follows.

Meter decameter, 10 meters hectometer, 100 meters, kilometer, 1000 meters

Fractions of the meter are indicated by prefixes derived from the Latin, as follows

Meter, decimeter, 1/10 meter centimeter, 1/100 meter, millimeter, 1/100 meter

In microscopy, the unit of measure is  $\cancel{1}_{1000}$  of a millimeter, called micron (symbol  $\mu)$ 

#### Weights

The unit of weight is the gram or gramme, abbreviation, usually g or gm It is practically the weight of one cubic centimeter of distilled water at its maximum density (4°C) Multiples of this unit are designated by prefixes derived from the Greek numerals as follows

Gram, decagram, 10 grams, hectogram, 100 grams kilogram (abbr kilo), 1000 grams

Fractions of the gram are designated by prefixes, derived from the Latin numerals as follows

#### Fluid measure, volume, capacity

The unit of volume is the cubic decimeter, called a liter one liter of water weighs practically 1 kilogram. It is divided into the deciliter,  $\frac{1}{10}$  liter

Kilograms	Ounces (Avoirdupois)	Kilograms	Pounds (Avoirdupois)
1	35 27	1	2 2048
5	176 37	5	11 0240
10	852 74	10	22 0480

#### Fluid Measures

One liter = 1 76 imperial pints or 2 1 U S pints, I cubic centimeter = 17 minims (British) or 16½ minims (U S)

One imperial gallon = 4.55 liters 1 U S gallon = 3.79 liters, 1 imperial pint = 568 cubic centimeters, 1 U S pint = 473 cubic centimeters, 1 fluid ounce (U S) = 29.5 cubic centimeters 1 fluid ounce (U S) = 29.5 cubic centimeters 1 fluid drachm (U S) = 3.7 cubic centimeters 1 fluid drachm (U S) = 3.7 cubic centimeters, 1 minim = 0.065 cubic centimeter

To convert liters to imperial gallons, multiply by 22 and divide by 100 to convert liters to U S gallons multiply by 265 and divide by 1000 (by moving the decimal point three places to the left), to convert imperial gallons to liters divide by 22 and multiply by 100 to convert U S gallons to liters divide by 365 and multiply by 1000 to convert U S gallons to liters divide by 365 and multiply by 1000

#### Metric equivalents of U S measures

Minims	Cubic	Fluid	Cubic	Fluid	Cubie
	Centimeters	Drachms	Centimeters	Ounces	Centimeters
1	0 061618	1	3 697086	1	29 5"6686
5	D 308091	5	18 485431	6	177 460116
10 15	0 616181 0 924272			16 (pt)	473 226976

## Metric equivalents of British (Imperial) measures

Minims	Cubic Centimeters	Fluid Drachins	Cubic Centimeters	Fluid O inces	Cubic Centimeters
1	0 059905	1	3 5523	1	28 4184
5	0 296025	5	17 7615	5	142 0920
10	0 590050			10	284 1840
15	0 888075			20 (pt.)	568 5680
90	1 184100			. 4 . 7	

#### Equivalents of metric in U S measures

Cubic Centimeters	Minims	Later	Fluid ounces	Pints
0.1	1 62341	1	33 82108	2 11381
0.5	8 11706			
10	16 23412			
5 0	81 17061			
10 0	162 34122			

### Equivalents of metric in Imperial measures

Cubic Centimeters	Minims	Liters	Fluid ounces	Pints
0 1	1 68911	1	35 19691	1 75981
0.5	8 44556			
10	16 89112			
50	84 45560			
10 0	169 91123			

#### MEASURES OF ENERGY

One kilogrammeter is the energy or force expended in raising a weight of 1 kilogram to a height of 1 meter. One foot pound is the energy or force expended in raising a weight of 1 pound avoirdupois to a height of 1 foot. One kilogrammeter = 724 foot pounds. One foot pound = 0 1931 kilogrammeter.

	SYM	BOLS	
収分 5 5 0 μ μ 抽 十 一 土	minim scruple drachm ounce pint micron micron micron micron micron misce mix plus excess acid reaction positive minus deficiency alkaline reaction negative plus or minus either positive or neg ative, indefinite	> V8 * † \$ ° ° 6	greater than whence from which is derived less than from derived from infinity ratio is to equality between ratios as lurth death female, male
	Att of Withamson		

#### COMPARATIVE TEMPERATURE SCALES

Centigrade	Fahrenheit.
	220
100	🗏
· · ·	210
	200-
90	
	100-
	180-
80	
	170
I ∄	160
70日	100
	150-
Ⅰ目	<del>.</del> 🗒
60 🗐	140-
l ≠≡	180-
I 50 🗐	-
I <b>"</b> "⊟	120
	110
40=	
\	100
	90-
80=	-
	80-
目	70-
20	
I■	60-
10-	50
し""員	
	40-
- 6■	20-
	***
	20
10=	10-
20	0-
"冒	10
一目	10
	F

The zero of the centigrade scale marks the tem perature of melting ice (32°F), the zero of the Fahr enheit scale is an arbitrary point, that of the lowest temperature observed by the deviser of the scale during the winter of 1700 practically the temperature of a mixture of ice and salt, it corresponds to ~177°C. The temperature of boiling water, at sea level, is marked 100° on the centigrade scale and 212° on the Fahrenheit scale A degree F is therefore \(^2\)6 degree C

The following are rules for the conversion of the temperature of one scale into that of one of the others.

Above O°C or \$2°F

F to C subtract 32 multiply by 5 divide by 9 C to F multiply by 9, divide by 5 add 32

Between 0° and 32°F or -17 77, and 0°C

F to C subtract from 32 multiply by 5, divide by 9 C to F multiply by 9, divide by 5, subtract from 32

## A CLASSIFICATION OF THE PROTEINS

CIAM OF PROTEIN	CHARACTERISTICS	EXAMPLES
	A. Somple proteins	
(t) Albumins	Soluble in water and congulable by heat. Present in both animal and plant tissues	Soram albumin egg albumin lactalbumin and various vegetable albumins such a leucoins (in wheat rye and barley) legu milin in lentilla, soy bean beans and peas and phased in m kidney bean
(I) Gobalias	Soluble in dilute saline solutions insoluble in water Animal globulins are coagu- lated by heat. Vegetable globulins in- perfectly or not coagulated by heat	Serum globul n firmsogre (and forum) suid ins of egg yolk and vegetable globulum such as execton (Brazil-out) ederin (bermp) phasesiss (Biology bean) lega- min (pean and lentils) and tuberin (po- tato) A number of other vegetable globulum shaw been besideted and named
(3) Gluidins	Found only in plants. Insoluble in water saline or alcohol but soluble in very dilute alkali	Glatenia of wheat, organia of nice and glatelia of maite
(4) Prolomines or Gladins	Found in cereals (except rice) soluble in 70-90 per cent alcohol. Insoluble in water They contain a large proportion of proline and ammonia nutrogen	Glissis of wheat, kordens of barley and keen of maize
(5) Albuminoids or sclar-proteins	Especially renstant to the usual reagents. They enter into the construction of protective and connective tissues, e.g. thin tendons, ligaments and hones	Kerain of hair skin bone feathers, tor touse shell and egg shell, elastin colla- gen extens and gelains of tendons, hig aments, bone etc.
(6) Histones	Soluble in water and precipitated by ammonia solution and by alkaloids. They contain a large percentage of diamino acids (p. 625)	Globin of hemoglobin thymus histone scombron and godur histone in sperma- tozon of mackerel and cod fish respec- tively
(7) Protomines	Found in combination with nucleic acid in heads of fish spermatozos. Con- structed predominantly of diamino- acids	Salmins and sturgeon respectively
B Conjugates	polerns Proteins whose molecule is combine	d with another non protein group
(1) Nucleoprotesus	Nucleic acid in combination with a protein belonging usually to the class of histones	See ch. 48
(2) Chromoproteins	or protamines. Found in cell nucles Protein in combination with a pigment (e. g., bemain) containing from copper or other metal	
(3) Glycoproleins	Proteins other than nucleoproteins in combination with a carbohydrate group	
(4) Lipoprotesus (5) Phosphoprotesus	Proteins in combination with lipid Proteins other than nucleoproteins and lenthoproteins in combination with a phosphorus-containing group	Present in plasma, milk cell nuclei Carenogen (and casem) mielles of egg yolk
C. Deresatives of prote	enzymes upon certain of the proteins	the action of acids, alkalis or proteolytic hsted above
(a) Primary derivatives (1) Proleans	Insoluble products formed in the early stage of the action upon proteins of water dilute acids and enzymes	

## CLASSIFICATION OF THE PROTEINS (CONT.)

	C. Derrestrees of proteins—derivated proteins.—Continued							
(2) Metaproteus	Formed in a later stage of the action of acid or alkali	And metaprotes alkali metaprotes						
(3) Cospulated pro- teins	Formed by the action of heat or of akohol upon solutions of proteins	į						
<ul><li>(b) Secondary derivatives</li></ul>		}						
(1) Protesses	Formed by the action of pepsin or trypus upon proteins. They are soluble in water from which they are preoputated by satu ation with ammon un sulphate. They are incongulable by heat	Albument from albumen globulese from globulin, excesse from cusein						
(2) Poplones	These represent a further stage in action of protecylur enzymes. They are soluble in water but are not precipitated from an acurous solution by ammonium sulphate. They are not congulated by heat	}						
(3) Peptides di-pep- tides Impeptides	Products formed in the final stages of proteolytic digestion	Glycyl-alanine lencyl-glutanic add, etc.						

# A LIST OF THE AMINO ACIDS PRESENT IN MOST COMPLETE PROTEINS

I. ALIPSATIC ANDVO-ACTOR

A. Mondomino-monocorborrhic acids

Glycine (or glycocoli) C<sub>1</sub>H<sub>2</sub>NO<sub>2</sub> or amino-acrtic acid

COOR

Alanine C.H.:NO: or e-atnine-propionic acid CH: CH-NH:

COOM

Serion C.H.NO. or e-amino-8-hydroxy proponic

OH COOH

Threonine C.H.NO.

CH,CH CH—NH,

Valine C.H., NO.

Notleucine C.H ,NO2

CH, CH, CH, CH, CH—NH,

Leucine, Call aNO:

Isoleucine CoHuNOs,

Sulphur-containing monoamine-monocarbest acids

Cystine Call sN.S.O.

# AMINO ACIDS PRESENT IN MOST COMPLETE PROTEINS (CONT.)

Methionine, C.HnSNOs,

CH1-S-CH1-CH1-CH-NH1

COOR

R. Monoamino-dicarhovelic acids

Aspartic acid. C.H.NO..

COOH-CH:-CH-NH:

COOR

Glutamic acid, C<sub>4</sub>H<sub>2</sub>NO<sub>4</sub>, or α-amino-glutaric acid

COOH-CH,CH,CH-NH,

COOH

Hydroxyglutamic acid, C,H,NO,

COOH CH,CHOH-CH—NH,

C. Diamino-monocarboxylic acids

Arginine, C<sub>4</sub>H<sub>14</sub>N<sub>4</sub>O<sub>24</sub>

IN⇒C—NH·CH₁·CH₁·CH₁·CH—NH₁

:00

Lysine, Calla N.Oz.

NH.

CH1-CH1-CH1-CH1-CH-NH1

ייטים

II. AROMATIC AMINO-ACIDS

Phenylalanine, C.H., NO.

Tyrosine, C.H., NO.

III. HETEROCYCLIC AMING ACIDS
Tryptophane, CuHuNrOn

Histidine, C.H.N.O.,

Proline, C.H.NO.

Pyrrole nucleus

Hydroxyproline (oxyproline), C,H,NO,

# ASH CONTENT OF THE EDIBLE PORTION OF SOME COMMON FOODS

(Modified from Lusk)

	1		tM 100	CSTR	S TREE	a zas	STANCE	L
	Inn		Calcium	Magnesium	Sodium	Potessium	Phosphorus	Chlorus
	-	r	×t.	=:		m:	mr.	m.
Beefsteak, lean	3	8	8	24	6	35	22	50
Liver (beef)	8	0	11 (	o l	1	1	i	1
Eggs	3	0		9				
Milk, whole	0	2	120	11	51	142	94	120
Commeal	1	1		İ	1	1	1 1	
Oatmeal	3	7	93	127				35
Rice, polished	0	7	8	27				50
Wheat flour	1	5	26	30	69	146	86	76
Wheat, entire		I			Į	l		
grain		2	44	170		515		88
Beans, lima, dried	7	2	71	187	245	1743	336	25
Beans, string,		1			1 :			
fresh		6			i	i	- 1	
Cabbage		옛	49	14	20	243	27	13
Corn, sweet		8				1	- 1	
Peas, dried.			100	145			397	40
Potatoes	1	2	11	22	19	410	61	30
Spinach		8	1	l			- !	
Turnips		6	64	169	59		51	40
Apples		3	10	8	15	125	13	4
Raisins	3	6	57	9	141	830	126	70

# AVERAGE COMPOSITION AND ENERGY VALUES OF EDIBLE PORTIONS OF SOME COMMON FOOD MATERIALS

	1		PLS CENT			ENTRO	ANTOE
_	Water	Protein N X 6.25	Fat	Carbo. Sydra e	Ash	Per kg	Per
	Frame	- Front	trems.	grams.	frem!	celories	caleries
Mest.	í	1	ł .	(	1	ĺ	l
Beef, round steak, medium fat	54.8	23 5	20 4	1	12	2 860	1 300
Mutton, leg roast	509	250	22 6		12	3 125	1 420
Pork, ham, luncheon bacon, side	1 49 2	22 5	21 0	·	58	2 870	1 305
Fork, mam, mincheon bacon, ande	18 8	99	67 4	1	4.4	6 665	3 030
Chicken		1			İ	!	
Broilers	74 8	21 5	2.5	1 1	11	1 110	505
Fish, cod, whole	82.6	16.5	04	)	12	715	325
Herring, whole	72.5	19 5	71		1.5	1 455	660
Salmon, whole	64.6	22 0	128	1	14	2 090	950
Trout, brook whole	77 8	19 2	21		12	980	445
I four, proce whose	" "	19 2	21		1 2	700	443
Fats		1				1	
Butter	110	10	85 D	,	30	7 930	3 605
Lard	1	I	100 0			9 285	4 220
Suet	13 7	47	818		0 3	7 790	3 540
Cheese				i '		[ ]	
American, red	28 6	296	38 3		3 5	4 765	2 165
Milk	87.0	33	40	50	07	715	325
Eggs, hens', boiled	73 2	13 2	12 0		0.8	1 685	765
Flour white, wheat	11 5	11 4	10	75 6	0.5	3 650	1 660
Bread, white	35 6	93	12	52 7	12	2 650	1 205
Fruit	- 1	1 1		} }	'	1	
Apples	84 6	04	0.5	14 2	0.3	640	290
Banana	75 3	13	0.6	220	0.8	1 010	450
Chernes	80 9	10	0.8	16 7	0.6	805 Ì	365
Grape fruit	936	06	01	57		267	120
Oranges	86 9	0.8	02	16	0 5	528	240
Vegetables	- 1	1 1		1			
Beans, dried	12 6	22 5	18	596	3 5	3 530	1 605
Cabbage	91.5	16	03	56	10	320	145
Lettuce	94 7	12	0.3	29	0.9	206	90
Potatoes	75 5	25	01	209	10	968	440
Sugar granulated	1			100 0		4 090	1 860
Chocolate	59	12 9	48 7	303	22	6 295	2 860
Cocce, powder	4.6	21 6	28 9	377	72	5 105	2 320

#### RECOMMENDED DIETARY ALLOWANCES\*

(Food and Nutrition Board National Research Council)

	CALCULES	PROTEIN	CHICLE	n	ertage Al	(A)	BURO- PLAYER	STACTO TORRE ACUS)	1400 845 1400 1400 1400 1400 1400 1400 1400 14	ALLEYSTE D
	[	-			10	ert	-	=1	-41	14
Man (70 Kg)	1	i	1	1	!	ł			1	1
Sedentary	2500	1 -	I I	۱	1 -	1 2 5	2 2	15	I _	١ -
Moderately active	3000	70	08	12	5000	18	27	18	75	
Very active	4500	-	- 1	_	-	2 3	33	23	-	-
Woman (56 Kg)							1 1		١.	
Sedentary	2100	1 1	1 1	_	l	1 1 2	118	12	l _	1
Moderately active	2500	60	108	12	5000	1 1 5	22	15	70	
Very active	3000	-	-	_	-	18	27	18	-	
Pregnancy flatter			1 1							
half)	2500	85	1 2 5	15	6000	18	25	18	100	400 to 800
Lectation	3000	100	20	15	\$000	23	30	23	150	400 to 800
Children up to 12 years			ĺĺ					- 1		
Under 1 years	100/kg	3 to 4/kg	10	6	1500	04	06	4 1	30	400 to 800
1-3 years I	1200	40	10	7	2000	061	09	6 1	35	
4-6 years	1600	50	10	8	2500	08	12	8	50	-
7-9 years	2000	60	10	30 İ	3500	10	251	10	60	
10-12 years	2500	70	12	12	4500	12	18	12	75	-
Child en ove 12 years	- 1		1	- 1	- 1	- 1	- 1	- !	- (	
Cirls, 13-15 years	2800	80	13	13	5000	14	20	14	80	••
16-20 years	2400	75	10	12	5000	1 2	1.8	12	\$0	-
Boys, 13-15 years	3200	85	14	15	5000	16	24	15	99	••
16-20 years	3800	100	14	45	6000	20	30	20	100	_

<sup>\*</sup> Tentative goal toward which to sim in planning practical detaries can be met by a good dot of natural food Such a duct will also provide other mnerals and vilamens, the requirements for which are less well known. § 1 mg thanine equals 333 IU I mg ascorbic and equals 20 LU

Requirements may be less if provided as vitamin A greater if provided chirdly as the pro-vitamin excetene.

Needs of infants increase from month to month. The amounts given are for approximately 6-8 months.

The amounts of protein and calcium needed are less if derived from human milk.

¶ Allowances are based on needs for the middle year in each group (as 2 5 8, etc.) and for moderate activity

"V tamm D is undoubtedly necessary for older children and adults. When not available from sombline it
should be p ovided probably up to the minimum amounts recommended for individent probably.

Further Recommendations Adapted 1947.

The requirement for sed as is small probably about 0 002 to 0,004 milligram a day for each kilogram of body.

weight. This amounts to about 0.15 to 0.30 milligram duly for the adult. This need is easily met by the

regular use of foduced salt; its use as especially important in adolescence and pregnancy. The requarement for copper for adults is in the neighborhood of 10 to 20 milligrams a day. Infants and children require approximately 0.05 per hilogram of bodyweight. The requirement for copper is approximately one-term of that for the million of the control of the foreign of the control of the control of the foreign of the control of the foreign of the control

The requi ement of vikewis K is usually satisfied by any good diet. Special consideration needs to be given to newborn infants. Physicians commonly give vitamin K either to the mother before delivery or to the

infant immediately after birth.

## STANDARD WEIGHT ACCORDING TO HEIGHT AND AGE

HEIGHT AND AGE																		
Men																		
Αp	11	6 Ft. 1 To.	6 Ft. 2 Ia.	8 Pt. 3 Ia.	17.614	1 Pt. 6 It.	# Ft. # Ja.	8 PL 7 ft.	41.4	6 Pt. 9 In.	1 5 Ft. 10 In.	6 P. 15 Ia.	4	6Ft. 1 Ea	6P. 3 Th	4 P. 1 1 1a	57.4 In	471.6
15 16 17 16 19	10° 109 111 113 115	109 111 113 115 117	112 114 116 118 120	115 117 119 121 123	118 120 122 124 126	122 124 126 128 130	126 128 130 132 134	130 132 134 136 138	134 136 138 140 142	138 140 142 144 146	142	147	152 154 156 158 160	1 150	164 166	16: 16: 17: 17: 17:	17	179
	•	1	3	<u> </u>	13	3	•	Ī	Œ	Ī	10	11	1.	ĮΞ	3	3	1	<u> </u>
20 21 22 23 24	117 118 119 120 121	110	123 124 124 125 126	125 126 127 128 129	128 130 131 132 133	132 134 135 136 137	136 138 139 140 141	140 141 142 143 144	344 145 146 147 148	148 149 150 151 152	152 153 154 155 156	156 157 158 159 160	161 163 164 165	166 167 169 169 171	172	371 177 178 186 183	183 183 184 2 183	183
25 25 27 28 29	122 123 124 125 126	124 125 126 127 128	126 127 128 129 130	129 130 131 132 133	133 134 134 135 135	137 138 138 139 140	141 142 142 143 144	145 146 146 147 148	149 150 150 151 152	153 154 154 155 156	157 158 158 159 160	162 163 163 164 165	167 168 169 170 171	173 174 175 176 177	179 180 181 182 183	184 186 187 188 189	192	194 196 197 198 199
	-	1	1	7	4	•	•	7	•	•	10	11	•	1	3	7	1	-
30 31 32 33 34	126 127 127 127 127 128	128 129 129 129 129 130	130 131 131 131 132	133 134 134 134 135	136 137 137 137 138	140 141 141 141 142	144 145 145 145 146	148 149 149 149 150	152 153 154 154 155 155	156 157 158 159 160	161 162 163 184 165	166 167 168 169 170	172 173 174 175 176	178 179 180 181 182	184 185 186 187 188	190 191 192 193 194	196 197 198 199 200	201 202 203 204 206
i	7	7	1	-	1	•	•	f	•	•	ΩĐ	n	7	1	7	3	7	-
85 86 87 88 89	128 129 129 130 130	130 131 131 132 132	132 133 133 134 134	135 136 136 136 137 137	128 139 140 140 140	142 143 144 144 144	146 147 148 145 148	150 151 152 152 152	155 156 157 157 157	160 161 162 162 162	165 166 167 167	170 171 172 173 173	176 177 178 179 179	182 183 184 185 185	189 190 191 192 192	195 196 197 198 199	201 202 203 204 205	207 208 209 210 211
	-	-	그	1	<u>-</u>	-	-	-	1	-	10	-11	-	-	-	-	-	8
43 43 41 41	131 131 132 132 133	133 133 134 134 135	135 135 136 136 137	138 138 139 139 140	141 141 142 142 143	145 145 146 146 147	149 149 150 150 151	153 154 154 154 155	158 158 159 159 160	163 163 164 164 165	168 168 169 169 170	174 174 175 175 176	180 180 181 181 182	186 186 187 187 188	193 193 194 194 195	200 200 201 201 202 202	206 207 208 208 209	212 213 214 214 215
	3	-	,	3	7	3	•	7	3	<u> </u>	10	11	•	工	3	3	₹,	7
45 65 47 48 49	133 134 134 134 134 134	135 136 136 136 136 136	137 138 138 138 138	140 141 141 141 141	143 144 144 144 144	147 148 148 148 148	151 152 152 152 152 152	155 156 156 156 156 156	160 161 161 161 161	185 186 166 166 166	170 171 171 171 171 171	176 177 177 177 177 177	182 183 183 183 183	188 189 190 190 190	195 196 197 197 197	202 203 204 204 204 204	209 210 211 211 211 211	215 216 217 217 217 217
		1	3	3	<u>.</u>	•	ات	ᄀ	5	•	10	11	<u>-</u>	1	1	-	4	<u> </u>
50 51 52 53 54	134 135 135 135 135	136 137 137 137	138 139 139 139 139	141 142 142 142 142	144 145 145 145 145	148 149 149 149 149	152 153 153 153 153 153	156 157 157 157 157 158	161 162 162 162 163	166 167 167 167 168	171 172 172 172 173	177 178 178 178 178	183 184 184 184 184	190 191 191 191 191	197 198 198 198 198	201 205 205 205 205 205 205	211 212 212 212 212 212	217 218 218 218 218 219

# STANDARD WEIGHT ACCORDING TO HEIGHT AND AGE

TT7 .

Women																		
λr	48.49	(Pt. 9 la.	47.1014	47t.11 Ia	Ę	414	2									10.1		:
15 16 17 18 19	10 10 10	2 10 3 10 4 10	4 100 5 100	10	10	9 11	1 1 3 1 4 1	6 1	15 1 17 1 19 1 20 1 21 1	18 1 20 1 22 1 23 1 24 1	24 1 25 1 26 1	23 1 29 1 30 1	32 1 33 1 34 1	36 1 37 <sub>1</sub> 1 38 1	39 1 40 1 41 1	43 1 44 1 45 1	49 1	83
	1 8		10	11		1-	3	-	-1-				-	-	-1-	-1-	<u> </u>	-
20 21 22 23 24	10 10 10 10 10	10 10 11	9 111	113		57 11.	7 12 7 12 8 12	0 1	241 1	25 1 26 1 26 1 27 1 27 1	29 I 29 I 30 I	32 1 33 1 33 1 34 1 34 1	7 1	11 1 11 1 12 1	64 1 65 1 66 1	48 1 49 1	51 1 52 1 53 1 53 1 54 1	56 57
	1		10	11	1.	1	. 2	Ŀ	- 6	_		-1-	1.	-1-	_ [	~	_!_	_
25 26 27 26 29	100 110 110 111 111	113	114	115 116 116 117 117	117 116 118 119 119	120 120 121	) 12 12 12	1 12 2 12 2 12 3 12 3 12	5 13 5 13 6 13 6 13	8 13 9 13 0 13	1 13 2 13	5 13 6 14 7 14	9 14 0 14 1 14	3 14 4 14 5 16	7 1. 3 1. 9 1.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	55 15 55 15 56 16	999
		1	10	11	-	1	3	3	1 .		10	7			20	11	•	_
30 31 32 33 34	112 113 113 114 115	114 115 115 116 117	116 117 117 118 119	118 119 119 120 121	120 121 121 122 123	122 123 123 124 125	125 125	12 13 13	13 13 13 13	2 13 2 13 3 13 4 13	13: 14: 14: 14:	143 144 146	146 146 146 150	15 15 15 15	15 15 15 15	15 15 15 16	7 16 8 16 9 16 0 16	1 2 2
	ᆣ	•	10	-11	-	1	1		1	1.	8	1			10	,11		
25 36 87 38 39	115 116 116 117 118	117 118 118 119 120	119 120 120 121 121 122	121 122 122 123 124	123 124 124 125 125 126	125 126 126 127 127	127 128 129 130 131	130 131 132 133 134	13:	5 13: 3 14: 7 14:	) 144    145	148 148 149	151 152 153	150 150	150 150	163	164 165	
	•	•	10	11	<u> </u>	1	3	1	4	1	•	,	•	1.	30	11	•	
40 61 62 43 44	119 120 120 121 121	121 122 123 123 124	123 124 124 125 125	125 126 126 127 128	127 128 128 129 130	129 130 130 131 132	132 133 133 134 135	135 136 136 137 138	130	143 143 144	147 147	151 151 152	154 155 155 158 157	159 150	161 162 163 163 164	163 163 167 168	163 169 170 171	
		•	10	刯	-	1	•	匞	1	8	•	7	•		10	11	•	
45 45 67 48 49	122 123 123 124 124	124 125 125 126 126	126 127 127 128 128	128 129 129 130 130	130 131 131 132 132	132 133 133 134 134	135 136 136 137 137	139 139 139 140 140	142 142 143	146 147 147	150 151 152 152	155 156 156	157 158 159 160 161	161 162 163 164 185	164 165 166 167 168	169 170 171 172	174 175	
	1	-	10		-	IJ	1	•	٠	•	•	,	•	•	10	11	1	
50 51 52 63 54 55	125 125 125 125 125 125 125	127 127 127 127 127 127	129 129 129 129 129 129	131	133 133 133 133 133 133	135 135 135 135 135 135 135	138 138 138 138 138 138	141 141 141 141 141 141	144 144 144 144 144 144	149 143 143	152 152 153	156 157 157 157 158 158	161 162 162 162 163 163	165 166 166 166 167 167	169 170 170 170 171 171 171	173 174 174 174 174 174	176	

## Index

Abduction, 483 Abscess, 89 Absorption, 20, of food, see under Food Ac globulin of plasma, 103

a-c interval of venous pulse, 182

Acceleration effect of on circulation

154, 155, of heart, 188 Accelerator nerves, see under Nerve Accessory food factors, see Vitamins

Acclimatization to high altitude, 262 Accommodation of eye, 593ff-in new born, 679, of nerve, 467

Acetone 370 Acetylcholine, 567, action of on heart.

191, 192, sleep and, 572 Achromatic interval, see under Interval

Achromatism 601

Achylia gastrica, 318

Achroödexitrin, 300 acetic, 13, 360 acetoacetic, Acid(s) 370, 436-in urine, 275 amino, see Amino acids, ascorbic 391 carbonic, 12, 13, 14, 111, 112, 113, cholic, 331 373, description of, 14, excretion of 110, fatty-essential 399, 400 401, from fat, 334 372, folic, 76, 384 free in stomach, 313, 315, hyaluronic 51, hydrochloric, 7, 12, 13-and secretin 325, of gastric contents, 313, of gastric juice, 307, 308, 309, hydrocyanic-poisoning by, 261, 263, smell of, 643, hydroxybutyric, 370, 436, lactic-ac cumulation of in thiamin deficiency, 389, action of on capillaries, 203, buffering of in blood, 111 and fatigue of muscle 491 492, as a glyco gen former, 369, from glucose, 369 498, 499, in muscular con traction, 498, 499, oxidation of, 498, 499, in urine, 279, 280, linoleic, 400, 401, linolenic, 400, 401, nicotinic, 384, nitric, 12, 13, 14, nucleic, 361, 366, oleic, 372, osmic, 46, palmitic, 372, pantothenic, see Pantothenic acid. para aminobenzoic see Para aminobenzoic acid, phosphoric, 12 13, 14, 366, 367, pteroylglutamic, 76, stearic, 324 372, sulfuric, 12, 13, total in gastric contents, 313, uric, 366, 367, 374metabolism of, 366, in urine, 279, 280 Acid base balance 108ff kidney and

Acidosis, 113, in diabetes, 370, dyspnea in, 260, in nephritis, 284

Acinus (1), 39, 294, 295 Acoustic area, see under Area Acromegaly, 440

ACTH see Hormone, adrenocortico trophic

Actin, 497

Action currents of brain, see Electro encephalogram of heart 177ff also Electrocardiogram, of muscle, 494 of nerve, 469-71 of optic nerve 585, of retina, see Electroretinogram

Actomyosin, 42, 497

Adaptation dark and light of eye 580 581 of olfactory receptors, 643 of receptors see Glossary

Addison's disease, 420

Adduction 483

Adenine, 366, 367

Adenohypophysis, 430 431 432 436, effects of extracts of 432 433 436 functions of, 431ff overactivity of, 440

Adenoma of thyroid gland, 406 408 412 Adenosinediphosphate, 497, 498 499 Adenosinetriphosphatase (myosin), 500

(footnote) Adenosinetriphosphate, 497, 498, 499,

500

Adiposity, see Obesity ADP, see Adenosinediphosphate Adrenal cortex, 418, 419, 420 425-29

action of pituitary on, 455, in elasmo branch fish, 418 419, estrone in, 428 excessive activity of, 429 functions of, 425, and milk secretion, 455, pro gestin in, 428, steroids of, 425, 426 427, vitamin C in, 391, 428

Adrenal glands, 122, 418-29, see also Adrenal medulla and Adrenal cortex Adrenal insufficiency, 425, 426, basal

metabolic rate in, 359

Adrenal medulla, 418, 419-25, action of commercial extract, 420-23, nerves to, 565, overactivity of, 425

Adrenalectomy, effects of, 425 Adrenaline, 420-23, 425, action of on capillaries 203-on coronary arteries, 205, on heart, 192, 193, on pupil 608, calorigenic action of 380, 381, 423, and clotting of blood, 106, excessive production of, 425, liberated in heart

by sympathetic, 192, at sympathetic endings, 567 Adrenocorticotrophic hormone, see un der Hormone

Adrenosterone, 628 Adsorption, 19, 20

Aerobic phase of muscular contraction, 498, 499 African sleeping sickness, 210, 212

Afterbirth, 676, 679, 680 After discharge, 516

After images, 604, 605

Age and basal metabolic rate, 358, 359, and blood pressure, 143, and crystal line lens, 595, 610 611, and velocity of arterial pulse, 157

Agglutination of blood cells, 78, 79, 80, of bacteria 215

Agglutinins, 78, 79 215 Agglutinogens, 78, 79

Air, alveolar, 247, atmospheric composition of 245, 246-relative humidity of 56 317, of dead space 242 sub divisions of in lungs, 240, 242

Air passages 226ff Air sacs 229, 230, exchange of gases

and, 247

Alanine, ethylamine from, 342 Albumin(s), 361, egg 362, human for transfusion 83, serum 57 58, 361

Alcohol, effect of on body temperature, 382

Aldosterone, 427, 428

Alfalfa, vitamin K in 396 397 Alimentary tract 294ff sensitivity of, 346 347, in thiamin deficiency, 387 Alkalemia, 114

Alkalı reserve, 111

Alkalıs, 15 Alkalosis 113, 114

Allergy, 224, cosmophil leucocytes in, 90 All or none" law, see under Law

Alpha particle, 5 Alvarez, Dr W C., 343

Alveolar ducts, see under Duct Alveolus(i), 39, of glands, 294, 295, pulmonary, 229

Amboceptor, 217 American Indian, Rh positive, 81 (foot

Americanum, 3, 4 Amidopyrine, a cause of reduction of

leucocytes, 92 Amino acids 360, see also Appendix, chains of in muscular contraction 500 deamination of, 363, essential 364, 365, indispensable, see essential number in protein molecule, 362, par tial list of, 360

Amitosis, 25, 27, 28 Amitotic division, see Amitosis Ammonia, in urine, 274 Amnion, 667, 669, 670 Amoeba 22, of dysentery, 210, 212, respiration of, 225

Amoebic dysentery, 210, 212
Ampulla(ae) of semicircular canals,

630, 631, of Vater, 322, 332, 333 Amyl nitrite, in angina pectoris 205 Amylase(s), 292, 293, pancreatic see Amylopsin, salivary, see Pivalin Amylopsin, 292, 293, 323, 324

Anabolism, 350 Anacidity of gastric contents, 314 Anacrotic limb, see Pulse, arterial Anaerobic phase of muscular contrac-

tion, 498, 499 Analysis of gastric contents, 313 Anaphase, 26 27 Anaphylactic shock, see Shock Anaphylactogen, 223

Anaphylactoid reactions, 224 Anaphylaxis, 222, 223 Androsterone, 452

Anelectrotonus, 466 Anemia, 73ff anoxia in, 261, 262

aplastic. 76, cardiac output in, 153 due to poisoning of bone marrow, 76 dyspnea in 260, hemolytic 73, iron deficiency, 73, nutritional, 74, per nicious, 74, 75-jaundice in, 335, post hemorrhagie, 73, protein deficiency,

73, 74, vitamin deficiencies and, 74 Anemic anoxia, see Anoxia Anesthetics effect of on body tempera ture 382, effect of on nerve, 472, ef

fect of an reflex centers, 515 Anestrus, 446, 447 Aneurin, see Thiamin Angina pectoris, 205, referred pain in

Angiotonin 145 Anhydremia, 61, 148 "Animal starch," see Glycogen Anons 13

Anisotropic bands, 44 Ankle jerk, 519 Anode, 12, 13, 465, 466 Anomaloscope 602

Anomalous irichromatism, 602 Anoxia 261, and carotid body, 256 ef fect of on heart, 193, 205 effect of on reflex centers 515, effect on respira

tion, 255, in heart failure, 262, of heart muscle, 205, types of 261, 262 Anterior chamber of the eye 575 Anterior poliomyelitis 210 521, arti

145-51, fluctuations of, see phases of,

human, 141, 142 and labyrinth, 634,

mean, 138, measurement of-in ani-

mals, 137-39, in man, 141, 142, in

725

ficial respiration in, 239, natural immunity to, 217 Anthrax bacilli, 212 Anti A (antibody, agglutinin), 78 Antianemic factor, 74 Antı B (antibody, agglutınin), 78 Antibacterial agents, 22 Antibiotics, 221, 222 Antibody(ies), 215, against corpuscles, 78, 79, 80, gamma globulin and, 91, produced by lymphocytes, 101 Antibody antigen reaction, 216 Anticoagulants, 104 Antigen(s), 216, 218, of blood corpuscles, 80 Antigen antibody reaction, 216 Antimony tartrate, as an emetic, 319 Antiperistalsis, 337 Antipernicious anemia factor, see Vitamin B Antipyrine, effect of on fever, 382 Antirachitic vitamin, see Vitamin D Antisepsis, 211 Antitoxin, 215 Anus, 322 Aorta, 118, 121, 122, 123, 167, of cold blooded heart, 166 Apex beat, 183 APL hormone, see Hormone, anterior pituitarylike Apnea, 255, 258, 259 Apocrine gland, see under Gland Apomorphine, as an emetic, 319 Apoplexy, see Cerebral hemorrhage Appendix, see Vermiform appendix Appetite, 347, 348 Aqueous humor, 575, 591 Arachnoid granulations, 556, 560, 561 Arachnoid mater, 556, 557 Area(s): auditory, 535, 536, acoustic, 535, 536, motor, 431, 432, 433, 533, 634 olfactory, 641, premotor, 533, 541, psychoauditory, 536, psychovisual, 536, "silent" of cerebral cortex, \$35, somesthetic, 534, 535, 546, visual, 536, 578, 589 Areola(ae), 46 Aristotle's experiment, 646

Arrectores pilorum, 35

Arterial blood pressure, 136ff, action of adrenaline on, 420, 423, action of

noradrenaline on 423, action of piturtrin on, 437, in anaphylaxis, 224, and

carotid sinus, 258 and coronary circu-

lation, 205, diastolic, see phases of, effect of physiological conditions on

142, effect of splanchnic stimulation

on, 147, 195, excessive, see Arterial

hypertension, factors maintaining,

muscular exercise, 504, phases of, 137, in pulmonary artery, 131, 206, 207, in secretion of urine, 277, 278, sex difference, 142, in sleep, 143, 572, systolic, see phases of, in traumatic shock, 223 Arterial disease, 144, 145 Arterial hypertension, 143, kidney disease and, 143, in nephritis, 284, in obesity, 374 Arterial pulse, see Pulse, arterial Arteriole(s) action of adrenaline on. 421, afferent of renal glomerulus, 269, 270, 271, 272, 277, constriction of, 195-effect of in overcoming gravity, 153, control of, 195ff, dilatation of, 196-in inflammation 93, efferent of renal glomerulus, 269, 270, 271, 272, in hypertension, 144, nerves to, 195, reflex changes in, 198, in shock, 223, structure of, 124 Arteriosclerosis, 144, 145 Arteriovenous anastomosis 376 Artery(ses), 126, aorta, abdominal, 122, axillary, 123 brachial, 123, carotid 118, 122, 123, 200-and sleep, 572 celiac axis (branches of), 122, central of retina, 575, coats of, 123, common iliac, 121, 122, corkscrew in endometrium 448, coronary, 204, 205absence of in frog's heart, 164, action of adrenalme on 421, disease of, 144, 145, and perfusion of heart, 162, clas ticity of, 145, 148, 149, external iliac, 121, hepatic, 122, 208, 330, hypogastric, see internal iliac, inferior mesenteric, 122, innominate, 118, 122, internal iliac, 121, "pipestem," 145, 148, pulmonary, 108, 118, 119, 121, 122, 206-in fetus, 673, pressure in, 131, secoil of, 149, 150, renal, 122, 270, 272, structure of, 123, subclavian, 118, 122, superior mesenteric, 122, umbilical, 672 Artificial respiration, 236-40 Arytenoid cartilages, 265 Asphyxia in anaphylactic shock, 224, in tetany, 415, 517, resuscitation from see Artificial respiration, see also Апохіз Aspirin, effect of on fever, 382 Aster(s), 25

Asthma 228, adrenaline in, 421, cortisone in, 427, dyspnea in, 260

Astigmatism, 612-14

Astral rays see under Rays	synthesis of vitamin k by 397 syn
Astrocytes 506	thesis of vitamins by 212 390 391
Ataxia cerebellar 555	397
Atheromatosis 144 145	
	Bacteriolysins 215
Atherosclerosis 144 145	Bacteriophage 210
Athletics 503 504 505, and cardiac	Bacteriostatic substances 221 222
disease 194 505	Bad lighting 614 615
Atmosphere composition of 245 246	Balance acid base see Acid base bal-
Atom(s) 1 carbon 4 hydrogen 7 in	ance of body 630ff energy 374
destructibility of 11 sodine 1 iron	heat 375 nitrogen 363, water 55
1 oxygen 1 sodium 8 structure of 1	Barcroft Joseph 97
Atomic numbers 2 3	Barlow's disease 392
Atomic weights 3	
	Barr Murray 507
ATP see Adenosinetriphosphate	Basal metabolic rate 356-59 action of
Atrium(a) of heart 115 116 117 118	adrenatine on 422 calculation of 358
122 of lungs 229	at different ages and in two sexes 359
Atropine action on heart 188 191 ac	effect of thyroid on 405 in hyper
tion on pupil 508	thyroidism 411 in hypothyroidism
Audiogram 630	410 in planning a diet 400 under
Aud tory ossicles 617 618 622	various physiological and pathological
Auditory pathway 621	conditions 359
Auditory strings 624 626	Bases description of 14
Aureomycin 222	Basophil(s) 87
Auricle of ear 617 of heart see Atrium	Bayliss Sr William 325
Auricular flutter 185	"Beading of ribs" in rickets 394
Auticuloventricular block 160 186	Beat of heart see Heart beat
Auriculoventricular bundle 166 168	Beaumont William 317 318
169 182 189 action of digitalis on	Bed wetting 287
191 conduction in 168 183 185 187	Benedict Roth apparatus 357
188	Benzine as hemolytic agent 85
Auriculoventricular valves 118	Berger Hans 536
Auscultatory method of measuring blood	Beriberi 386 387
pressure 141 147	Bernhard Claude 52 198 368 372
Autointoxication 342	Biceps jerk 519
Autonomic nervous system 561ff cen	Bile 328ff circulation of 330 334 com
ters of 562 563 functions of 564	position of 378 329 330 concen
	tration of 332 discharge from gall
	bladder 331 formation of in Iver
sion 56? 563 564 565 566 sympa	330 functions of 333 334 hydro-
thetic division 563 564 565 566	330 functions of 335 334 man of 329
Autosomes 665	trop c action of 334 origin of 329
Avidin and biotin deficiency 390	required for absorption of viamin k
Avitaminosis see Vitamins deficiencies	107 397 secretion of 331 333, 334
of various vitamins	storage of 331 332
Axis(es) of lens 591 visual 607	Bile pigments 329 330
Axis cylinder 460	Bile salts, 329 331 334
Axolotyl 405 407	Bilirubin 329 330 in jaundice 335
Axon 508 509 hillock 508 reflex see	Biliverdin 329 330
under Reflex	Binary fission 27 28
	Biotin 384 390 396
Bab nsky reflex or sign 519 527 539	Blackout " 155
in newborn 679 in sleep 571	Bladder see Urinary bladder
Bacilli 212 213	Blastocyst 667 668 669
Bacitracin 222 (footnote)	"Bleeder " 107 108
Bacteria 210ff destroyed by leucocytes	Bleeding time 106
89 90 factors of vitamin B complex	Blind spot 578 579
required by 391 of feces 345 fever	Blinding glare 615
required by 391 of feces 345 fever carsed by 381 pyogen c 215 syn	Blindness 615 616 night see hight
thesis of vitamin B complex by 391	blindness
synthesis of vitamin C by 212 391	Block auruniar ventricular 169 186

auru-ular ventricular 169 186

bundle branch, see intraventricular, intraventricular 169, 182

Blood 52ff buffers of, 110, cells of, see Erythrocytes, Leucocytes, Thrombocytes, circulation of, see Circulation of blood, clotting of, see Coagulation of blood, coagulation of, see Coagulation of blood, corpuscles of, see cells of, defibrinated, 105, functions of, 53, 54, glucose of, see Blood sugar, hemolysis of, 84, incompatibility of, 79, laking of, see Hemolysis of, plasma, 57, see also Blood plasma oxygen absorption by, 69, pH of, see pH, platelets of see Thrombocytes. proteins of, see Plasma proteins reaction of, 108, regeneration of 71, Rh negative, 81, 82, Rh positive, 81, 82 rouleaux formation in 59, specific gravity of, 57, transfusion of, 77 velocity of, 133, see also Blood ve locity, viscosity of, 148, volume of, see Blood volume, water of, 55

Blood counts 60

Blood dust, see Hemoconia

Blood groups, 78, 79 80 Blood plasma antithrombin of, 104, bilirubin in, 335, colloids of, 18 com position of (Table), 58, deproteinized, 278 globulins of, 144, osmotic pres sure of, 65, pooled 82, proteins of, 57, 58 separated, 252, transfusion of, 77, 82, "true " 252

Blood pressure throughout vascular sys tem, 130, see also Arterial, Capillary,

and Venous blood pressures Blood proteins, see Plasma proteins Blood sugar, 349, 368, 369, adrenaline

and, 422, in diabetes 370, 371, and hunger, 349 Blood velocity, 133-36

Blood vessels control of 195 see also Arteries, Arterioles, Capillaries, and Veins

Blood volume, 63, 64, 66, 67, in control of body temperature, 377, in lungs 206 measurement of, 64, regulation of, 64, 66, 67 "Blue babies" 674

BMR, see Basal metabolic rate

Body(ses) carotid 201, central of cell

25 chemical composition of, 3, ciliary, 591, defence mechanisms of 210ffsympatho-adrenal system and, 423, 424, geniculate, 545-lateral 587, 589, medial, 621, heat production by, see Basal metabolic rate, inclusion 24, 25, Nissl, 507, 508 pineal 458 pituitary, see Pituitary body, polar, 661, 662 664 (footnote), restiform, see Peduncles, cerebellar; surface area of, 356, 358-chart for determining, 358, temperature of, see Body temperature, tigroid 507, vitreous, 575, 591, vellow, see Corpus, luteum

Body temperature, 375ff, centers controlling, 380, 381, chemical factors controlling, 379, 380, effect of drugs on, 382, physical factors controlling, 376, 377, 378, in sleep, 571

Body weight excessive, see Obesity, loss of, in diabetes 370, 371

Bohr effect 252 Boil " see Furuncle Boiler maker's disease, \$25

Bomb calorimeter, 352, 353 Bone(s), 48ff, and vitamin A deficiency, 386, cancellous, 49, cells of, 49, com pact, 49 composition of, 48, 59 de mineralized, 48, 49 effect of para thyroids on, 417, 418, femur, 50 in hyperparathyroidism, 417, isotopic

phosphorus and, 6, pituitary and, 455 red marrow of, 50, 51-failure of as cause of anemia, 76, in hemopolesis 70, 75, increase at high altitudes 62, in rickets, 393, softening of, 395 spongy, see cancellous, structure of 48, 49, trabeculae of, 50, turbinate 226 227, yellow marrow of, 49

Bowel(s), see Intestine Bowman Sir William, 276 277 Bowman's capsule, 269, 270 271, 276 277

Brachium(a), 545, pontis, 553, conjunc tıvum, 553

Bradycardia 186 Brain 529ff buoyancy of 559, interpre tation of sensory impulses by, 477

Brainstem see Midbrain, Pons, Medulla obiongata Bran laxative action of, 346

Branchial arches, 671, 672 Branchial clefts, 671, 672 Break shocks, 465, 466 467 Breathing, see Respiration Brewer's yeast, 396

Bright's disease, see Nephritis Bronchiole(s), 226 227, 228 229, ac tion of adrenaline on, 421, in anaphy

laxis, 224, in asthma 228, respiratory, 229, terminal, 229, 230

Bronchus(1), 226, 227, 228, 229 Brown Robert, 19

Brownian movements, 19 Brush border, 270

Bucket handle movement of ribs in respiration 235 236

Budding of yeast cells, 28 "Buffalo neck," 444 Buffers, of blood and body fluids, 110, in perfusion fluid, 165

Bulb, olfactory, 226 Bulbus arteriosus, 166

Bundle branch block, see Block, intraventricular block Butter, caloric value per pound, 400

Caffeine, diuretic effect of, 282

Calcuferol, 392

Calcium in blood, 58, in bone, 48, in clotting of blood, 103, daily requirement of, 401, equivalent weight of, 9, in hypoparathyroidism 416, increased excretion of in hyperparathyroidism, 417, isotopes of, 5, metabolism-effect of vitamin D on, 393, and parathyroid glands, 415, paracasein, 309, rigor,

162, in tetany, 416, in urine, 274 Calone(s), 351, evolved per liter of oxygen consumed, 354, 355, per square meter of body surface, 359, requirement of diet, 399, values of different

foods, 354

Calerimetry, 353ff

Canal(s) anal, 321, central, of spinal cord, 523, Haversian, see Haversian canals, semicircular, see Semicircular canals

Canaliculus(1) biliary, 328, 330, of gas-

tric glands, 308 Cane sugar, see Sucrose

Cannon, W B 423

Capillary(ies) actions of adrenaline and pituitrin on, 203, actions of chemicals on, 203, blood pressure, 151, circulation, 201, 204, see also Capillary circulation dilatation of, in inflammation, 93, loops in skin, 203, of lung, 230, nerves of 200, permeability of, 66, in purpura, 108, of renal glomerulus 269, 271, 272, 277, in shock, 223, of skin, 203, 204, structure of, 124, 125, 126, vitamin C and, 391

Capillary circulation 201-04 "Capillary lake," 135

Capon, 445, action of testosterone, 452 Carbohemoglobin, 252

Carbohydrate(s), 367ff, absorption of, 343, 344, 345 combustion of in colorimeter, 353, cycle, 367, dietary require

ment of, 400, digestion of, 324, 327, 368, history of in body, 368, 369, metabolism of, see Carbohydrate me tabolism, oxidation of, 369, 371, source of energy in muscular contraction, 497, 499, 503, sparing effect of, 364

Carbohydrate metabolism, 367ff, action of adrenaline on, 422, adrenal cortex and, 426, 427, in hyperthyroidism, 411, hypothalamus and, 542, pituitary and, 432, 435, 436

Carbomycin, 222 (footnote)

Carbon equivalent weight of, 9, isotopes

Carbon are light, 393

Carbon dioxide, 11, action of on capil laries 203, in alveolar air, 246, in artificial respiration, 240, in atmosphere, 245, 246, 'blowing off" of, 260, and carotid body, 258, in combination with hemoglobin, 252, determinationof BMR from, 356 357, of cardiac output from, 133, diffusibility of, 248, effect of on heart, 193, effect of on respiration 254, 255, elimination of, 248, 255-in control of blood reaction, 110, 112, exchange of in respiration, 243, in expired air, 246, from oxida tion of carbon, 255, 351, partial pres sure of-in arternal blood, 248, in in spired air, 246, in venous blood 248, production of by nerve 478, in saliva, 298, transport in blood 252, volumes percent-in arterial blood, 248, in venous blood, 248

Carbon monoxide, 11; poisoning 262, Carbonic acid, sodium bicarbonate ratio,

Carbonic anhydrase, 112

Cardiac cycle, 169ff Cardiac failure, ve Heart failure Cardiac muscle, see Muscle, cardiac

Cardiac nerves, see under Nerve Cardiac output, 132, 133, measurement of, 133, in muscular exercise, 504 un-

der various conditions, 132, 133 Cardiometer, 174

Cardiospasm, 303 Carlson, Dr A J, 310, 348

Carnosine, 497 Carotene, 385

Carolfd body, 257, 258, 260 Caroted sinus, 119, 256, 257, 258, in control of circulation, 199, in control of respiration, 257, 258, hypersensi tivity of, 200, in muscular exercise, 504

Carpopedal spasm in tetany, 416 417 Cartilage, 47, 48, elastic, 47, 48, hyaline,

Caryocinesis, see Mitosis Caryoplasm, 23 Caryosomes, 24 Casein, 361, 365

Caseinogen, 309 Castle, Dr W B, 75 Castle's intrinsic factor, 75 Castor oil, 346 Castration, 445 Catabolism, 350 Catacrotic limb, see Pulse, arterial Catalysts, 290, 291 Cataract, 591, 615, 616 Catelectrotonus, 466 Cathartics, 346 Cathode, 12, 13, 465, 466 Cathode ray oscillograph, 471 Cations, 12, 13 Cattle, hemorrhages in, 104, 107 Cavity(ies): amnionic, 669, nasal, 226, 227, pleural, 231, tympanic, see Tympanum Cecum, 289, 320, 321, 339 Cell(s), 23ff, acidophil, 431, 432, of anterior horns of cord, 507, basophil of pituitary, 431, 432, border, of gastric glands, see parietal, cartilage, 47, of cerebral cortex, 532, chief of gastric glands, 307, 308, chromaffin, 418, 419, chromophobe, 431, 432, connective tissue, 45, daughter, 25, 26, 27, 662, differentiation of, 29, egg see Ovum, Oocyte, endothelial, 38-form ing capillary wall, 124, 125, 201, fat, 45, 46, ganglion-in heart, 187, of parasympathetic, 562, of retina, 576, 577, 589, of sympathetic, 563, germ, see sex, giant pyramidal of cerebral cortex, 507, goblet, 28, 326, 327, hair -of organ of Corti, 620 of semi circular canals, 631, 634, interstitial of testis, see of Leydig, Kupffer, 95, 329, of Leydig, 452, 454, 656, 657, of liver, 208, mitral, 643, mucus secreting, 295, multiplication of, 25, see also Mitosis and Meiosis, muscle, 41, nerve, 506, 507, 508, neuroglia, 506, 507, oligodendroglia, 506, 507, oxyntic of gastric glands, see parietal, of Paneth 326, 327, parietal, of gastric glands 306, 307, 308, pigment, of retina, 577, 578-of frog s skin, 438, of pitui tary body, 431, 432 Purkinje, of cere bellum 507, 552, 553, red, see Ery throcytes, reticulo-endothelial, 86 88, 94, 95, 97, 329-in spleen 97, of retina 576, 577, 578, Rouget, 125, 202, Sertoli, 657, sex, 648, 649-con jugation of, 662, 663, maturation of, 660, 661, meiosis of, 660, 661, see also Ovum oocyte Spermatogonia, Spermatozoa of skin 33, spermatogenic, 656, 657, of spinal cord, white,

see Leucocytes, yeast, 27, 28, zymogenic, of gastric glands, see chief Cellulose, 368, as roughage, 345, 346 auditory, 627, cardioinhibi-Center(s) tory, 187, 189, expiratory, 253, 254, "feeding" 349, hearing, 535, 621, 627, heat controlling 380, inspiratory, 253, 254, micturition, 287, olfactory, 642, pneumotaxic, 253, 254, primary visual, 587, reflex, 511, respiratory, 252, 253, 254, 257, 260-effect of anoxia on, 255, salivary, see salivatory, salivatory, 296, 297, 567, sweating, 378, taste, 640, vasoconstrictor, 196, 197, vasodilator, 196, 197, vasomotor, 196, 197, visual, 536, vomiting, 319 Central body, 25 Central canal, of spinal cord, 523, 524 Central excitatory state, 516 Central inhibitory state, 517 Central nervous system, 506ff Central sulcus, 531, 534, 535 Centriole(s), 25 Centripetal acceleration, effect of on circulation, 155 Centrosome, see Centrosphere Centrosphere, 25 Cephalic phase of gastric secretion, see Psychic phase Cephalm, 373 Cerebellar ataxia, 555 Cerebellar cortex, 552, 553 Cerebellum, 529, 531, 544, 551-55, 558, functions of, 555 Cerebral apoplexy, 539 Cerebral aqueduct, 542, 543, blockage of, 561 Cerebral circulation, 208, 209 Cerebral cortex 530ff, areas of, 531-36, cell layers of, 531, 532, inhibition of, and sleep, 573, removal of, 542 Cerebral hemorrhage, 539, caused by flying maneuvers, 155 Cerebrospinal fluid 209, 557, 558, circulation of, 560 Cerebrum 529ff Cervix, of uterus, 651 cH. 110 Chalone, 314, 403 Cheese caloric value of per pound, 400, proteins of, 365 Chemical elements 1, of body, 3, table of, 2, "trace," 402 Chemical formulas, 6 Chemical reactions, 7 Chemical symbols 6 Chemical transmission of nervous effects. 191, 192 Chemoreceptors, 258

Chemotherapy, 220, 221, 222 Cheyne Stokes respiration 259, 260 Chickenpox, 210

Childbirth, see Labor, transfusion of Rh positive blood after, 82 'Chill' at commencement of a fever.

382 Chinese, Rh positive blood, 81 (foot

note) Chloride shift, 112

Chlorine, 1, equivalent weight of, 9, isotopes of, 5

Chloroform, and hemolysis, 85 Chloromycetin, 222 (footnote) Chlorophyll, 69, 71, in photosynthesis,

350 Chlortetracycline, 222 (footnote)

Cholecystokinin 333 Cholesterol 373 in

Cholesterol 373, in adrenal cortex, 428, in atherosclerosis 145, in bile 329, in blood 57, in erythrocytes 59, of skin, irradiated 392, and testosterone, 452

Cholesterol esters 373 Choline, 191, 373, 384

Cholinester, see Acetylcholine Cholinesterase, 192

Chordae tendineae, 117, 175 Chorion, 667

Choroid of eye, 575 592, 593, 596 Choroid plexuses, 560 561 Chromatic aberration 610 611

Chromatin 23, 24, 666, sex, see Sex chromatin

Chromoprotein, 361 Chromosome(s), 26 27, 660, 661, 662, 663, number in various species, 663 (footnote), sex, 660 665, X and Y,

see sex Chronaxi 467-69 Chvostek's sign, in tetany, 417

Chyme, 317 Cilia, 30, 37, 220 of nasal mucous membrane, 226

brane, 226
Ciliary muscle 614
Circulation of blood 115ff, discovery

of 120, greater, 119, lesser, 119 206 models of, 126, in muscular exercise, 503 504, pulmonary, see lesser, sys temic, see greater, see also Capillary, Coronary, Cerebral, Pulmonary circu-

lation, etc Circulation rate, see Cardiac output Cisterna(ac) of subarachnoid space,

559, 560
Cisternal puncture 561 (footnote)
Citrates as anticoagulants, 104
Citrus fruits, vitamin C in 391
Clearance, see Plasma clearance
Cleft of pituitary, 430

Coagulation of blood, 102ff, action of adrenaline on, 422, failure of, 107, in hemorrhage, 76, means used to hasten 105, vitamin K and, 104, 105, 397, see also Thrombosis

Coagulation time, 106
Cobalt, necessary for blood regeneration,
72, 401

72, 401 Cocci, 212, 213 Cochlea 617, 618 619, 621, 622, 623,

piezoelectric effect in, 627 Cockerel, castration of, 445

Cod liver oil source of vitamins A and D, 396

Coffee, diuretic effect of, 282 Costus 657

Collisculus(1), 545, 621 Collisculus(1), 545, 621 Collist, Prof J B, 415

Colloid(s), 17, 18, hydrophilic and hy drophobic, 17 (footnote), internal and external phases of, 18, of thyroid

gland, 404, 405, 408
Colon, 320, 321, ascending, 289, de scending, 289, 320, 321, pelvic, 289

320, 321, spastic, 340, transverse, 289
320, 321

Color(s) blindness, see Color blind
ness, complementary, 603, confusion
602, primary, 600, qualities of 600

ness, complementary, 603, confusion 602, primary, 600, qualities of 600 retinal areas sensitive to, 600, vision see Color vision Color blindness, 601, partial, 602

Color bindness, 601, partial, Color vision, 599, 601
Colostrum, 455 (footnote)

Columns of spinal cord, 524 Coms, in diabetics, 370 Comfort zone, 380

Common cold, 210 Complement, 217 Compgunds, 8

Conception, 675 Concha(ac), 226, 227

Conditioned reflexes 567-71, inhibition of 569, 570, and salivary secretion, 298, and sleep, 573

298, and sleep, 573
Conduction in heart 161, 187, in nerve
464, property of protoplasm, 23, over
reflex arc, 515

Conductivity of heart 161, 187, of nerve 464, of protoplasm, 23

Cones of retina, see Rods and Cones Conjugated proteins, see under Protein Conjugation, of sex cells, 662, 663 Conjunctiva, 574, 575, axon reflexes in

521 Consciousness, thalamus and 539 Constitution 340, 341, and thiamin d

Constipation 340 341, and thiamin de ficiency, 387

Constrictor pupillae, see Iris Contractility, of cardiac muscle, 159, of striated muscle, see Muscular contractraction

Contraction heat, 501, 502 Convolutions of cerebral cortex, 53 Convulsions insulin, 371, in tetany, 417

Copepods, and vitamin D, 392 Copper in blood regeneration 72, essential in diet, 402, sulfate, 12, 13

Corium, see Dermis Corn, proteins of, 365

clusion

Cornea, 574, 575, dryness of, in vitamin A deficiency, 385, opacities of, 614in riboflavin deficiency, 389 Corona radiata, 649, 650 Coronary circulation, 204, 205

Coronary occlusion, 205, atherosclerosis and, 145, dicoumarin in, 104 Coronary thrombosis, see Coronary oc-

Corpus(ora) callosum, 531, 543, 558, cavernosum penis, 658, 659, caver nosum urethrae, 658, 659, luteum (corpora lutea), 448, 449, 451, 456 650, quadrigeminum 543, 544 545. 554, striatum, 533, 537, 539, 540, 541 Corpuscie(s) of blood, see Erythrocytes, Leucocytes, Hassall's 457, 458, Maipighian, of kidney, 268-of spleen, 95. Meissmer's, 645, pacinian, 34, 644, 645 Corresponding retinal points, 588, 589

Cortex of adrenal, see Adrenal cortex. of cerebellum, see Cerebellar cortex, of cerebrum, see Cerebral cortex, of kidney, 273 of thymus, 457, 458 Corti's organ, 619, 620, 621

Corticoids, 426 Corticospinal tracts, 526, 627

Corticosterone, 427 Cortisone, 426, 427

Cough, 220 252, 258 Cowpox, 218 Cranial outflow, of parasympathetic 562

Cramotabes 394

Creatinine, 363, 497, in urine, 274 Crenation, 62, 63, 68

Cretin, 408, 409, 410 Cretinism, 408, 409, 410

Cricoid cartilage, 265 Crista(ae), of semicircular canals, 630,

631 Crop gland, 456 (footnote) "Crosseyes," see Strabismus

Croton oil, 346 Crying, 258 Crypts of Lieberkühn, 326 327, 344 Crystalline lens, 575, 591, 592 593 595, changes in shape of with age, 410, de-

fects of, 609, 610 discission of, 599, extraction of, 615, 'needling" of, 599, opacity of, see Cataract Cumulus oophorus, 649

Cupula, 619, 630 Current(s) action, 469, 470 electrical, see action

Curve(s) of gastric acidity, 315, intraaortic pressure, 171, intra auricular pressure, 173, 174, intraventricular pressure 171, monophasic, 471, oxygen dissociation curve of hemoglobin, 249, 250, 251, strength-duration of nerve,

Cushing, Dr Harvey, 444 Cushing's disease, 444 Cuticle, see Epidermis Cutis vera, see Dermis Cyanide poisoning, 261, 263

Cyanosis 36, 204, 263, 264, in circulatory defects, 674, in heart disease 194 Cycle. carbohydrate, 369, cardiac 169ff, estrous, 446, menstrual, 447, 449, 453 Cyclotron, 5 Cystine, 360 Cytolysins, 215 Cytoplasm, 24

Dale, Sir Henry, 567 Dalton, John and atomic theory, 1, color blindness of, 601, 602

Dead space 242 Deafness, 629 Deaminase(es), 292 Deamination, 363

Decerebrate rigidity, 522 Decidua 669

Decomposition bacterial, of protein, 242, 243, chemical, 8 Decussation of corticospinal tracts see

of motor tracts, of Forel, 542, 545, of motor tracts 526, 546, of optic nerve fibers, 586, 587, of pyramidal tracts, see of motor tracts, sensory, 546 Defecation, 340, 341 Deflections, of electrocardiogram, see

under Waves Degeneration of nerve reaction of, 463,

retrograde, 462 Deglutition, see Swallowing

Dehydration, 56, fever caused by, 56,

382 7 Dehydrocholesterol, irradiated in skin. 392

Dehydrocorticosterone, 427 Dehydrogenase(s), 10

Demilunes of Gianuzzi, 295 Democritus, 1

Dendrites, 508, 509

Dental carres, 393 396 Dental enamel from ectoderm, 668 Dermis 33, 34

Desivae process, 82 Desoxycorticosterone, 427

Deuteranomaly, 602 Deuteranopia, 601

Dextran, as a transfusion material, 84 Dextrus 368

Dextrose, see Giucose Diabetes, 370, 371, cataract in, 616, caused by nervous lesions, 372, hunger in, 349, ketosis in, 370, 371, and obesity, 374, and pituitary, 432, 436,

susceptibility to infection in, 220 Diabetes insipidus, 281, 444

Diabetogenic effect of pituitary, 432, 436,

Dialysis, 18 Diapedesis, 88

Diaphragm, 122, 234 235, 256 Dicoumarin 104, in coronary throm-

bosis, 104

Dierotic notch of pulse tracing, 158 Diestrus, 447

Diet and atherosclerosis 145, and con stipation 342, following hemorrhage, 77, planning of, 397ff, and regenera tion of blood, 72, and regulation of intestinal movements, 337

Dietetics 397ff Differentiation of cells, 29, in condi-

tioned reflexes, 570

Diffusion of gases, 244 Digestion 288ff, bacterial 242, 243, of carbohydrate 324, 327, 368, of fat, 324, 334, 372, in intestine, 320ff in mouth 294 of protein 323, 324 327, by saliva, 298, 300 in stomach 304ff

Digitalis, action of on heart, 191 Dilatation, of heart, 177

Dilator pupilae, see Iris Dinstrophenol effect of on body tem perature, 383

Dioses, 367 Dipeptids, 324 Diphtheria antitoxin 217 Diphtheria bacilli, 213

Diplococci 212, 213 Diploid number of chromosomes, 661

Diplopla, 589

Disaccharides 367, 368 Discission of lens 615 Discus proligerus 649, 650, hyaluroni

dase and 657 Dissociation see Ionization, 12, 13

Diuresis 282, caused by water drinking. 282 283

Digretics 282

Domagk, Dr G, 221

Donor, 77, 78, universal, 79 Dorsiflexion, 483 Drinker's method of artificial respira-

tion, 239 Drugs, action of on heart, 191

Drum of ear, see Tympanum Duct(s), 39, alveolar of lungs, 229, 230,

bile, 330, cochlear, 619 620, 622, 631, common bile, 331, 333, cystic, 331, 333, ejaculatory, 655, 656, 659 endolymphatic, 631, hepatic, 331, 333, lacrimal, 574, 576, lactiferous 652, 653, pancreatic, 333, of Rivinus 294, Stenson's, 294, Wharton's 294

Ductus arteriosus, 673 Duodenal ulcer, 308, 309, 315, mucin in.

Duplicity theory of vision, see under

Theory Dwarfism 441, 442 Dwarfs, 441, 442

Dyspnea, 260 261, in angina pectoris, 205, in heart disease, 194, 260, in hyperthyroidism, 260, in muscular ex ercise 260

Dystrophia adiposogenitalis, 441, 442

Ear(\*), 617-638, in embryo, 671, in ner-nervous connections of 554, see also Labyrinih middle 617-model of, 623, outer, 617, "pricking of," 545,

structure of, 617-21 Ectoderm, 668 669

Ectopic gestation 658 Edema, 101, 102, in heart disease, 194.

in nephritis, 284 Efficiency, formula of, 496 Egg(s) proteins of 365, white, 17-

raw, avidin in, 390 excretion of in urine 343 (footnote) Ehrlich, Paul 216, 220, side-chain the

ory, 216, 219 Einthoven, Wilhelm, 178

Ejaculation 656 Elasmobranch fish, adrenal glands in. 418, 419

Elastin 361 Electrical changes in nerve, 469-71

Electrocardsogram, 178ff normal record, 180, 181

Electrocardiograph, 178 leads of 179 Electrodes, stimulating 484 485 Electroencephalogram, 536 537, 538

Electrograms, 178 Electrolysis, 12 Electrolyte(s) 14 amphoteric 361

Electromagnet of electrocardiograph 179 Electrons 3, 4

Electroretinogram, 585 Electrotonus, 465-67 Elements, see Chemical elements death of in vitamin E defi Embryo ciency, 395, definition of, 675, devel

opment of, 666ff Embryonic disc, see Embryonic shield Embryonic shield, 667

Emergency theory, of function of adrenal medulla, 423

Emetic(s), 319 Emmetropia, 611

Emotion and adrenal medulla, 424. after excision of cerebral cortex ("sham rage"), 542, cardiac output in, 132, effect of on blood pressure, 142,

effect of on respiration, 257, 259, hypothalamus and, 542, sympathetic

and, 424, 566 Empirical formulas, 6

Empyema, 231 Emulsions, 16, 17 Encephalon see Brain End feet, 509, 510 Endocardium, 117 Endocrine glands, 403-456

Endolymph, 620, 630, 633, 634 Endometrium, 651, implantation of ovum in 669, in menstrual cycle, 447, 448

Endomysium, 43 Endoneurium, 459, 460 Endothelium, 38 Endothermic reaction, 350 Enterogastrone, 314, 327

Entoderm, 668, 669 Energy balance, 399, 400, calculation of, 497, of food, 350, 351, forms of, 22, 351, of heart, 134, light, 583, solar, 350, of sound, 622, source of in muscular contraction, 497-501, 503, storage of in body, 352, stored in

arterial walls during systole, 150, transformation of chemical to me chanical by muscle, 499, 500 Enzyme(s), 288-293, digestive, 290, effect of morganic salts on, 292, factors

influencing action of, 291; of intestinal ruice, 327, nomenclature, 292, 293, optimum temperature, 291, poisoning of, 292, reversible action of, 290

Eosinophil leucocytes, see Leucocyte, granular Eosmophilia, 87, 91

Epinephrine, 420, see also Adrenaline

Epicardium, 115 Epidermis, 33, 34 Epididymus, 655 Epiglottis, 266 Epimysium, 43

Equilibrium nitrogen, 363, organs of, 630ff Equivalent weights, 9 Erb's sign, in tetany, 417 Erection, 659, nerves controlling, 196 Erepsin, 324, 327 (footnote) Ergosterol, 373, irradiated, 392, 393

Epithelium, 29ff, germinal, 449, 648,

glandular, 38, 39, olfactory, 226, 227,

642, origin of from ectoderm, 668, see

Erythroblastosis foetalis, 82 Erythroblasts, 70 Erythrocyte(s) 56, 57, 59ff, abnormal

also Tissues, epithelial

Epsom salts, 346

forms of 62, counting of, 60, crena tion of, 62, 63, 68, diameters of, 59, effect of spleen on action, 97, in fetus 679, formation of, 70, increase at high altitudes, 61, life of, 71, nucleated, 62, sickle shaped, 62, stroma of, 71

Erythrodextrin, 300 Erythromycin, 222 (footnote) Erythrophores, 439 Erythropotesis, 70 "Escape of heart," 188 Eserine action of on heart, 191, 192,

action of on pupil, 608 Esophagus, 301, 304, in act of swallowing, 302

Essential hypertension, 143 Esthesiometer, 644 Estradiol, 448, 449, 450, 653, and milk secretion, 455, and onset of labor, 678 Estrin, 446

Estrogen, 450 (footnote)

Estrone, 449 (footnate) Estrous cycle, see under Cycle Estrus, 445, 446, 447, 457 Ethylamine, formed in large intestine,

342 Eve's method of artificial respiration,

Excitability: of cardiac muscle, 159, of nerve, 467ff, of protoplasm, 22 Exophthalmic goiter, 411, 412 Exophthalmos, 411, 412

Exothermic reaction, 350 Expiration, 236

Expiratory reserve volume, 240, 241.

Extension, 483 Exteroceptors, 512

Extractives, gastric secretion and, 313 Extrasystoles, 161, 184, in paroxysmal tachycardia, 185

Extransic factor, 75 Eye(s), 574ff, accommodation of, 97—in new born, 679, compared to camera, 992, conjugate deviation of, 606, convergence of, 997, dark adapta tion of, see under Adaptation, defects of, 608–15, development of, 668 671, effects of bad lighting on, 614, 615, far and near points of, 594–95, fixation point of, 607, light adaptation of, exe under Adaptation movements of, 605, 606, 607—and labyrioth see also Mystagmus muscles of, 605, 606, 607, pressure within, see Intraocular pressure

Eye strain bad lighting and, 615 cause of headache 614, cause of nausea, 119

Eyelid(s), movements of, 567

Fainting, 154 200

Falx cerebell: 556, 557, cerebri \$56, 557

Fasciculus(i) cuneatus 528, gracilis, 528 see also Tracts

Faits) 372–75 absorption of, 334 of blood 57—n diabetes, 370, combus tion of in calorimeter 353 and contraction of gail bladder, 333, dietary requirement of, 800 digestion of, 334, effect of on gastne secretion and motility, 314 emulsification of in in testine, 334 beat equivalent of 354 metabolism of, 372ff in milk, 620 condation of 374, 375, storage of in

body, 352, 374
Fatigue of muscle, 491, of reflex center,

515

Fatigue toxin," 572
Fatty acids see under Acid

Fear on adrenal glands, 423, effect of on respiration, 257

Feces formation of, 345, loss of heat in 378 379 pigment of, see Stercobilin Femur, 48, 49

Fenestra ovalis, see Window, oval, rotunda see Window, round Ferments, see Enzymes

Fertilization of ovum 663 Fetal circulation, 672, 673

Fetus, 675, 679

Fever, 56 basal metabolic rate in 359 effect of drugs on, 382, heart rate in,

186, loss of water in 56 Fiber(s) adrenergic, 566, 567, association 530, cholinergic, 566, 567, collagenour 45, 46—corritione and, 426, 427 commicsural, 530, 538, elastic, 45—in vascular walls, 123, mantie 26 muscle 41, 480—single, 489, nerve, 499T—functional classification 459, 460, single, 475, optic, 464, 575, 576, 577, 578, 586—absence of neu rilemma from 464 decussation of 586, single 584, pre- and post gan glionic, 562, 563, 567—of cardiae nerves, 189, projection 530, vasocon strictor, 195, vasocontor, 195, vasocontor, 195, 196—to penis, 563, vasocontor, 195, 196, 196

penis, 563, vasomotor, 195 196 Fibrils in cells 25

Fibrin 102, 103 Fibrinogen, 18, 57, 103, 361, lack of,

Fibroblasts 45, 46
Fibrocartilage 47, 48
Fick principle, 133
Fillet see Lemniscus

Filtrable viruses, 211
Filtration, through renal glomerulus 276

Final common path 513 Fish respiration in 672 Fish liver oil 393

Fish liver oil 393
Fission, binary, 27 28
Fission(s) lateral con-

Fissure(s) lateral cerebral see of Sylvius of lungs, 229, of Rolando see Central sulcus, of Sylvius 531 533, 535 superior longitudinal 529 su perior sagrital, see superior longitudinal

Fistula esophageal, 310 gastric, 310 311

Flavors see Sensation taste Fleming Sir Alexander, 221

Flexion 483 Flexures of large intestine, 322

Florey Sir H , 221
Fluid(s) of body, 52ff—reaction of

108ff, in Bowman's capsule, 278, cerebrospinal see Cerebrospinal fluid ex change of between capillaries and tis sue spaces, 53, 65-67, intraocular 616 for perfusion of heart, 162, 163,

164, 165, for transfusion 77ff Fluorides as anticoagulants 104

Fitiorine, essential in diet, 402
Flying (military) effects of on circula

tion, 155, effects of on ear 629, at high altitude, 262 Folic acid, 76 384

Follicle(s) of glands, 39, Graafian see Graafian follicle, hair 35

Food(s) absorption of, 343-45, calone values of various, 400, center con trolling intake, 349, energy from 351 heat equivalents of 334, oxidation of in body, 351, 352, requirements of findet, see Diet, planning of, specific dynamic action of, see Specific dynamic action see also Carbohydrate Fat Protein

Foot-candle, 615 Foramen of Munro, 558, 560, ovale, 673 Forced respiration, see Respiration, forced

Fovea centralis, 575, 579 Foveola(ae), of gastric glands, 305, 306 Freemartin, 665 von Frey's hairs, 644, 645

Frohlick's type of dwarf, 441, 442 Frontal lobe, 531, 532, 533, 534 535

Fructose, 328, 367, 368, diphosphate, 497 Fruit(s), as laxatives 346 FSH see Hormone, foliale sumulating

Fundus of eyeball, see Fundus oculi, of stomach, 304, 305, 307

Fundus oculi, 578 579 Fungi, 210, 211, 212 Furred tongue 342 Furuncle, 89

G's. 155

Galactogogue action of pituitrin 437

Galactose, formula of 367 368 Galen Claudius 121 Gall bladder, 331, 132, 333

Galvanometer, 178, 470 Gametes, 648

Gamma globulin, 57, 58, 91, 219 and

antibodies, 91 formed in lymph nodes, 100, in passive immunization, 219, produced by lymphocytes, 91

Gamma rays 5 Gangliated cord of sympathetic, 419 563 Ganglion(a) celiac, 565, cervical, 565, facial 549, geniculate, see facial, hypogastric, 565, impar, 563, mesenteric, 565, otic, 296, of parasympathetic, 562, of posterior spinal roots, 525, prevertebral, 563, 564, 565, spiral of Corts, see vestibular, stellate, 189, 423 (footnote), submaxillary, 296, sympathetic, 563, 564, 565, thoracic, 565, of vagus, 550, vertebral 563 565, vestibular, 631, 632

Gas(es) of atmosphere 245, diffusion of 244, 247, kinetic theory of, 244ff, laws of, 244, 245, properties of, 244, respiratory, 245, 246, 247-exchanges of, 247, 249, transport of in blood, 249, 250, 252, see also Oxygen, Car

bon dioxide, Nitrogen Gastric analysis, 313 Gastric fistula, see under Fistula

Gastric juice, see under Juice Gastric phase of secretion by stomach. 309, 312, 313

Gastrin, 313 Gastrointestinal tract (diagram), 289 Geiger counter 5

Gel, see Hydrogel

Gelatin, 17, an incomplete protein, 365, as a transfusion material, 84 Germ layers, 668, 669

Gestation, see Pregnancy, ectopic, see Pregnancy, extrauterine

Giantism, 439, 440 Gianuzzi, demilianes of, 295

Gills, 672

Gland(s), 38, 39, 40, adrenal, see Adrenal glands, apocrine, 654, crop of pigeon, 456 (footnote), ductless, see Endocrine glands, endocrine, see En docrine glands, intestinal, 326, 327, lacrimal, 574, 576, lymph, see Lymph nodes mammary, see Mammary gland, parathyroid, see Parathyroid glands, parotid, see Salivary, pineal, see Pineal body, pituitary, see Pituitary body, prostate, see Prostate, racemose, 40, 41, 295, salivary (salivatory), 39, 294, 295, 296-excretion by, 300, sex 444-57, see also Ovaries, Testes, of skin 34, of stomach, 305, 306, 307, sublingual, see Salivary, submaxillary, see Salivary, submandibular, see Salivary, sweat, 378-cholinergic nerves to, 567, thyroid, see Thyroid gland, tubular, 38, 39, types of, 38, 39

Glans penis, 658, 659 Glare, 614, 615

Glaucoma, 616 Gliadin, 361, 365

Glisson, Francis, 481, Glisson's experi ment, 480, 481

Globin, 361 in blood regeneration 72 73, a constituent of hemoglobin 68 Globulin(s), 361, serum, 57, see also Gamma globulin

Globus pallidus, 540 Glomerular tuft, 268 269, 270, 271, 272 Glomerulus of kidney, 268 269, 270

271, filtration by, 276, 277 Gluconeogenesis, 369

Glucoreceptors, 349 Glucose, 328, 367, 368, absorption of,

343, 344, 345, blood, see Blood sugar. combustion of, 353, excretion of in diabetes, 370, 371, formulas of, 367 as fuel for muscular work, 503, of liver, 368, 369, of muscle, 368, 369, oudation of, see combustion of, from protein, 363, 369

Glucose phosphate in muscle 497, 498 Glucostatic theory of hunger, 349 Glucosuria 370, 371, pituitrin and, 438

Glue, 17 Glutamine, în urine, 274

Glutelin(s), 361

Glutenin, 361, 365 Glycerm, see Glycerol Glycerol (glycerm), 372, from fat, 324, 372

Glycine, 331, 360, 364 Glycocoll, see Glycine

Glýcogen, 368, 369, in hyperthyrodism, 411, of liver, 368, 369—action of adrenalme on, 423, in muscular contraction, 497, 498, 499, of muscle, 368, 369—action of adrenalme on, 422, stores of in body, 352, synthesis from lactic acid, 498, 499, used during fast, 352

Glycogenesis, 368 Glycogenolysis, 369

Glycoproteins, 37, 38, 309, 361 Glycosuria, see Glucosuria

Glycotropic action of pituitary, 432, 435.

436 Glycuronates, 343 Goster, 406, 408, 412

Goldblatt, Dr H, 143 Golgi apparatus, 24, 25

Gonad(s), see Ovary, Testis "Goose flesh," 25

Gout, 366

Graafian follicle(s), 446, 449, 450, 452, 648, 649, hemorrhages into, 456, maturing, 649, primary, see primordial, primordial, 648, 649, rupture of, 449, 450

Granules in leucocytes, 86 Nissl, 507, 508, secreting, see zymogen, zymogen, 25, 40, 41—of pancreatic cells, 323,

Granulocytes, life of, 92 Graphic formulas, 6

Gravity effect of on circulation, 152-54, force of increased by maneuvers in flying, 155

Gray matter, of nervous system, 506, of cerebellum, 552, 553, of cerebrum, 531, 537, 539, 541, of spinal cord, 523, 524

Growth: amino acids and, 365, of bone, vitamin D and, excessive, see Giantism, nitrogen balance in, 364, and riboflavin deficiency, 389, retarded, see Crein, Dwarf, thiamin and, 387 Guanine, 366, 367

Gunne pig, in anaphylactic shock, 224 Gunn acacia, 83 Gunna 215

Gunpowder fuse, and nerve fibers, 472

Habit, and constitution 341 Habitual abortion, vitamin E and, 397 Hair(s), 35, von Frey's, see von Frey's hairs, proteins of, 500, touch sensation and, 645

Haldane, J. S., 254 Hales, Stephen, 136

Halibut oil, source of vitamin A, 396 Haploid number of chromosomes, 661

Haptins, 216 Haptophore group, 216 Hartline, H. K., 584

Harvey, William, 119, 120
Hassall s corpuscles, see under Corpuscles
Hay fever, 224

HCO<sub>2</sub>/NaHCO<sub>2</sub> ratio, 113 Headache, in consupation, 341

Headache, in consupation, 341
Hearing, 622, thresholds of, 627, 628, see
also Ear

Heart, 115ff, acceleration of, 188, 189, action of, 169ff, anatomy of, 115-19. apex beat of, 183, beat of, see Heart beat, blood supply of, 204, 205, see elso Coronary circulation, calcium rigor of, 162, cold blooded, 166, depervated, 421, diastole of, 171, 172, 173, dilatation and hypertrophy of, 177, 194-in athletic training, 505, disease of-cardiac output in, 133 compensation in, 177, dilation in, 177, and muscular exercise, 505, effect of cations K+ Ca++ and Na+ on 1, effect of drugs on, 191, effect of oxygen lack, 237, electrical changes in, 177ff, see also Electrocardiogram, of embryo, 672, escape of, from yagal inhibition, 188, excised, 162, excitation wave of, 167, extrasystoles of, 161, hypertrophy of, 177, 194, 505, inhibition of-by potassium, 163, by vagus, 188, 189, minute volume of, 132, see also Cardiae output, nerves of, see Nerves, cardiac, output of, 132, see also Cardiac output, pacemaker of, 167, perfusion of, 162ff, position of, in chest, 232, premature beats of, 161, pressure curves of, 171, 172, Purkinge's system of, 167, 168, rate of, see Heart rate, regulation of, 186ff, relation of length of muscle fibers to contractile force, 193, reserve power of, 194, sensory nerves to, 190, sounds of, see Heart sounds stroke volume of, 132, systole of, 171. 172, 173, valves of, see under Valve, valvular disease of, 176, 177-murmurs in, 177, work of, 130-in muscular exercise, 505 Heart beat, 132, 165ff, conduction of,

165, disorders of 183ff, origin of, 165 Heart block, 169, 183, 184, 186, caused by digitalis, 191, in auricular flutter, 185

"Heart burn," 315

Heart failure, 194, anoxia in, 262, oxygen administration, 264 Heart rate, 186, action of adrenaline on, 421, effect of vagus on, 187, 188, in

fetus, 679, in hyperthyroidism, 411, in muscular exercise, 503, 504, in

sleep, 572, in tetany, 415 Heart sounds, 176, 177

"Heat." 445

Heat balance, 375, daily loss of, 378, dissipation of, from body, 376, 377, 378, evolved in combustion of differ ent foods, see Heat equivalent, equivalents, 354, latent, of evaporation, 377 (footnote), produced by body, see Metabolism, radiation, and convection from body, 377, 378, sensation of, 644 Heat centers, see under Center

Heat cramps, 383

Heat exhaustion, 383

Heat production by muscle, 496, 497, 501, by nerve, 478, in sleep, 571, see

also Metabolism Heat rigor of muscle, 494

Heatstroke, 383 Hematin, 69

Hematinic principle, 75 Hematocrit, 63, 64

Heme, 69, 71

Hemianopia, 586, 587 Hemiplegia, 539

Hemisphere(s) cerebellar, 551, 552,

cerebral, 529ff Hemocoma, 71

Hemocytoblast, 51

Hemocytometer, 61 Hemodynamic principles, 126ff

Hemoglobin, 60, 68ff, affinity for carbon monoxide, 262, bile pigment from, 329, 335, as a carrier of carbon dioxide, see Carbohemoglobin, concentration of in blood. 70, in cyanide poisoning, 26, in cyanosis, 263, in muscular exercise, 504-05, oxygen dissociation curve of, 249-51-in fetus, 674, oxygen saturation of, 249, 261, 262, oxy genation of, 69, reduced, 69, 263, as

a weak acid, 111

Hemolysins, 85, 215 Hemolysis, 84, and raundice, 335

Hemolytic disease of new born 82 Hemophilia 107, 108

Hemopotesis, 70, copper essential for,

Hemorrhage 76 77, action of "buffer" nerves in, 200, after childbirth, 680, cerebral, see Cerebral hemorrhage, due to low prothrombin level of blood, 105, 107, heart rate in, 186, in new born, 107, 397, in obstructive jaundice, 107, shock caused by, 222, transfu sion in. 77, 82

Hemorrhagic shock, 223 Hemorrhagic states, 107 Hemostasis, 106 Hemostatics, 106 Hench, P S . 427 Henle's loop, 270 281

Heparin, 104 Hepatic circulation, 207, 208

Heptoses, 367 Hermaphrodite, 665 Hermaphroditism, 665, 666

Hexoses, 367, 369 Hibernation, periodic breathing in, 260 High altitude, 261, 262, erythrocyte

count at, 61, 62, red bone marrow at, 62, respiration at, 255 High-threshold substances 280

H ton concentration, see Hydrogen ton concentration

Hirschsprung's disease, 340 Hirudin, 104

Histamine in anaphylaxis, 224, in gastric mucosa, 313, in leucocytes, 93

Histidine, histamine from, 224, 342 Histones, 361 Histotoxic anoxia 261, 263 "Hives," 106

Holger Nielsen method of artificial res piration, 238 Holmes, Oliver Wendell, 211, 214

Homeostasis, 53, 220 Homosothermic animals, 375

Hopkins, Sir F Gowland, 383 Hordein, 361, 365

Hormone(s), 403ff, of adrenal cortex and renal function, 282, adrenocorti cotrophic, 428, 432, 435-and produc tion of plasma globulin, 101, effect of on eosinophil leucocytes, 90 exces sive secretion, 444, anterior pituitary like 456, 457, antidiuretic, 56, 277, 281, 283, 438, 444, 538, antidiabetic, see Insulin, chemical nature of, 403, and contraction of gall bladder, see Cholecystokinin, follicle-stimulating 455, 649, follicular, 449, see also Estradiol, galactogogue, 438 gall blad

der and, see Cholecystokinin, gastric,

313, gonadotrophic, 432, 433, 441, 452-455, growth, 432, 433, 434, 440,

and heat regulation, 380, interstitial

cell-stimulating, 454, lactogenic, 456

(footnote), see also Prolactin, lutein-

stimulating 452 453 455 futeinizing vee lutein stimulating luteotrophic 453 (footnote) male 451 452 see also Testosterone and mammary de velopment, 653 654 of neurohypoph 437ff ovarian 449 450 also Estradiol, Progesterone pituitary 404 of pituitary body 432ff placental 456 and renal function 281 secretion of milk see Prolactin sex 373 404-in adrenal cortex 428 429 see also Estradiol Progesterone Tes tosterone of testis see Testosterone thyroid 404-in treatment of hypo-408 409 410 thyroidism trophic 404 432 434 435 441-fac tors controlling secretion of 413 word coined 325 (footnote) Horns of spinal gray matter 523 524 Hot flushes " 449 Howship's lacunae 49 Humeral transmission of nervous ef fects 191 192 Hunger 347 348 contractions panes 348 Hyaluronic acid see under Acid Hyaluronidase 51 in semen 657 Hydrocephalus 560 561 Hydrogel 18 Hydrogen 1 Hydrogen ion concentration 108 blood and body fluids 108ff effect of on contraction of muscle 490 ef fect of on enzyme action 291 292 effect of on oxygen dissociation curve of hemoglobin 251 of intestinal con tents 345 see also pH Hydrophobia see Rabies Hydrosol 18 Hydroxide(s) 9 12 13 14 17 Hydroxy 11 dehydrocorticosterone 426 427 Hyord bone 265 Hypercalcemia caused by excessive doses of viosterol 393 due to excess of parathyroid hormone 417 Hyperchlorhydria 314 315 Hyperglycemia 370 371 in diabetes 370 after pancreasectomy 434 pitus tary and 435 438 Hyperinsulinism 371 Hypermetropia 611 612 Hyperparathyroidism 417 418 Hyperpituitansm 440 441 Hyperpnea 259 Hypertensin 144 Hypertension see Arterial hypertension Hyperthyroidism 411 412 basal meta

bole rale in 359 cardiae output in

132 dyspnea in 260 heart rate in 186 Hypertrophy of heart 177, 194 Hypocalcemia 415 Hypoglycemia 371 after hypophysec tomy, 434 436 in Simmond's disease 413 Hypophysectomy 432 433 436 453 454 effect on uterus 453 Hypophysis cerebri see Pituitary body Hypopituitarism 441 442 Hypoprothrombinemia 107 Hypothalamus 541 control of body tem perature and 380 diabetes insinidus and 444 hunger and 349 sleep and Hypothyroidism 408 409 410 bisal metabolic rate in 359 cardiae out put in 133 Hypoxia 261 ICSH see Hormones interstitual cell stimulating Icterus see Jaundice Idiocy due to thyroid deficiency 410 Heocolic valve see under Valve fleum 321 339 Image(s) after 604 605 retinal 498 592 593 598-fusion of dissimilar by brain 498 Imbeculity due to thyroid deficiency 410 Immune body 215 217 Immunity 215 216 217 Impregnation of ovum see Fertilization Impulse(s) of heart see Heart beat nerve see Nerve impulse Inclusion bodies, 24 25 Incompetence of heart valves 177 Incus 617 618 622 Indoxyl sulfuric acid 343 Inductorium 484 485 486 Infantile paralysis see anterior pol omye Infection 89 90 92 93 210ff puer peral 680 Infectious hepatitis 219 Inflammation 92 93 Influenza 210 Infundibulum of pituitary 429 430 431 Infundin 437 Infusion 77 Initial heat in musculir contraction 501 502 Inoculation 218 Inositol 384 390 396 Insensible perspiration 377 Inspiration 234 235 and smell no of swallowing 302 Inspiratory expansity 240 24

```
Inspiratory reserve volume, 241 (foot-
  note)
Insulin, 370, 371
Insulin shock, see under Shock
Intercellular cement, see Intercellular
  substance
Intercellular substance, 50, 51
Intermedia, 440
Internal capsule, 533, 537, 538, 539, 540
Interoceptors, 512
         a-c, see a c interval, achro-
Interval
  matic, 581, colorless, see achromatic,
  P-R. see P-R interval
Intestinal intoxication, 342
Intestinal phase of gastric secretion, 309,
```

Intestine(s) absorption from, 343, effect of adrenaline on, 421, effect of pitu itrin on, 439, evacuation of, 340, 341, large, 320, 321, nerves to, 563, 566, small, 320

Intraocular pressure, 616 Intraventricular block, see under Block Intrinsic factor of Castle, 75 Invert sugar, 368 (footnote) Involution, of uterus 680 Iodine, 1, essential of diet, 401, isotope

of, 6 lack of, as cause of gotter, 406, in soil, 407, sources of, 407, test for starch and dextrins 299 thyroid affinity for, 414, in thyroid hormone, 408, 412 lodized salt, 401 Iodopsin, 583

Ion(s), 13, bicarbonate 13, 112, chloride, 14, 112, hydrochloride, 14, hydrogen, see Hydrogen ion, nitrate, 13, phosphate, 14, sodium, 14, sulfate, 13, 14. table of, 14 Ionization, 12, 13, of acids and bases (Table), 13

Iris 575, 607-608, see also Pupil Iron deficiency of, 73, in hemopoiesis, isotopes of, 5, radioactive, in measure ment of blood volume 64 requirement of, in diet, 401, in tirine, 274 Irritability of protoplasm, 22

Isinglass, as a transfusion material, 84 Islets of Langerhans 323, 370, effect of adenohypophysis on, 436 Isoelectric precipitation of protein, 361 Isoelectric range, 361 Isotopes, 4, 5, radioactive, 5 Isotropic bands, 44 Ivy, A C, 333

I bands of striated muscle, vee I discs J discs 41, 42 Jaundice 335 failure of absorption of

vitamin K in, 107, 397, hemorrhage in, 107, 397 Jejunum, 320 Jenner, Edward, 218 Juice: appetite, 310, gastric, 307, 308, 309-secretion of, 307, 308, 309, 310, intestinal, 326, 327, 328-enzymes of, 328, secretion of, 325, 328, pancreatic, 323-26-secretion of, 325

Junctional tissues of heart, 165 Karyoplasm, see Caryoplasm Kendall's compound E, see Cortisone Keratin, 33, 361 Keratomalacia, 385, 386 Ketogenic action of pituitary, 432, 436 Ketone bodies, 370, 371, 374 Ketonuria, 370 Ketosis, 370, 371 Kidney(s), 122ff, action of hormones on.

281, in control of acid base balance of blood, 110, disease of, 283, 284and hypertension, 143, 144, 384, in fetus and new born, 679, as food in blood regeneration, 72, functions of, 283, physiological anatomy of, 269ff, structure of, 269ff, and surgical shock, 274, water of, 55

Kinetic theory, 19, of gases, 243, 244 Kite, in experiments on digestion, 317 Knee jerk, 511, 518, 519 Krause's end bulbs, 645 Krause's membrane in muscle fibers, 41, Kymograph, 137, 138, 484, 485 Kyphosis, in rickets, 394

Labor (parturition), 675, 676, 677, 678. factors in onset of, 678 Labyrinth of ear, 618, 619, 630, non auditory, 630ff reflexes from, 634ff Lactalbumin, 361, 365 Lactase, 327 Lactation, see Milk, secretion of Lacteal(s), 99, 344 Lactic acid, maximum in muscle, 498 see also under Acid Lactose, 367, in milk, 654 "Laking" of blood, see Hemolysis Lamina propria see Membrana propria Langerhans, islets of, 323 Laryngeal spasm in tetany, 415, 417

Laughing, 257, 259 Lavoisier, Antoine, 351 Law(s). "all or none"-of cardiac muscle, 159, 160, of striated muscle 488, of nerve, 471, 472, of biogenesis,

Larynx, 226, 227, 264, 265, 266, 301,

in swallowing, 302

see of recapitulation, Boyle's, 244, Charles', 244, of conservation of rergy, 357, Dallons 245, of gases, 244, 245, Gay Lussae's, see Charles', of bemodynamus, 126, 127, 128, Henry's, 190, of mass action, 291 (footonet), Mendelian in inheritance of blood groups, 30, of partial pressures see Dallons of recapitulation, 670 671, 672, Startings, of heart, 193, Van Hoffs, 492

Laxatives, 346
Layer(s) of cerebral cortex, 531, 532, of retina, 576 577, of skin, 33, 34
"Lead pipe" rigidity, 541

Lecutini 373, in bile, 329, in erythrocytes, 59 Legumin, 365

Lemniscus lateral, 542, 545, medial 545, 546, trigeminal 546

Lens of eye, see Crystalline lens, glass 590, 591—achromatic, 610, contact, 614

Letting down" of milk, 438, 455 Lettuce, as source of vitamin E, 395 Leucemia, see Leukemia

Leucocytes 85ff basophil, see granular in cerebrospinal fluid, 561, counting of, 60° differential count of, 91, cosmophil, see granular functions of, 87, 88, granular 86, 87, 88, histamine in 93, 224, in inflammation, 92–93, life of, 92, numbers of, 87, 91, neutrophil see

granular polymorphonuclear, see gran ular Leucocytosis 91 promoting factor, 92 Leucopenia, 92

Leucoiaxin 93 Leukemia, 92 Levulose, see Fructose Leydig, cells of, 452, 454

Light adaptation of eye to 581, 582 effect of on pupil, 608, red rods of retina insensitive to, 581, refraction of

reter to the pupil, 608, ret rods of retina insensitive to, 581, refraction of 589, 590, 591, 602, 603, sensitivity of retina to, 583, see also Visual actus, split by prism, 581, ultraviolet, anti-rachitic action of, 392 393—effect of on skin, 393, visual purple bleached by, 582, 583, wavelengths of 599, 600

Lillie, Ralph, 472 Lillie's iron wire model of nerve, 472-

73 Lipase(s), 292, 293, 327, of gastric puce, 307, 309, of intestinal puce, 328,

pancreatic, see Steapsin Lipemia, 370 Lipoproteins, 361, in atheroscierosis, 145 Liquid paraffin as a laxative, 346 Liquor amnii 675

Liquor folliculi, 649, 650 Lister, Lord Joseph, 211, 213

Lister, Lord Joseph, 211, 213
Litmus 15
Liver, 289, and carbohydrafe metabolism
369, circulation of blood in, 207, 208
cords of, 208 328, detoxicating func
tion of, 342, 343, and fat metabolism
373, glycopen of, 368 origin of from

cords of, 208 328, defoxicating function of, 342, 343, and fat metabolism 373, glycogen of, 368 origin of from entoderm, 669, production of bile by 328 in regeneration of blood (Table) 72, sinusoids of, 208, as a source of serum albumin and fibrinogen 57, water of, 55

Liver extract in the treatment of per nicious anemia 74

"Living test tube," 105

Lobe(s) of cerebrum 531-38 frontal 544, of lungs, 229; neural of pituriary body, see Neurohypophysis occipital 533, 536, parietal 533, 535, of pitul tary body, 430 431, temporal, 533,

535 536, 544 Local excitatory state, 465

Locomotor staxia, 521 Loewi Otto, 191

Longsightedness, see Hypometropia Lorain type of dwarf, 441, 442 Lovén reflex 151

LSH, see Hormone Jutein stimulating Ludwig Carl, 137

Lumbar puncture 558, 561 Lung(s), 227, 228, 229, air of, 240-42 buoyancy of, 241, dead space of, 242, elasticity of, 232, evaporation of wa ter from, 377, 378, exchange of gases in 247, 248, expansion of 233, hyper ventilation of, see Respiration, forced,

stiffness of, 206 Lung capacity, 241

Lung unit, 229, 231 Lutern, 450, 469

Lutein-stimulating hormone see under Hormone

Lye, 14 Lymph, formation and composition of,

98 Lymph glands, see Lymph nodes Lymph nodes, 99, 130 220, as filters, 100 functions of, 100, 101

Lymphatics 98, 99, 100
Lymphocytes 85, 90, 91, 98, in lymph
98, produced by spleen 97, produced
by thymus 457, production of gamma
globulin by, 101, production of plasma
globulin by, 57, 91, 101

Lymphocytous 91 Lymphopenia, 92

Lysine, 360, 365 Lysozyme, 575

Macrocytes, in permicious anemia, 74 Macrophages, 94, 95, 97 Macula lutea, 579

Magendie, François, 319

Magnesium in blood, 57, 58, in diet, 401, in inositol, 390, in urine, 274

Magnesium sulfate, see Epsom salts Maize pellagra and, 390, proteins of,

Make shocks, 465, 466, 467 Malaria, 210, 212

Malleus, 617, 618, 622, 623

Malpighi, Marcello, 202, 276 Malpighian corpuscle, see Glomerulus

of kidney

Maltase, 300, 327

Maltose, 300, 368, hydrolysis of 289 Mammary gland, 455, 652, development of, 653, secretion of, see Milk, secre-

tion of

Manganese, as an essential of diet, 402 Manometer, 137, 138, 139, 141, 151,

152, for measuring venous pressure, 152

Masculinization, 429, 665

Mass movement, see under Movement Mastication, 301

Mating season, 445, 447

Measles, 210, 219 Meat calorie value of, per pound, 400, proteins of, 365, minerals of, 401

Meatus(es) external auditory, 617, in ternal auditory, 620, 621, 622, of nose,

226, 227 Meconium, 679

Mediastinum, 231, 232

Medulla of adrenal, see Adrenal medulla, of kidney, 273, of thymus, 457 Medulla oblongata, 529, 531, 543, 546,

Megacaryocytes, 93, 94

Megacolon, 340

Megaloblasts, 70

Meiosis, 661, 662 Meissner's corpuscles, see under Cor

puscles

Melanin of skin, 36 Melanoblasts, 36 Melanophore expanding action of pitui-

trin, 437, 438 439 Melanophores 437, 438

Membrana granulosa 649, 650 Membrana propria, 29

Membrane(s) basement 29, basilar, 619, 620 622, 624, 625, 626, of brain and spinal cord, 555-57, cell, 25, drum, see tympanic, fetal, 676, 677, 679, Krause's, in muscle fiber, 41, 42, mucous, 37-of intestine, 326, of stomath, 304, 305, 306, of red cell, as semipermeable membrane. Reissner's, see tectorial, semipermeable, see Semipermeable membrane, tectorial, 620, tympanic, 617, 618, 622,

Membrane theory of nerve conduction, 472

Memory, 536

Meninges, see membranes, of brain and spinal cord Meningitis, 561

Menopause, 449

Menstruation 447, 448, 450 precocious 429, suppression of, in pituitary disease, 443

Mercury, diuretic effect of, 282 Mercury vapor lamp, 393

Mesencephalon see Midbrain Mesoderm, 668 669

Mesothelium, 38 Metabolism,

350-70. carbohydrate. 367ff-in diabetes, 370 hormones and 371, insulin and 370ff, and nerv ous system, 372, dyspnea caused by increase, 260, effect of fever on 381, fat, 372ff-in diabetes, 370 of nerve 478, protein 359-65 of protoplasm,

7, 23, purine, 366 367 Metalloporphyrins 68 Metaphase, 26 27 Metaprotein, 361 Metarterioles 202, 203

Micellae, of muscle, 42, 497, 500 Microglia, 506, 507

Microgram, 401 (footnote) Micromicron, 17 (footnote) Micron, 17 (footnote)

Microphages, 95 Microvolt, 537 (footnote) Micturition, 285ff

Midbrain, 529, 531, 543ff

Midget, 441, 442 Midget ganglion cells of retina, 576, 577

Milieu interne, 54 Milk calcium of, 407, calorie value of,

400, coagulation of, by rennin, 309, composition of, 654, irradiated 393 "letting down" of, 438, 455, proteins

of, 363, secretion of, 454 653, 654 680-hormones and, 454 455, pitui tary and, 432, 435, 454, vitamins of, 654, 655, "witch's" 456 (footnote)

Milk sugar, see Lactose Millimeter, 17 (footnote) Muscle twitch phases of 487 recording of 484-87
Muscular contraction 494ff chemistry of 497ff—phases in 498 499 503

of 497ff—phases in 498 499 503 and intensity of simulus 488 dys trophies vitamin E and 395 effect of loading on 490 491 effect of stretching fibers on 490 491 effect of temperature on 492 efficiency of 495 496 497 heat production in 501 502 and intensity of stimulus 488 isometric 491 496 inotonic 491 496 latent period of 487 488 phases of 497 498 499 source of energy for 497 499 503 total energy expendi

ture of 497
Muscular exercise 502 05 dangers of in elderly 505 dyspinea in 260 eryth rocyte increase in 61 heart rate in 186 and heat production 379 380 metabolism in 339 physiological ad

justments in 103 504 Muscular work 495 carbohydrate usage

Mustard as an emetic 319 Mustard 460

Myelin sheath 460 461 481 487 508 Myelinization 461

Myeloblasts 92 Myelocytes 92

ſ

Myocardium see Muscle cardiac Myofibrals 41

Myoglobin 497 Myoneural junction fatigue of 487 492 Myopia 611 612

Myzedema 410 411

Nails 35

Nasal conchine 301 641 Nature healing power of 219 Nauvea 310 and 1 byrnith 634 Navel rer Umb licus Near au-hitedness are Myopia Necrous of heart muscle 205 Needling of lens 615 Negatine exponent 109 Negroes Rh positive blood 81 (footnote) Nephritis 283 284 Nephron 269 270 Nerwe(3) abducent 544 547 548 549 accelerator 188 189 accessory 544

Neinton 269 270
Nerve(s) abducent 544 547 548 549
accelerator 188 189 accessory 544
547 551 accommodation of 467
acoustic 544 547 549 afferent 460
all or nibne Jaw of 471 472 maye
linated 460 aortic 199 200 audi
torv fails to regenerate 464 aug
mentor to heart see accelerator of
blood vessels 195 buffer 200 car

diac 186 189 190-hberation of acetylcholine and adrenaline by 192 of cerebral vessels 209 chemical changes in 478 chorda tympani 296 297 549 567 618 640 chronaxie of 467-69 of coronary vessels cranial 547-51 degeneration of 467 525 depolarization of 473 474 ef ferent 460 electrical changes in 469 71 electrotonus in 465 67 excitability and conductivity of 464 465 of face 549 facint 297 544 547 549 of gall bladder 332 glossopharyngeil 199 200 296 544 547 548 549 550 640-in act of swallowing 303 hypoglossal 544 547 551-in act of swallowing 303 of large intestine 340 lingual 296 640 mandibular 549 mixillary 549 medullated see myelinated mixed 459 motor and sensory 459 400 of muscle 482 myel nated 400 myelinization of 461 nonmedullated see amyelinated ocu lomotor 542 offactory 227 643 oph thalmie 549 optie 547 574 575absence of neur lemma 4f1 fails to regenerate 464 parasympathet c 567 565-to sal vary glands 296 297 vasodilator fibers of 190 see also pelvic variis etc pelvic 563 565io bladder 287 of penis 659 phren c 253 256 [lexuses (cervical brach il etc.) 523 polarization of 473 of pulmonary vessels 207 re fractory periods of 473 474 regen eration of 461 464 renal 272 273 sensory to heart 190 s nus 190 199 200 201 of small intestine 338 339 spinal 524 525 526-primary d vi sions of 575 526 splanchnic 195 564 stimulation of 465 466 stomach 315 316 strength-duration curve of 468 structure of 459ff of swallowing 303 sympathetic 209and act of ejaculation 657 to adrenal medulla 424 to bladder 287 to blood vessels 195 to heart see accelerator to salivary glands 296 297 to small intestine 338 339 vasoconstrictor 195 196 565 vasodilator 196 565 vasomotor see vasoconstrictor vasodiator of taste 638 639 640 tri geminal 544 547 548-in act of swallow ng 303 trochlear 547 twoway conduct on in 515 570 521 vagus 544 547 150 562-acetyl choline from in heart 191 in act of swillowing 303 action of on auriculoventricular conduction 188 in card o

spasm, 303, to heart, 187, 188, 191, and pancreatic secretion, 325, to intestine, 338, 339, vasconstrictor, 195, 196, 565—distribution of, 196, vascodifator, 196, 565, vestibular, 631, 632, of vomiting, 319

Nerve impulse(s), 465ff, afferent, in relation to sleep, 572, 573, antidromic visodilator, 196, frequency of, 474, 475, 476, 477, 478—in auditory nerve, 624, 628, and magnitude of response, 476, 477, 478, velocity of, 474, 475 Nervus intermedius 549 Nervus intermedius 549

Nervus intermedius 549 Neural stalk of pituitary body 429 Neurilemma, 461, 463, 464, 508 Neurofibrils, 462, 464, 506, 507, 508 Neurofila, 506, 507

Neurohumors, 567

Neurohypophysis, 430, 431, 437 Neuromuscular junction, see Myoneural

junction Neuron(s), 506, 507, 508, effector, receptor, and connector, 510, 511, 512, intercalated, see connector; internun

cial, see connector; motor, 512, upper and lower motor, 527 Neutral fats, 372 Neutron(s), 3, 4 Neutropenia, 92

Neutrophilia, 97 Neutrophilis 87, 88, 89, 91

New born, 679, erythtrocytes in, 679, hemolytic disease of, 82—hemorrhage of 105, 107, 397, 679, vitamin K de

ficiency in 679
Niacin see Nicotinic acid
Nicol prism, 43

Nicotine, action of, on heart, 191 Nicotinic acid, 284, 289, 296 Night blindness, 386, 583 Nissl bodies, see under Body

Nitrates in angina pectoris, 205, cause of cyanosis, 263 Nitrogen in air, 245, 246, equivalent weight of, 9, in feces, 345 ment in

weight of, 9, in feces, 343 men in respiration 245, in urine, 274 Nitrogen balance, see Nitrogen equilibrium

Nitrogen equilibrium, 363, 364

Nitroglycerine, in angina pectoris, 205 Nocturnal entiresis, 287 Nodal tissues, see under Tissue

Node(s) aunculocentricular, 167, 168, 169, 186 188—effect of vagus on, 188, lymph, see Lymph nodes, of Ran-vier, 461, 462 annoauricular 167, 168,

Nonthreshold substances, 280

18, 191. Noradrenaline, 420, 423, 425, excessive to inproduction of, 425 Norepinephrine, 420, see also Noradrenaline aline 11, 632, Normoblasts, 63, 70

Normoblasts, 63, 70 Nostrils, 226

Nuclear sap, 24 Nuclear satellite, see Sex chromatin

Nucleon, 366 Nucleolus, 24, 25, 27

Nucleoprotein, 24, 361, 366, 367 Nucleosidase, 327 Nucleosides, 366, 367

Nucleotidase, 327 Nucleotides, 366, 367

Nucleus(1): 23, 24, cerebellar, 525, 240, of cell, 23, 24, cerebellar, 525, 26, definition of (footnote); 340, dentate, see rebellar, dorsal of vagus, 550, emboliforms, see cerebellar, fastigu, see cerebellar, gracults, 228, hypothalamic, 430, 541, lentiform, 538, 540, of neu nlemma, 461, of oculomotr nerve, 542, paraventrular, 541, pontine, 542, paraventrular, 541, pontine, 546, red, 542, 545, 554, ruber, see

red, salivatory (salivary), 296, 549,

550, 640, supraoptic, 541, of trigem-

inal, 549, vēstibūlar, 522, 523, 553, 554, 632 Nutrition, 351ff Nyctalogia, see Night blindness

Nystagmus, 634

Obesity, 352, 374, 375, dinitrophenol in 383, in pituitary disease, 441, 442, 443 Occipatal lobe, 536 Occupations, calorie requirements, 399 Ocular accommodation, 593-95 Ocular hypertension, see Glaucoma

Odors, 641, classification of, 642 Oestrin, see Estrin Oestrus, see Estrus Olfactory nerves, see under Nerves

Olfactory nerves, see under Nor Olfactory sensations, 640 Oligodendroglia, 506, 507 Onion, chromosomes of, 27

Oocyte(s), 468, 661, 662 Oögenesis, 661 Oogonia, 661 Oosperm 663, see also Zygote

Ophthalmoscope, 578 Optic chiasma, 430, 531, 541, 558, 586, 587

Optic papilla, 578
Optic radiation 589
Optic thalamus see Thalamus
Optical defects, 608-16
Optical illusions 605, 606

Optical principles, 589
Optiogram, 583
Organ of Corti, 619, 620, 625, degeneration of, 625
Organdles, 24
Organods 24
Organods 25
Osmoties, 56, 66, 67
Osmotie pressure, 65, of plasma 67
Osmotie work of kidney, 281 (footnote)
Ostetitis fibroar cystica, 417, 418
Osteoclasts, 49
Osteomalcan, 395, tetany in, 416
Otcoman, see Otolith
Otoliths 631, 6335

Ovalocytes, 62
Ovarian pregnancy, 658
Ovariectomy, 445
Ovary(tes), 648, 649, action of gonadotrophic hormones on 455, effect of pituitary extract on 453, hormones of, see under Hormones pregnancy on,

see Ovarian pregnancy
Oviducts, 650
Ovovitelline, 365
Ovulation, 446, 447, 448 450 649

Ovalbumın, 361, 365

Ovum(a), 449, 450, 648, fertilization of, 451, 650, 657, 658 662, 663, implantation of, 669, impregnation of, see fertilization of see also Blastocyst

Oxalates as anticoagulants, 104
Oxidase(s), 292

Oxidation, 9
Oxidative phase of muscular contraction, see Aerobic phase

Oxygen absorption of, in respiration 243, in alveolar nir, 246, in arterial blood, reduced pressure of, 255, atomic weight and number of, 2, capacity of blood, 70, consumption, 354, 355, 359 -basal metabolic rate from, 356, 357, of in combustion of food stuffs 355, of in muscular exercise, see Oxygen debt, of by nerve, 478, effect of in fatigue of muscle, 491, 492, equivalent weight of, 9, in expired air, 246 in inspired air, 245, 246, lack of, see Anoxia isotope of 5, in measure ment of cardiac output, 133, partial pressure of-at high altitudes 261, in arterial blood, 248, in venous blood 248, in plasma 249, in saliva, 298, want of, see Anoxia

Oxygen administration 264, in artificial respiration, 240, 264
Oxygen capacity of blood, see under

Oxygen debt, 502, 503

Oxyhemoglobin, 69, 250 Oxytetracycline, 222 (footnote) Oxytocic action of pituitrin, 437 Oxytocin, see Pitocin

Oxygen transport, 249, 250

249, 250

Oxytocin, see Pitocin
Pacemaker, of heart, 167

Pacinian corpuscles, see under Corpuscles Pain in angina pectoris, 205, cerebral cortex and, 535, cutaneous, 644, 645, fast and slow conduction of 475 of

Oxygen dissociation curve of hemoglobin

fast and slow conduction of, 475, of hunger, 348, intestinal, 347, protective value of, 219, 220, referred, 347, thalamus and, 439, 540

Palate, in swallowing, 301, 302
Pancreas, 289, 322, 323, action of pituitary on, 432, 436, 455, and diabetes, 370, removal of, 370

Pancreatectomy, 370, 436
Pancreatic diastase, see Amylopsin
Pancreozymin, 325
Panting, 378

Pantothenic acid, 390, 396
Papillae of skin, 33, 34, of tongue, 638
Papillary muscles of heart 117

Para minobenzoic acid 221, 384, 390, 396 Paracasein, 309 Paraffin oil 346

Paraganglia, 419
Paralax, 599
Paralysis, caused by cerebral hemorrhage, 539, flaccid, 525, 527, spastic, 525.

527, upper and lower neuron types, 527 Paralysis agitans, 541

Parasites, 212, eosinophil leucocytes and,

Parasympathetic nerves, see under Nerve Parathyroid extract, 416, and vitamin D 393

Parathyroid glands, 414ff
Parathyrotrophic action of pituitary, 432,

Parietal lobe, 535 Paroxysmal tachycardia, 185

Pars distalis of hypophysis, see Adenohypophysis

Pars glandularis of pituitary, see Adeno-

Pars intermedia of pituitary, 430, 431, 440
Pars nervosa, see Neurohypophysis

Pars tuberalis 430, 431
Parthenogenesis, 664
Particle(s) alpha, 5 colloid 17

Pasteur, Louis, 211, 213 of trypsin, 324, scale, 110 Pathogens, 210ff Phagocytosis, 88 89, 90 Payloy, Ivan. 310, 567, 568, 573 Pharyngeal pouches, in embryo 671, 672 Pharynx, 226, 301, muscles of, 550, sen Peas, proteins of, 365 Peduncle(s); cerebellar, 543, 544, 551, sation of thirst in, 349 554, cerebral, 529, 543 Phenylalanine, 360 Pellagra, 389, 390, 396 Phipps, James, 218 (footnote) Pelvis female organs of, see under Re-Phonation, see Voice productive organs, of kidney, 272 Phosphate(s), in perfusion fluids, 163 273, 285, male organs of, see under Phosphatides, see Phospholipids Phosphenes 578 Reproductive organs Pendular movements, see Movements, of Phosphoereatine in muscle, 497, 498 small intestine 499, in nerve, 478 Penicillin 221, 222 Phospholipids, 373, in atherosclerosis 145 Penicillum notatum 222 Penis 659 perves of, 659 Phosphoprotein(s), 361 Phosphorus in bone, 49, in diet, 401 in Pentose, 366 367 inosital 390 isotope of, 6, radioactive Pepsin, 288, 292, 307, 308 see isotope of in urine, 274, 275, vita Pepsinogen, 308 Pentidases 324 327 min D and 393 Peptids 323, 361 Photophobia, in riboflavin deficiency, 389 Peptones 307, 361 Photosynthesis, 350 Pericardial sac, 115 Physostigmine, see Eserine Pied termineaux, 509 510 Pericardium, 115 Pericaryon 506 Piezoelectric effect in cochlea, 627 Penlymph, 620, 630 Pigeon crop gland of, thiamin deficiency Perimeter 587, 588 in, 387, 388 Perimysium, 43 Pigeon breast, 394 Perincurium, 459 460 Pigment(s) of bile, 329, 330, of feces Period(s) of auricular systole, 170 171 see Stercobilin, of retina, 577, 578 172, 173, of cardiac cycle 169ff, of 583, of skin, 34-pituitrin and 437 diastasis, 170 171, 172 173, of ejec 438 439 of unne, see Urochrome tion, 170 171, 172, 173 isometric visual purple, 583 contraction, 170, 171, 172 173, 150 Pilocarpine, see Eserine metric relaxation, 170, 171, 172 173. Pincal body, 458 latent, of muscle 487, menstrual 447, Pinna 617 puerperal, see puerperium refractory Pitch of sound, 624, of voice 266 -of cardiac muscle, 160, of nerve, Pitocin 439 473, of striated muscle, see latent, re-Pitressin, 439 productive, of females, 445 Pituicytes, 431 Periosteum union of muscle tendon Pituitary body, 429ff, disorders of in with 480 man, 440-44, and onset of labor, 678 Peripheral resistance, 146, 147, and ar Pittiitary cachexia, 443 ternal hypertension 143, effect of nor Pituitary extract see Pituitrin adrenaline and adrenaline on, 423, Pituitrin 437, action of on capillaries splanchnic nerves and 147 203 in diabetes insipidus, 444

of steapsin, 324, optimum for action

Place theory of hearing see under The 337, in stomach 316 Peristaltic rush, 337 Placenta, 670, 675, 677, 679, hormones Pernicious anemia 74 gastric amacid of 456, passage of Rh antigen through ity in 314, jaundice in, 335 Perspective, 599 Plantar flexion 483 Petechiae, in scurvy, 391 Plasma see Blood plasma Peyer's patches, 101 Plasma accelerator globulin 103 plf. 108ff, of blood and body fluids Plasma clearance, Glossary 108ff of gastric june, 308, of in-testinal contents, 345 and isoelectric Plasma proteins, 57 58, and formation of urine, 277, restoration of after hemor precipitation, 361, opt mum for action thage, 77

Peristalsis 302, 341 in intestine, 336

Pleura(ac), 231 Pleurisy, 231 Plexus(es) choroid, 559, 560, nerve.

195, 525-of Auerbach, 338, 339, 562, of Meissner, 338, 339, subpapillary venous, 35, 204, 377

Plica circulares, 344 Plutonium 3 Pneuma, 120

Pneumonia, 210, 220 Poikilocytes, 63

Poikilothermia, 375 Polar bodies, see under Body

Polished rice, cause of thiamin deficiency, 386, 387

Polycythemia, cyanosis in 264 Polyestrus species, 446

Polygraph, 182 Polymorphonuclear leucocytes, 87, 88,

Polyneuritis due to thiamin deficiency, 386 388

Polypeptids, 323, 361 Polysaccharides 367

Pons, 529, 531, 538, 544, 545, 546, 554, 558

Porphytin 69 Posterior chamber of eye 575

Postestrus, 446 Postural tone, of rectum, 341

Potassium action of on heart 163, in adrenal insufficiency, 425 in blood, 57, 58, equivalent weight of, 9, sulfocyanate in saliva, 298, in urine, 274

Potassium inhibition of heart, 163 Pot belly" in rickets 393, 394 P-R interval, 182, 184

Precipitins, 216

Pregnancy, 447, 448, 674ff, abdominal. 658, Ascheim Zondek test for, 456, 457, cardiac output in, 132, dietary cal cium in, 401, ectopic, see abdominal extrauterine, see abdominal, growth of uterus in, 651, 652, injury of pitus tary in, 443, ovarian, see ectopic, Rh factor in, see Rh factor, test, see Ascheim Zondek, tubal, 658, vomit

ing in, 320 Premature beats, see Extrasystoles

Premotor area, see under Area Presbyopia 610, 611

Pressure blood, see Arterial Venous, Capillary blood presure, in Bowman's capsule, 277, 278 of cerebrospinal fluid 558, 559, filtration, in urine formation 278, hydraulic of blood see Hydrostatie, 64 hydrostatic of blood, 64, 65, 67, intra abdominal, 341, intrapleural, 232, 233, intrapulmonary, 232, intrathoracic, 232, 233effect of on blood pressure, 207, 'negative" in pleural cavities, 233, osmotic -in urine formation, 277, 278, see also Osmotic pressure, sensation of cutaneous, 644, 645, 646

Primary hypertension, 143 Prism nicol, 43, refraction by, 589,

white light split by, 581 Process(es), definition of, Glossary, of

nerve cells 506, 507, 508, 510, see alsa Dendrites Axons Proestrus 446, 447

Progesterone, 448, 450, 669, see also Progestin

Procestin, 450, 653, and milk secretion 455, and onset of labor 678 see also Progesterone

Prolactin, 432, 453 (footnote) 454, action of on crop gland of pigeons, 456 (footnote)

Prontosil, 221 Prophase 26

Proprioceptors 512, of carolid sinus 199 256, 257, 258 of lungs 257, of semicircular canals 630 631 of stri

ated muscle 518, of utricle, 638 Prostate 659, 660

Protamines, 361 with insulin, 371 Protanomaly, 602

Protanopia, 601

Protease(s) of eastric fuice see Pepsin of intestinal juice, see Erepsin leucocytes 80, of pancreatic juice see Trypsin

Protective mechanisms, 219 220

Protein(s), 359ff, as amphotenic electrolytes, 361, classification of, 361, combustion of in calorimeter, 353, con jugated, 361, deficiency of as cause of anemia, 73, derived, 361, dietary requirement, 399, 400, digestion of, 307, 323, 324, 327, elemental composition of, 359 foreign reaction to 223, 224, glucose from, 363, 369 of hair, 500, heat equivalent of 354, history of in body, 362-63, incomplete 365, metabolism of, 359-65-adrenal cortex and, 427 of plasma, 57, 58 of skin, 33, in urine, 284, in various foods, 365, of various foods, 359

Proteinase(s), see Protease Proteinates 361

Proteoses 307, 361

Protons 3, 4

Prothrombin, 103 reduction of in blood 105 107 and vitamin K, 397

Parturition, see Labor Pasteur, Louis, 211, 213 Pathogens, 210ff Pavlov, Ivan. 310, 567, 568 573 Peas, proteins of, 365 Peduncle(s) cerebellar, 543, 544 551. 554, cerebral 529, 543 Pellagra, 389, 390, 396 Pelvis female organs of, see under Re productive organs, of kidney, 272, 273, 285, male organs of, see under Reproductive organs Pendular movements, see Movements of small intestine Penicillin, 221, 222 Penicillium notatum 222 Penis, 659, nerves of, 659 Pentose, 366, 367 Pepsin, 288, 292, 307, 308 Pepsinogen, 308 Peptidases, 324, 327 Peptids 323, 361 Peptones, 307, 361 Pericardial sac 115 Pericardium, 115 Pericaryon 506 Perilymph, 620, 630 Perimeter, 587, 588 Perimysium 43 Permeurium, 459, 460 Period(s) of auricular systole, 170, 171 172, 173, of cardiac cycle 169ff of diastasis 170, 171, 172, 173, of ejection, 170 171, 172, 173, isometric contraction 170 171, 172, 173, iso metric relaxation, 170 171, 172 173. latent of muscle, 487, menstrual 447, puerperal, see puerperium refractory -of cardiac muscle, 160, of nerve, 473, of striated muscle, see latent, re productive, of females, 445 Persosteum, union of muscle tendon with, 480 Peripheral resistance, 146, 147 and ar terral hypertension, 143, effect of nor adrenaline and adrenaline on, 423 splanchnic nerves and 147 Peristalsis 302, 341 in intestine, 336 337, in stomach 316 Peristaltic rush 337 Pernicious anemia 74 gastric amacid ity in, 314 jaundice in, 335 Perspective, 599 Petechiae, in scurvy, 391 Peyer's patches, 101 pli, 108ff, of blood and body fluids 108ff, of gastrie juice 308 of intestinal contents 345, and isoelectric

precipitation, 361, optimum for action

of steapsin, 324, optimum for action of trypsin 324, scale, 110 Phagocytosis, 88, 89, 90 Pharyngeal pouches, in embryo 671, 672 Pharynx, 226, 301, muscles of, 550, sen sation of thirst in, 349 Phenylalanine, 360 Phipps, James 218 (footnote) Phonation, see Voice Phosphate(s), in perfusion fluids 163 Phosphatides, see Phospholipids Phosphenes, 578 Phosphocreatine in muscle, 497, 498. 499, in nerve, 478 Phospholipids, 373, in atherosclerosis Phosphoprotein(s), 361 Phosphorus in bone, 49, in diet, 401, in mositol 390, isotope of, 6, radioactive see isotope of, in urine, 274, 275, vita min D and, 393 Photophobia, in riboflavin deficiency, 389 Photosynthesis, 350 Physostigmine, see Eserine Pied termineaux, 509, 510 Piezoelectric effect in cochlea 627 Pigeon, crop gland of, thiamin deficiency ın 387, 388 Pigeon breast, 394 Pigment(s) of bile, 329, 330, of feces, see Stercobilin of retina 577, 578 583, of skin 34-pituitrin and 437, 438, 439 of urine, see Urochrome, visual purple 583 Priocarpine, see Eserine Pineal body, 458 Pinna, 617 Pitch of sound, 624, of voice, 266 Pitocin 439 Parressin, 439 Pituicytes 431 Pituitary body, 429ff disorders of in man, 440-44, and onset of labor, 678 Pituitary cachexia, 443 Pituitary extract, see Pituitrin Pituitrin, 437, action of on capillaries 203, in diabetes insipidus, 444 Place theory of hearing, see under The Placenta 670, 675, 677, 679, hormones of, 456, passage of Rh antigen through 81 Plantar flexion, 483 Plasma, see Blood plasma Plasma accelerator globulin 103 Plasma clearance, Glossary Plasma proteins 57, 58, and formation of urine, 277, restoration of after hemorrhage, 77

Platelets, see Thrombocytes Pleura(ae), 231 Pleurisy, 231 Plexus(es) choroid, 559, 560, nerve, 195 525—of Auerbach, 338, 339, 562, of Meissner 338 339, subpapil

Plica circulares, 344
Plutonium, 3
Pneuma, 120
Pneumonia, 210, 220
Pneumonia, 62

lary venous 35, 204, 377

Pneumonia, 210, 220 Poikilocytes, 63 Poikilothermia, 375 Polar bodies, see ur

Polar bodies, see under Body Polished rice, cause of thiamin deficiency, 386, 387

Polycythemia, cyanosis in, 264 Polyestrus species, 446 Polygraph, 182

Polymorphonuclear leucocytes, 87, 88, 89, 90

Polyneuritis, due to thiamin deficiency, 386 388

Polypeptids, 323, 361 Polysaccharides 367 Pons, 529, 531, 538, 54

Pons, 529, 531, 538, 544, 545, 546, 554, 558 Porphyrin 69

Posterior chamber of eye 575 Postestrus, 446

Postural tone of rectum 341
Potassium action of on heart 163, in adrenal insufficiency, 425, in blood

57, 58, equivalent weight of 9, sulfo cyanate in saliva 298, in urine, 274 Potassium inhibition of heart, 163

"Pot belly" in rickets, 393, 394 P-R interval, 182, 184

Precipitins, 216

Pregnancy, 447, 448, 674ff, abdomnal, 658, Ascheim Zondek test for, 456, 457, cardiac output in, 132, dietary cal cum in 401, ectopic see abdomnal, extrauterine, see abdomnal, extrauterine, see abdomnal, growth of uterus in, 651, 652, injury of pitul tary in, 443, owarian, see ectopic, Rh factor in, see Rh factor, test, see Ascheim Zondek, tubal, 658, vomit ing in, 320

Premature beats, see Extrasystoles Premotor area, see under Area Presbyopia, 610 611

Pressure blood, see Arterial, Venous Capillary blood presure in Bowman's capsule 277, 278 of cerebrospinal fluid 558, 559, filtration, in urine formation 278, hydraulic of blood see Hydrostatic, 64 hydrostatic of

blood, 64, 65, 67, intra abdominal, 341, intrapletural, 232, 233, intrapulmonary, 232, intrathoracic, 232, 233—effect of on blood pressure, 207, 'negative'' in pleural cavities, 233, osmotic—in urine formation, 277, 278, see also Osmotic pressure, sensation of cutaneous, 644, 645, 646

Primary hypertension, 143
Prism nicol, 43, refraction by, 589,
white light split by, 581

white light split by, 581 Process(es), definition of, Glossary, of nerve cells 506, 507, 508, 510, see

also Dendrites, Axons
Proestrus, 446, 447
Progesterone, 448, 450, 669, see also

Progestin, 450, 653, and milk secretion

455, and onset of labor, 678, see also Progesterone
Prolactin, 432 453 (footnote), 454, ac-

tion of on crop gland of pigeons, 456 (footnote)

Prontosil, 221 Prophase 26

Proprioceptors, 512, of carotid sinus 199, 256, 257, 258, of lungs 257, of semicircular canals 630, 631, of stri ated muscle, 518, of utricle, 638

Prostate, 659, 660 Protamines, 361, with insulin, 371

Protanomaly, 602 Protanopia, 601

Protease(s) of gastric juice, see Pepsin of intestinal juice, see Erepsin in leucocytes, 80, of pancreatic juice,

see Trypsin
Protective mechanisms 219, 220
Protein(s), 359ff, as amphoteric electro-

lytes, 361, classification of, 361, combustion of in calorimeter, 353, con jugated, 361, deficiency of as cause of anemia, 73, derived, 361, dictary requirement, 399, 400, digestion of, 307, 323, 324 327, elemental composition of, 359, foreign, reaction to 223, 224, glucose from, 363, 369, of hair, 500, heat equivalent of, 354 his tory of in body, 362-63, incomplete 365, metabolism of, 359-65—adrenal cortex and 427, of plasma 57, 58 of skin 33, in urine 284, in various foods, 365, of various foods, 359

Proteinase(s), see Protease Proteinates, 361

Proteoses 307, 361

Prothrombin, 103 reduction of in blood 105 107, and vitamin K, 397

Protons 3, 4

Protoplasm, 21ff, physiological properties of, 22, 23 Protozoa, 210, 211, 212

Provitamin(s) A, see Carotene, D, see
7 Dehydrocholesterol
Providencial (pp. 22, 88)

Pseudopodium(a), 22, 88 Pseudopregnancy, 446, 447, 451

Psittacosis, 210

Psychic phase of gastric secretion 309-12
Psychic phenomena of menopause 449
Psychoauditory area see under Area

Pteroylglutamic acid 76 Ptyalin, 288, 291, 292, 298, 300

Puberty, 445, 449, 454 Puerperal fever, 211 Puerperaum, 679, 680

Pulmonary circulation, 206 Pulmonary valves, 118

Pulse arienal, 156, 157, 158—dicrotic, 158, length of, 158 production of 156 tracing of, 158, velocity of, 156, 157,

velocity increase with age, 157, venous, 182

Pulse pressure, 138
Punctum proximum, 595
Punctum remotum, 594

Pupil, 575, action of adrenaline on, 421, action of drugs on 608, effect of accommodation on, 597, effect of light on 582, and labyrinth, 634, in sleep,

572, see also Iris Purgatives, 346

Purines 366, 367 Purkinje cells, of cerebellum, see under

Cell Purkinje phenomenon, 581 Purkinje system, 167, 168 181

Purkinje system, 107, 166 181 Purkinje Sanson images, 596 597 Purpura, 108 Pus, 89

Putamen, 540
Pylorus, 304, 305, 306
Pyramidal tracts, see Tracts.

Pyramidal tracts, see Tracts, corticospinal Pyramids of Medulla, 526, 546, decussa

Pyramids of Medulla, 526, 546, decuss tion of, 538 Pyrexia, see Fever

Pyridoxin, 384, 390 396 Pyrimidine, 366, 367

Q discs, 41, 42 Quickening, 675 Quinine, effect of on fever, 382

Rabies, 218, 301 Rachitic rosary, 393 Rachitis, see Rickets

Radiation, of heat, 376, 377, 378

Radioactivity, 5

Radium, 3, in treatment of malignancy, 6 Ramus communicans (Rami communicantes), 563, 564

Ranvier, node of, 461, 462 Rathke's pouch, 430

Rathke's pouch, 430 Rayleigh, Lord, 602

Rays astral 26, 27, from radioactive atom, 5, from ultraviolet, antiraching action of, 392

Reaction(s) anaphylactoid, 224, antigen antibody, 216, of blood, 108ff, chemical—enzymes and, 290, velocity of, 291 (footnote), of degeneration,

463, 527, of intestinal contents, 345 Reamur, René, 317

Receptors, 477, 512, 513, auditory, 620, cutaneous 644ff, gravity, 638 olfac tory, 641, 642, 643, rotary, 634, in side-chain theory of immunity, 216, of semicircular canals and utricle 634, 635, 638, taste, see Taste buds, visual, see Rods and Cones

Recipient, universal, 78 79 Reciprocal inhibition, 516, 517, of eye

muscles, 606
Reciprocal innervation, see Reciprocal inhibition

Recovery, heat in muscular contraction 501, 502

Recovery phase of muscular contraction, see Aerobic phase

Reduced hemoglobin, 69, 250

Reduction, 9
Reduction division see Meiosis

Referred pain see under Pain

Referred pain see under Fain Refeax(es) abdomnal, 1519, 527, as commodation of eye, in new born, 679, allied, 512, antagonistic, 512, ant tudinal, 635, 636, 637, arx, see Refeax, cano, 520, Pain'n, dee Re cardiac, 189, 190, center, 511, condcardiac, 189, 190, center, 511, conddept. see tendon, defectation, 241, 340 dept. see tendon, defectation, 251, 341, 518

deep, see emount contains, see, see emount contains, see, see escape and see esca

lar, 198, withdrawal 475 Reflex arc, 511, conduction over 513 Refraction of light by eye, 592, by lenses, 589, 590, 591

Refractory periods, see under Period Regeneration of nerve, 463 Regurgitation in valvular disease of the

heart, 176, 177 Reinforcement, in conditioned reflexes,

Relaxation heat of muscle, 501, 502

Renal calculi, in hyperparathyroidism, 418

Renal circulation, 272, 273

Renal clearance, see Plasma clearance Renal disease, see Kidney, disease of

Renal failure, 284

Renal function, 283

Renal glomerulus, 269, 270, 271 Renal stones, see Renal calculi

Renin, 143, 144

Rennin, of gastric juice, 307, 309 Replacement method of investigating endocrine function 404

Reproduction, 648ff Reproductive organs, female, 648-54

Reproductive organs, male, 655-60 Residual volume, 241

Respiration 225ff, abnormal 259, at high altitudes, 255, 256, chemistry of, 243, Cheyne Stokes see periodic, con trol of, 252ff-chemical, 254-257, chemoreflex, 257, reflex, 257, diffi cult, 236, see also Dyspnea, effect of on arterial blood pressure, 207, of embryo, 673, forced, 255, inhibition of during swallowing, 252, internal, 225, 226, 351, mechanics of, 231-36, movements of, 235, 236-effect of on cir culation, 153, 154, in muscular exer cise, 502 503, 504, nervous centers of, 253, 254, organs of, 226ff, periodic, 259, in unicellular organisms 52

Respiratory quotient, 354, 355, 356, 358, nonprotein, 355

Rest healing value of, 219

Restiform body, see Peduncle, cerebellar Rete testis, 655 Reticulocytes, 62, 63, 70

Reticuloendothelial system, 89, 94, 95, 215, 506, and bile pigments, 329 Retina, 574, 575, 576-579, areas sensi

tive to color, 600, 601, extrafoveal, 580, formation of image on, 592, 593, layers of, 576, 577, and release of intermedin 439, sensitivity of, 583relation of duration to intensity of light, 583

Retinene, 582 Rh factor, 80, 81, 82 Rh negative, 81

Rh positive, 81 Rheobase, 468 Rheoscopic preparation, 494 Rhesus monkey and Rh factor, 81 Rheumatism, cortisone in, 426, 427 Rheumatoid arthritis, cortisone and, 426,

Rhodopsin, see Visual purple Rhythmicity of cardiac muscle, 160 Riboflavin, 384, 389, 390, 396

Ribose, 366 367 Ribs, 231, movements of in respiration,

235, 236 Richards, A. N., 279 Rickets, 393, 395 396 Right lymphatic duct, 99

Rigidity decerebrate, 522, 523, "lead pipe," 541 Rigor mortis, 494 Ringer, Sidney, 162

Ringer's solution, 162 Ringworm, 210 Rods and cones of retina, 576, 577, 581, 582, 589

Rotation, effects of, 634 Roughage in diet, 346 Rouleaux formation, 59 Round window of ear, see under Window R Q, see Respiratory quotient Ruffini's end organs 644, 645 Rugae of stomach, 305

S-A node, see Node, sino, auricular St. Martin, Alexis, 318 Sac, lacrimal, 576 Saccule, 630, 631 Saccus endolymphaticus, 631 Sacral outflow, of parasympathetic, 562, 563

Salicylates, effect of on fever, 382 Saliva composition of, 298, daily quan tity of, 298, functions of, 298 299, 300, pH of 298, secretion of, 296-98,

specific gravity of, 298 Salivary amylase, see Ptyalin Salivary reflexes, see under Reflex Salmine, 361

Salt metabolism action of adrenal cor tex on, 426, 427

Salts, 14 Santorio, Santorio, 11, 12, 377 Sarcolemma, 41, 42

Sarcoplasm, 41, 42 Scala media, see Duct, cochlea

Scala tympani, 620, 622 Scala(ac), vestibuli, 619, 620, 622, 623 Schafer, Sharpey, 238

Schafer's method of artificial respiration,

238

Scleroprotein(s), 361

Scorbutus, see Scurvy

Scrotum, 655

Schemer, Christopher, 593, 595, 596

Sclerotic coat of eye, 574, 575

7.50

Scurvy, 391, 392, 396 Sea food source of jodine, 401 Seasickness, 319, 638 "Second wind," 503 Secondary sex characters 429, 444, 445 Secondary shock, see Shock Secretin 325 Secretion of adrenal medulia, 424 of cerebrospinal fluid, 559, gastric, 309ff -abnormal, 314, in sleep 571 572, internal, see Hormones intestinal, 328 lacrymal 574 milk 652, 653, 654 680-in vitamin A deficiency 385, pan creatic, 325 prostatic 657, reverse 281, salivary, 296 297—conditioned reflexes and 299 of seminal vesicles

656 657, sweat, 377, 378 380 383in sleep 571 of urine, 268 276-80 Secretogogues, 313 Segmentation of zygote, 666 667 Segmenting movements, see Movements of small intestine

Semen, 656, 657 Semicircular canals 630ff Seminal vesicles 451, 452 655, 656 action of testosterone on, 451, 452

Semipermeable membrane, 65 Sensation(s), sense auditory, see Hearing balance, see equilibrium cortical center for 534 535 cutaneous 644ff -cold, 644, due to vasoconstriction in skin 382 heat, 644 localization of, 645 646 pain 644, 645, 646 see also Pain pressure 644, spatial (two-point) 646 647, touch, discrimination in 644, 645 646 647, equilibrium 630, of hearing 617-30 intensity of 476. 478 kinesthetic, 528, muscle, see kin esthetic nature and quality of, 477, 478 olfactory, 640, pain, see Pain, smell, see olfactory, specificity of 477 taste, 297, 638-40-center for 640, pathway, 640 relation of to smell, 639, and saliva, 300 and thalamus 539, touch see under cutaneous, visual, see Vision Sense organs, see Receptors

interauricular, 116, interven tricular. 116, 167-defects of, 674, nasal, 226

Serum, 57, frozen 82, transfusion of 77 Serum albumin 57

Serum globulin. 57

Serum sickness 224

Sex: abnormalities of due to adrenal tumor, 428 determination of, 665, glands of, 444-57, see also Ovaries Testes, hypothalamus and 542 pre cocious development of due to adrenal tumor, 428, secondary characteristics of, see Secondary sex characteristics

Sex chromatin, 508, 666 Sex desire, 445 "Sham feeding," 310 311 "Sham rage," 542

Shell shock, see under Shock Sherman, H. C. 407

Shivering, 380 Shock anaphylactic, 222, 223, deferred see traumatic, hemorrhagic, 222, in-sulin 222 primary 222, 223 second ary, 222, 223, shell, 222, spinal, 522, 523, surgical, see traumatic, transfu-

sion, 80, traumatic, 222, 223 Shortsightedness, see Myopia "Shunts" in circulation 673

Sickle cells 62

Side-chain theory, 216 Sighing 259

Signal magnet 484 "Silent areas" of cerebral cortex 535

Silicone, and elotting of blood 105 Simmond's disease, 443 Sinus(es) venosus, 165, 164 venous,

of brain, 556

Sinusoids of liver, 208 Skeletal muscle, see Muscle, striated Skeleton origin from mesoderm, 668

Skin, 32, 33, 34, 35, appendages of, 34 35, axon reflexes in, 521, capillaries of 203, color of, 35, 36 see also Cyanosis, color in (rog, 437, 438, 439, 7-dehydrocholesterol in, 392, of different races, 36 from ectoderm 668 evapo ration of water from, 377, 378, func tions of, 36 37, glands of, 34 hairs of, 34, 35, infection of 89, 90 irra diation of, 393 loss of water from

56, papillae of, 34, temperature of, 35, vitamin D in. 37 "Skyshine" in prevention of rickets, 395 Sleep, 571-573, basal metabolic rate in

359, blood pressure, 143 effects of loss of \$71, hours required \$71, \$72 metabolism in 399 400, theories of 572, urmation in, 287 Small cell infiltration, 91

Smallpox, 210, 215 Smell, see Sensation, olfactory Snake venom, 85

Sneezing, 259 Snellen's test type 584 585

Soap, 372, formation in intestine 324 Soda lime, used to absorb carbon diox ide, 357

Sodium, loss of in adrenal insufficiency, 425

Sodium bicarbonate and acid base balance, 111, in intestinal juice, 327, in pancreatic juice, 323, in Ringer's, Locke's, and Tyrode's solutions, 165, in saliva, 298

Sodium chloride, 14, of blood, 58, formula of, 7, loss of in sweat, 383, solutions of, 163, 164, 165

Sodium glycocholate, 331

Sodium iodoacetate, poisoning of muscle by, 499

Sodium sulfate, as a laxative, 346 Sodium taurocholate 331

Sol, see Hydrosol

Solute, 15

\*\*Outuon(s), 15 68 colloidal 17, crystalloid, dextran for transfusion, 84, gelatin, for transfusion, 84, gelatin, for transfusion, 83, 84, hypertonic, 68, hypotomic 68—as a hemolytic agent, 68, 85, isunglass for transfusion 84 isotomic, 56 Locke's, 163, melar, 16 normal, 16 physiological saline, 16 Ringers 163, 164, 165, saline 16—for transfusion, 83, supersaturated,

15, Tyrode's, 163, 164, 165
Somesthetic area see under Area
Sound(s), 622, analysis of by ear, 624,
625, 626 intensity of, 627, pitch of,
624-26 waves, see under Wave

Space(s) cavernous of pens 658, 659, dead, see Dead space, subarachnoid, 556 557, 561, subdural, 556, 557 Spallanzani Lazaro, 317

Spatial discrimination in skin, 646 647 Spaying, 445 Specific dynamic action, 359, allowance

for in diet 400
Spectrum 599, 600

Speech 267, see also Vocalization Spermatids, 660 661 Spermatocytes 660 661

Spermatogenesis, 660 Spermatogonia, 660 Spermatozoon(a), 656 657, 660 662

Spermatozoon(a), 656 657, 660 662 663, life of, 657, not produced in vitamin E deficiency, 395, protamines in, 361

Spherical aberration, 609, 610
Sphincter(s) anal, 322—action of ad renaline on, 421, cardiac, 308, 318, definition of, 303 (footnote), ileocolic, 339—action of adrenaline on, 421, of Oddi, 332, 333, pupillae, see Iris, py

loric, 304, 305, 316—action of adren aline on, 421 Sphygmograph, 157

Sphygmograph, 157 Sphyngomyelin, 373 Spinal cord, 523-529, nerve roots of,

524, 525, 526, tracts of, 526, 527, 528 Spinal shock, 522, 523

Spindle in mitosis, 26, of muscle, see
Proprioceptors, of striated muscle
Spiral Green, see Contro propri

Spiral organ, see Corti's organ Spireme threads, 26

Spireme threads, 26 Spirillum(a), 212, 213

Spirochetes 212 213, 214 Spirometer 242

Spirometer, 242
Spleen 95-98, action of on blood cells
97, contractions of, 61, 97, functions
of 97, lampaged tests of 95, 07

of, 97, lymphoid tissue of, 95, 97 101, and production of lymphocytes 97

Splenectomy, 97, for purpura, 108 Spoiled clover, dicoumarin in, 104 "Spots" touch heat, and cold, etc., of

skin, 644 645
Spreading factor, 51
Sprue, tetany in, 416, 417
Squint, see Strabismus
Stagnant anoxia, 262

Stags, castration of, 445
'Staircase' phenomenon see Treppe
Stannius ligatures 165
Stapes, 617, 618, 622 fixation of, 629

Staphylococci 212 213 Starch(es), 367, digestion of by saliva 299

Starling Sir Ernest 325 Starvation, negative nitrogen balance in 363

'Steady state," 502 Steapsin, 292, 393 394 Stenosts, of heart valves, 177 Stercobilin 330

Stercobilinogen 330 Stereoscope, 598

Sterility, due to vitamin E deficiency, 395 Sternum, 231

Steroids 372, 373, of adrenal cortex, 426 Sterols, 372, irradiated, 392 Stethoscope, 176

Stimulus conditioned 569, liminal 467, maximal, 488 minimal, 488 qualities of, 467, subliminal 467, threshold, 488 Stomach, 289, 304-07, 310, 311, cancer

of, 314 digestion in 304f, early studies of functions, 317, emptying of 316, glands of, 305, 306, 307, miniature 310 311, movements of, 316–20 and permicious anemia, 75

Stomatitis, 300 Strabismus 607 Stratum (strata): corneum, 33, germi ganglia, 567-acetylcholine liberation, nativum, 33, 34, granulosum, 33, 34, 567, strychnine and, 513 lucidum, 33; mucosum, see germina-Synaptic knob, 510 tivum, of skin, 33 Syncope, see Famune Streptococcs 212, 213 Syncytium, 159 Streptomycin, 222 Syphilis, 213, 214 Stress, pituitary body in, 429 Systole, 170, 171, 172 Striate body, see under Body "Stroke," see Cerebral apoplexy Tachycardia, 186 Styptics, 106 Tadpoles skin color of and putuitary, Stroma, of ovary, 449, 469 438, effect of thyroid hormone on. Structural formulas, 6 Submucosa, 36, 37 Tail of child (footnote), of human em Subpapillary venous plexus, 35, 204 bryo, 671, 672 Substantia nigra, 542 Tartar emetic, 319 Substitution method, see Replacement Taste see under Sensation Taste buds, 638, 639 Substrate, 282 Taste pore, 638 Succus enterious, see Juice, intestinal Taurine 331 Suckling, 455 Tears, 574 Sucrase, 327 Tectum, 545 Sucrose, 327, 328, 367, 368 Tegmentum, 545 Sugar, calorie value of per pound, 400 Telereceptors, 514 Sulcus(1), of cerebral cortex, 529, 531, Telophase, 26, 27 532, central, 529, 531 Temperature, body, see Body tempera-"Sulfa" drugs, 221 ture, critical, 379, effect of on enzyme Sulfacetamide, 221 action, 291, effect of on muscular con Sulfadiazine, 221 traction, 492, effect of on oxygen dis-Sulfanilamide, 221 sociation curve of hemoglobin, 251, Sulfapyriadine, 221 environmental-effect of on body temperature, 378, 379, effect of on Sulfates, as anticoagulants, 104 Sulfathiazole, 221 cardiac output, 132, effect of on me Sulfur atomic weight and number of, tabolism, 359 3, in urine, 274, in protein, 359 Temporal lobe, 535, 536 Surrenation absence of in cardiac mus-Tendo Achillis, 484, 519 cle, 160, in reflex center, 516, in stri-Tendon(s), 479, 480, central of dia a ed muscle, 488, 489 phragm, 234, reflexes, see under Re-Surlight, antirachitic action of, 392 flex Sunshine, in preventing rickets, 394, 395 Tentorium cerebelli, 556, 557 Sunstroke, 383 Terramycin, 222 Test(s): Aschem Zondek, 456, 457, of Suprarenal capsules, see Adrenal glands Surface area of body, 158 color vision, 603, fragility, 85, hearing, 629, 630, for human blood, 2, 6 Surgical shock, heart rate in, 186 (footnote), for incompatibility of Suspensions, 16 Suspensords, 17 blood, 85 Suspensory ligament of lens, 575, 591, Test meal, 313 592, 595 Test sera, 78 Swallowing 301, 302 Testis(es), 655 action of gonadotrophic Sweating, 378, and labyrinth, 634 hormones on 453, 454, action of pitu-Swift, Dean, 211 (footnote) stary on, 455, excision of, see Castra Sympathetic resonance, 220, 424, 425, tion, hormone, see Testosterone, hy-aluronidase in, 5, secondary sex char Sympathetic nervous system, 563, 564, acters and, 452 565, see also under Nerve Testosterone, 451, 452, 656 Tetanus bacilli, 212, 214 Sympathin 425 Sympathoadrenal system, 566 Tetanus of muscle, 488, 489 Tetany, 414ff, due to alkalosis, 417, in-Sympathomimetic action of adrenaline,

fantile 416, latent, 417, parathyroid,

414, 415

Synapse, 510, 511, 512, at autonomic

Tetroses, 367 TH. see Hormone, thyroid Thalamus, 537, 538, 539, 540, 543 Theca, interna and externa of Graafian follicles, 649 Theelin, 449 (footnote) Theobromine, diuretic effect of, 282 Theophyllin, diuretic effect of, 282 Theory(ses) of color vision, 600, du plicity of vision, 579-82, emergency of adrenal function, 423, frequency, of hearing, see telephone, Helmholtz, of pitch, 624, membrane of nerve con duction, 472, place, of hearing (pitch), 626, 627, resonance, of hearing (pitch), 624, 625, Rutherfords, 624, see also telephone, of hearing, of sleep, 572, telephone, of hearing, 625, 626, volley of hearing, 626, Young's, see of

color vision Thiamin 384, 386, 387, 396 (Table), deficiency of, 384, 386 387, 388, 396, and gastrointestinal tone, 386

Thirst, 349 Thoracic duct, 99

Tetrapeptids, 324

Thoracolumbar outflow, 563 Thorax enlargement of in inspiration,

235, expansion of at birth, 232 Threonine, 360

Threshold substances, 277 Thrombin, 103

Thrombocytes, 57, 93, 94, in coagulation of blood 102, reduction, a cause of purpura, 108

Thromboplastin, 103, 105

Thrombosis of coronary arteries, 205 Thymus, 457, 458

Thyroglobulin 412 Thyroid cartilage, 265, 301, 405

Thyroid gland, 404ff adenoma of, 408.

calorigenic action of 380, enlargement of, see Goster; from entoderm func tions of, 412ff, isotopic iodine and, 6 Thyroidectomy, 381, 405, 406, 407, 411,

effect of on heat production 381

Thyrotoxicosis 411, 412

Thyrotrophic hormone and exophthal mos, 411 (footnote), see also under Hormone Thyroxine, 404 412 413

Tidal volume, 240, 241 Tigroid substance, 507, 508 Timbre, of voice 266 Time marker, 484

Tissue(s), 29ff, adipose, 46 47-water of, 55, collagenous 426 427, connective, 44ff-water of 55, delivery of oxygen to 249, development of, from

germ layers, 668, 669, endothelial, 38, epithelial, 29ff-cuboidal, 32, squa mous, 31, 32, stratified, 32, 36 transi tional, 32, erectile of penis, 659, glan dula, 38, infection of, 213, 214, junctional of heart, 165, 166 167, lymph oid, 101, muscular, 40ff, see also Muscle, striated, smooth cardiac nervous, 506 507, 508-water of, 55, nodal, of heart, 165, 166 167, 169, osseous, 48, 49, see also Bone, pul monary, blood supply of, 206 (foot note), renal, see kidney, stimulation of oxidation in by thyroid hormone 405, 413, thyroid, 405, types of, 29ff Tissue fluids, 53, 54, 55, see also under

Fluid Tochopherols 395

Tomatoes vitamin C in, 391 Tone of cardiac nerves 188 189 of

skeletal muscle, 521-23-in paralysis, 527, in spinal shock, 522 and vestibu lar nuclei, 522 of smooth muscle, 293 --postural 293, 294 of vascular nerves 196

Tongue, distribution of taste sensation over, 639 640, papillae of, 638 639, touch spots on, 645, 647

Tonometer, 250

Touch, see under Sensation, cutaneous Toxin 215

Toxoids 219 Toxophore group, 216 Trace elements 402

Trachea, 226, 227, 228 301 Tract(s)

act(s) alimentary, see Alimentary tract, of Burdach, 528, cerebelloves tibular, 553, corticospinal, 526 527 528, 542, 546 554-in new born, 679, decussation of, 546, digestive, see Ali mentary tract, frontopontine 543 553, of Golf 528, hypothalamicohypophyseal, 430 541, optic, 544, 589. of posterior spinal columns, 528, soli tary, 549, of spinal cord, 526, 527, 528, spinocerebellar, 528, 553, 554, tectospinal, 545, temporopontine, 553,

vestibulocerebellar, 553, vestibulospi nal, 528 553, 632 Tractus solitarius see Tract, solitary Transfusion 77ff, indications for, 80, isinglass in 84, with materials of non

human origin, 83, of Rh positive blood, 81, of serum or plasma, 82, in shock, 223, specifications for material used 83

Traumatic shock, see under Shock Tremor in hyperthyroidism, 411, in paralysis agitans, 541

Treppe: in cardiac muscle, 160, in muscular contraction, 490 Tricalcium phosphate in bone, 48 Tricuspid valves of heart, see under

Triceps jerk, 519

Valves Triglycerides, 372 Trigone, 286

Triolein 354 Trioses 367 Tripalmitin 372

Tripeptids, 361 Tristearin, 324, 372

Trophic versus tropic as suffix, 435 (footnate)

Trophobiast 667, 669 Trousseau's sign in tetany, 417

True skin, see Dermis Truncus arteriosus, see Bulbus arteriosus

Trypanosomes, 212 213 Trypsin 288, 292, 323, 324

Trypsinogen, 324 Tryptophane, 360

Tube(s) auditory, 623, 629 Eustachian see auditory, Fallopian see uterine, uterine, 449, 650-fertilization of

ovum in 669 Tuber cinereum, 541

Tubercle, 215

Tubercie bacilii, 212, 213 Tubule of kidney, see renal, renal 269 270, 271, 277, 279-concentrating

function of, 279 reabsorption from 279, 280, seminiferous, 655, 656 Tunica adventitia, 123

Tunica interna, 123 Tunica media, 123

Tuning fork, as time marker, 484, 485 Turbinate bones, 226, 227

Twins, 664 Two point discrimination, see Spatial dis

crimination Tympanum, 617, 618

Tyndall's phenomenon, 17 Typhoid bacillus, 212, 213, 215, 218 Typhoid fever, 215

Tyrosine, 360 tyramine from, 342 Tyrothmem, 222

Umbilical cord, 670, 675, 676 677, 678 Umbilical vein see under Vein

Umbilious 673 Unicellular organisms 23 52, respira-

tion *in*, 225 Unit, of heat see Calorio Universal donor, 79

Uranium 3

Urea, 363, diuretic effect of, 282, in muscle, 497, retention of in blood in nephritis, 284, in urine, 274 Ureter, 285

Urethane action of on capillanes, 203 Urethra, 285, 658 Uric acid, in urine, 274

Urinary bladder, 285, 286 action of adrenaline on, 421, nerves to, 563, postural tone of, 293

Urine composition of, 274, excretion of raw egg white in, 343 (footnote), of fetus, 679, formation of, 276ff, glucose in, 280, 370, 371, loss of heat in 378, 379, of new born 679, pH of, see reaction, phosphorus of 283, pigment of, 274, of pregnancy, 450, 457 (footnote), protein in 284, reaction of, 275, 280, specific gravity of, 275, 284 ultrafiltration and, 278, vitamin C in 280, voiding of, 286, volume of, 275, 276 280 283

Urobilin, 330, 331, in urine in obstruc tive jaundice 335

Urobilinogen 330 Urochrome 274

Uterus, 650, 651, action of adrenaline on, 421, action of pituitary on 437, 455, development of, 651, 652, growth of in pregnancy, 651, 652, involution of, 680 in labor, 676ff, mucosa of

see Endometrium Utricle, 630, 631, 634, in seasickness, 638

Vaccination, 215, 218 (footnote)

Vaccines 218 Vacuoles 22, 24 25

Vagina, 651

Vagustoffe, 192 Valve(s) of heart 116, 118 174 175, 177-aortic, 150, auriculoventricular, 175, bicuspid, see mitral disease of 176, 177, incompetence, 177, mitral, 175, pulmonary, 175, semilunar, see sortic and pulmonary, stenosis of, 177, tricuspid, 175, of Heister, 333, ileo coinc, 321, 339, of veins 125 Valvular desease, 176, 177

Vasoconstriction, caused by noradren aline, 423

Vasodilation caused by adrenaline 423, in sleep, 572

Valves of Heister, see under Valve Vas deferens, 655, 656

Vater ampulla of, 322 Vegetables, laxative action of 346 Vein(s), 126 blood pressure in 151.

cardiae, 204, central of liver, 208, coats of, 123, common iliac, 122, hepatic, 208, interlobular of liver, 208, jugular, 122-pulse in, see Pulse, venous, of liver, 208, pulmonary, 206, renal, 122, 270, 272, structure of, 123, 125, subclavian, 122, 123, of Thebesius 204, umbilical, 672, valves of.

125 Vena cava (venae cavae) in fetus 673 inferior, 118, 122. 167-of cold blooded heart, 166, superior, 118, 122, 167-of cold blooded heart, 166

Venous blood pressure 151, measure ment of, 152, and rate of heart, 190

Venous pulse, see under Pulse Ventricle(s) of brain-fourth 543, 544

554, 558, 559, third 531 537 541. 558 of heart, 115, 116 118, 122 Venule(s), 202

Veratrine causing contracture of mus

cle, 492 Vermiform appendix 321

Vermis of cerebellum 551 552

Vertigo 634 Vessel(s) blood see Arteries Capillar Arterioles Veins lymph, see

Lymphatics Ventricle cerebral 557-59

Vestibule of ear, 630 Vibrations, sound 622

Villus(i) arachnoid, see Arachnoid granulations chorionic 667, 669, in testinal 326 344 345 Viosterol, 393

Virgin birth see Parthogenesis

Virilism, 429 Virus(es) 210 211

Vis a fronte 153 Vis a tergo 153

Vis medicatrix naturae 219

Visceral arches, see Branchial arches 574ff, acuity of, see Visual acuity, binocular and monocular, 586 of birds color, see Color vision, con-

trast effects in, 602 603 604, daylight, see Photopic, defective, 585, duplicity theory of, 579, extrafoveal, 579 in moonlight, 580 (footnote), peripheral, 579, photopic, 580, scotopic, 580, stereoscopic, 597-99, twilight, see scotopic

Visual actity, 584, 585, 615 Visual area see under Area Visual fields, 586, 587

Visual pathway, 589 Visual purple, 582

Vital capacity, 241, 242 Vital red in measurement of blood volume, 63

Vital spirit, 121

Vitamin(s), 384ff, A, 384-86, 396-deficiency of, 385-86, and regeneration of visual purple, 385, 583, antihemorrhagic, see vitamin K, antineuritic, see Thiamin, antirachitic, see vitamin D, antiscorbutic, see vitamin C, antiste rility, see vitamin E, antixerophthal mic, see vitamin A, B complex, 384 386-91-B, factor of, see Thiamin, B. factor of, see Riboflavin, B. factor of, see Pyridoxin, B, factor of, 74, 75, 384, and enzyme systems, 391, C. 373, 391, 392, 396-in adrenal cortex 328, classification of, 384, 386 (Ta ble), D, 392, 393 396-in skin 37 and ultraviolet light 392, deficiencies of see individual vitamins, dietary re quirements of, 399 see also individ ual vitamins, E, 384, 395 396 G see Riboflavin K, 384 396 397-defi ciency in new born 679, and prothrombin, 104 synthesis of by bacteria, 397, in milk 654 synthesis of by bacteria 212, 391, 397

Vitellin 361 Vitreous body hyaluronidase 51 see also under Body

Vocal cords, 265 266 Vocalization center for, 534

Voice, 264ff Vomiting, 318 319, in angina pectoris 205 and labyrinth 634

Waller, A D . 462 Wallerian degeneration 462 "Warming up" of muscle 493

Water absorption of from gall bladder 332, absorption of from intestine, 321 341, 343, balance, 55 of body, 54 diuretic effect of, 282, 283, evapora tion from body, 377, 378 excretion of in nephritis, 284, extracellular, 54, in take of, 55, intracellular, 54, iodine in, 406, metabolism of-adrenal cortex and, 425, 426, 427, hypothalamus and, 542, of metabolism 225 in mus cle, 497, output of, 55 from oxidation of hydrogen, 351, produced by oxida tion of hydrogen, 55, reabsorption of from renal tubule, 277, 280, of urine,

Water vapor, in expired air, 55, 246 Wave(s) of arternal pulse, 158 of electrocardiogram, 180, of electroenceph

276, of various tissues, 55

alogram 537 538 of electroretinogram 585 sound 622 624-fre quency of and pitch 674 of venous pulse 187

"Wear and tear" of body protein 363 Weiner A S 80

Wever and Bray effect, 627 Wharton's jelly 51 Wheat prote as of 365 whole laxative

act on of 346 Wheat germ 395 396

Whey protein 309 Whipple G H 74 White matter of cerebellum 552 553

of pervous system 506 of spinal cord 523 524 Window oval 622 623 round 622

623 Window glass opaque to ultraviolet rays

"Witch's m lk " 456 (footnote)

X chromosome 665 Xanthophores, 439 Xerophthalmia 385 386 396

Y chromosome, 665 Yawning 259 Yeast enzymes of 290 irrad at on of 393

Yellow body see Corpus luteum Yellow spot see Macula lutea Yolk sac 667 670 Youngs theory of color vis on see un der Theory

Z l ne 41 47 43 Zein 361 365 Z ne essent al in det 40° with insulin 371 402

Zona granulosa see Membrana granu losa pellucida 649 Zygote segmentat on of 633 664 666 Zymogens, 25